

PHYSICIANS CARE ASPIRIN- aspirin tablet, coated
PHYSICIANS CARE ASPIRIN- aspirin tablet, film coated
Acme United Corporation

Physicians Care Aspirin

Drug Facts

Active ingredient (in each tablet)

Aspirin (NSAID*) 325 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

Temporarily relieves:

- headache
- minor arthritis pain
- pain & fever of colds
- backache
- menstrual cramps
- toothache
- muscle aches

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcohol drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or to any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have asthma
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking prescription drugs for gout, diabetes or arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- if ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms appear
- an allergic reaction occurs. Seek medical help right away.

If pregnant or breast-feeding,ask a health professional before use. **It is especially important not to use aspirin at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose

Adults and children 12 years and over:

take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in

24 hours

Children under 12 years:

consult a doctor

Other information

- read all product information before using
- store at room temperature
- avoid excessive heat and humidity
- do not use any opened or torn packets

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments? 1-800-835-2263

Physicians Care Aspirin Label

90014

Physicians Care®

Aspirin

Pain Reliever and fever Reducer (NSAID)

Pull To Open

This Package for Households without Young Children.

Coated Tablets

Tamper-Evident Packets of 2 Tablets

Do not use if packet is open or torn.

Aspirin (NSAID) 325 mg

100 Tablets (50 Packets, 2 tablets each)



Also Available from PhysiciansCare®
in individual dose packets:

- PhysiciansCare® Non-Aspirin (Acetaminophen)
- PhysiciansCare® Ibuprofen
- PhysiciansCare® Cold & Cough
- PhysiciansCare® Allergy
- PhysiciansCare® Sinus
- PhysiciansCare® Antacid

PhysiciansCare®

ASPIRIN

Pain Reliever and Fever Reducer (NSAID)

THIS PACKAGE FOR HOUSEHOLDS
WITHOUT YOUNG CHILDREN.

COATED TABLETS

Tamper-Evident Packets of 2 Tablets
Do not use if packet is open or torn.

Aspirin (NSAID) 325 mg
100 Tablets (50 Packets, 2 tablets each)

90014

Retain carton for complete product information

Drug Facts

Active ingredient (in each tablet) Aspirin (NSAID) 325 mg

Purpose Pain reliever / fever reducer / nonsteroidal anti-inflammatory drug

Uses
Temporarily relieve:
• headache • minor arthritis pain • pain & fever of colds
• backache • menstrual cramps • toothache

Warnings
Roy's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Roy's syndrome, a rare but serious illness.

Allergy alert: aspirin may cause a severe allergic reaction which may include:
• hives • facial swelling • shock • asthma (wheezing)
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
• are age 60 or older
• have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
• have 3 or more alcohol drinks every day while using this product
• take more or for a longer time than directed

Do not use
• if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if
• stomach bleeding warning applies to you
• you have asthma
• you have a history of stomach problems such as heartburn
• you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
• you are taking a diuretic

Ask a doctor or pharmacist before use if you are
• taking a prescription drug for • gout • diabetes • arthritis

Contact Us 1-800-835-2263 www.FirstAidOnly.com

Drug Facts (continued)

Stop use and ask a doctor if
• you experience any of the following signs of stomach bleeding:
• feel faint
• vomit blood
• have bloody or black stools
• have stomach pain that does not get better
• pain gets worse or lasts more than 3 days
• fever gets worse or lasts more than 3 days
• if ringing in the ears or loss of hearing occurs
• redness or swelling is present in the painful area
• any new symptoms appear
• an allergic reaction occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
• drink a full glass of water with each dose

Adults and children 12 years and over: take 1 or 2 tablets every 4 hours not to exceed 12 tablets in 24 hours

Children under 12 years: consult a doctor

Other information
• read all product information before using
• store at room temperature
• avoid excessive heat and humidity
• do not use any opened or torn packets

Inactive ingredients
corn starch, hydroxypropyl methylcellulose, polyethylene glycol, polyethylene glycol, polyethylene glycol, "may contain"

Questions? 1-800-835-2263

90014

PhysiciansCare®

ASPIRIN

Pain Reliever and Fever Reducer (NSAID)


THIS PACKAGE FOR HOUSEHOLDS
WITHOUT YOUNG CHILDREN.

COATED TABLETS

Tamper-Evident Packets of 2 Tablets
Do not use if packet is open or torn.

Aspirin (NSAID) 325 mg
100 Tablets (50 Packets, 2 tablets each)

90014



Also Available from PhysiciansCare®
in individual dose packets:

- PhysiciansCare® Non-Aspirin (Acetaminophen)
- PhysiciansCare® Ibuprofen
- PhysiciansCare® Cold & Cough
- PhysiciansCare® Allergy
- PhysiciansCare® Sinus
- PhysiciansCare® Antacid

PhysiciansCare®

ASPIRIN

Pain Reliever and Fever Reducer (NSAID)

COATED CAPLET

Individually packaged for workplace use

Tamper-Evident Packets of 2 Tablets
Do not use if packet is open or torn.

Aspirin (NSAID) 325 mg
100 Tablets (50 Packets, 2 tablets each)

90014

Retain carton for complete product information

Drug Facts

Active ingredient (in each tablet) Aspirin (NSAID) 325 mg

Purpose Pain reliever / fever reducer / nonsteroidal anti-inflammatory drug

Uses
Temporarily relieve:
• headache • minor arthritis pain • pain & fever of colds
• cramps with pain • toothache • aches with pain

Warnings
Roy's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Roy's syndrome, a rare but serious illness.

Allergy alert: aspirin may cause a severe allergic reaction which may include:
• hives • facial swelling • shock • asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
• are age 60 or older
• have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
• have 3 or more alcohol drinks every day while using this product
• take more or for a longer time than directed

Do not use
• if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if
• stomach bleeding warning applies to you
• you have asthma
• you have a history of stomach problems such as heartburn
• you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
• you are taking a diuretic

Ask a doctor or pharmacist before use if you are
• taking a prescription drug for • gout • diabetes • arthritis

Contact Us 1-800-835-2263 www.FirstAidOnly.com

Drug Facts (continued)

Stop use and ask a doctor if
• you experience any of the following signs of stomach bleeding:
• feel faint
• vomit blood
• have bloody or black stools
• have stomach pain that does not get better
• pain gets worse or lasts more than 3 days
• fever gets worse or lasts more than 3 days
• if ringing in the ears or loss of hearing occurs
• redness or swelling is present in the painful area
• any new symptoms appear
• an allergic reaction occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
• drink a full glass of water with each dose

Adults and children 12 years and over: take 1 or 2 tablets every 4 hours not to exceed 12 tablets in 24 hours

Children under 12 years: consult a doctor

Other information
• read all product information before using
• store at room temperature
• avoid excessive heat and humidity
• tamper evident sealed packets
• do not use any opened or torn packets

Inactive ingredients
corn starch, hydroxypropyl methylcellulose, polyethylene glycol, polyethylene glycol

Questions? 1-800-835-2263

90014

PhysiciansCare®

ASPIRIN

Pain Reliever and Fever Reducer (NSAID)

THIS PACKAGE FOR HOUSEHOLDS
WITHOUT YOUNG CHILDREN.

COATED CAPLET

Tamper-Evident Packets of 2 Tablets
Do not use if packet is open or torn.

Aspirin (NSAID) 325 mg
100 Tablets (50 Packets, 2 tablets each)

90014



© 2022 Bayer HealthCare Corporation
All rights reserved. Bayer is a registered trademark of Bayer AG.

04941212

90014
Physicians Care®
Aspirin
Pain Reliever and fever Reducer (NSAID)
Pull To Open
This Package for Households without Young Children.
Coated Tablets
Tamper-Evident Packets of 2 Tablets
Do not use if packet is open or torn.

Aspirin (NSAID) 325 mg

100 Tablets (50 Packets, 2 tablets each)

PHYSICIANS CARE ASPIRIN

aspirin tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0924-0106
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POVIDONE (UNII: FZ989GH94E)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	FR21
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0924-0106-01	50 in 1 BOX	04/01/2021	05/31/2023
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:0924-0106-00	2 in 1 PACKET; Type 0: Not a Combination Product	04/01/2021	05/31/2023
3	NDC:0924-0106-02	125 in 1 BOX	04/01/2021	05/31/2023
3		2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:0924-0106-03	10 in 1 BOX	04/01/2021	05/31/2023
4		2 in 1 PACKET; Type 0: Not a Combination Product		
5	NDC:0924-0106-07	25 in 1 BOX	04/01/2021	05/31/2023



Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/01/2021	05/31/2023

PHYSICIANS CARE ASPIRIN

aspirin tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0924-0105
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	Aspirin;44;157
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0924-0105-01	50 in 1 BOX	07/20/2020	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:0924-0105-00	2 in 1 PACKET; Type 0: Not a Combination Product	07/20/2020	
3	NDC:0924-0105-02	125 in 1 BOX	07/20/2020	
3		2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:0924-0105-03	10 in 1 BOX	07/20/2020	
4		2 in 1 PACKET; Type 0: Not a Combination Product		
5	NDC:0924-0105-07	25 in 1 BOX	07/20/2020	
5		2 in 1 PACKET; Type 0: Not a Combination Product		
6	NDC:0924-0105-04	6 in 1 CARTON	07/20/2020	
6	NDC:0924-0105-00	2 in 1 PACKET; Type 0: Not a Combination Product		
7	NDC:0924-0105-05	20 in 1 CARTON	07/20/2020	
7	NDC:0924-0105-00	2 in 1 PACKET; Type 0: Not a Combination Product		
8	NDC:0924-0105-06	250 in 1 CARTON	07/20/2020	
8	NDC:0924-0105-00	2 in 1 PACKET; Type 0: Not a Combination Product		


Also Available from PhysiciansCare®
in individual dose packets:

- PhysiciansCare® Non-Aspirin (Acetaminophen)
- PhysiciansCare® Ibuprofen
- PhysiciansCare® Cold & Cough
- PhysiciansCare® Allergy
- PhysiciansCare® Sinus
- PhysiciansCare® Antacid

PhysiciansCARE®

ASPIRIN

Pain Reliever and Fever Reducer (NSAID)



Tamper-Evident packets of 2 Tablets
Do not use if packet is open or torn.

Aspirin (NSAID) 325 mg
100 Tablets (50 Packets, 2 Tablets each)



90014

Retain carton for complete product information

Drug Facts
Active ingredient (in each tablet): Aspirin (NSAID), 325 mg. **Purpose:** Pain reliever / fever reducer / nonsteroidal anti-inflammatory drug.

Uses: Temporarily relieves: minor aches/pain, pain & fever of cold/flu, cramps with pain, headache, aches with pain.

Warnings: **Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, it changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** aspirin may cause a severe allergic reaction which may include: hives, facial swelling, shock, asthma (wheezing).

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 55 or older; have had stomach ulcers or bleeding problems; take a blood thinning (anticoagulant) or blood drug; take other drugs containing prescription or nonprescription NSAIDs (ibuprofen, naproxen, or others); have 3 or more alcohol drinks every day while using this product; take more or for a longer time than directed.

Do not use: if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if: stomach bleeding warning applies to you; you have asthma; you have a history of stomach problems such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, or kidney disease; you are taking a diuretic.

Ask a doctor or pharmacist before use if you are taking a prescription drug for: gout, diabetes, arthritis.

Stop use and ask a doctor if: you experience any of the following signs of stomach bleeding: feel faint; vomit blood; have bloody or black stools; have stomach pain that does not get better; stool gets worse or lasts more than 3 days; if ringing in the ears or loss of hearing occurs; redness or swelling is present in the painful area. Any new symptoms appear.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 30 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions: Drink a full glass of water with each dose.

Adults and children 12 years and over: Take 1 or 2 tablets every 4 hours, not to exceed 12 tablets in 24 hours.

Children under 12 years: consult a doctor.

Other information: Read all product information before using; store at room temperature; avoid excessive heat and humidity; use tamper-evident sealed packets; do not use any opened or torn packets.

Inactive ingredients: corn starch, hydroxyethylcellulose, polyethylene glycol, croscarmellose sodium.

Questions? 1-800-835-2263


Contact Us 1-800-835-2263 www.FirstAidOnly.com

90014

PhysiciansCARE®

ASPIRIN

Pain Reliever and Fever Reducer (NSAID)



Tamper-Evident packets of 2 Tablets
Do not use if packet is open or torn.

Aspirin (NSAID) 325 mg
100 Tablets (50 Packets, 2 Tablets each)

© 2023 Acme United Corporation
www.AcmeUnited.com
Manufactured by:
Unifirst First Aid Corporation
19900, CT 06488

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/20/2020	

Labeler - Acme United Corporation (001180207)

Registrant - Unifirst First Aid Corporation (832947092)

Revised: 12/2024

Acme United Corporation