

**COMPLETE LICE TREATMENT KIT- piperonyl butoxide, pyrethrum extract**  
**Walgreen Company**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Walgreen Co. Complete Lice Treatment Kit Drug Facts**

**Active ingredient**

Piperonyl butoxide 4%

Pyrethrum extract (equivalent to 0.33% pyrethrins)

**Purpose**

Lice treatment

**Uses**

treats head, pubic (crab), and body lice

**Warnings**

For external use only

**Do not use**

- near eyes
- inside nose, mouth, or vagina
- on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

**Ask a doctor before use if you are**

allergic to ragweed. May cause breathing difficulty or an asthmatic attack.

**When using this product**

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets into eyes, flush with water right away
- scalp itching or redness may occur

**Stop use and ask a doctor if**

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

Shake well before using

- Important: Read warnings before use

Adults and children 2 years and over:

#### Inspect

- check each household member with a magnifying glass in bright light for lice/nits (eggs)
- look for tiny nits near scalp, beginning at back of neck and behind ears
- examine small sections of hair at a time
- unlike dandruff which moves when touched, nits stick to the hair
- if either lice or nits are found, treat with this product

#### Treat

- apply thoroughly to DRY HAIR or other affected area. For head lice, first apply behind ears and to back of neck.
- allow product to remain for 10 minutes, but no longer
- use warm water to form a lather, shampoo, then thoroughly rinse
- for head lice, towel dry hair and comb out tangles

#### Remove lice and their eggs (nits)

- use a fine-tooth or special lice/nit comb. Remove any remaining nits by hand (using a throw-away glove).
- hair should remain slightly damp while removing nits
- if hair dries during combing, dampen slightly with water
- for head lice, part hair into sections. Do one section at a time starting on top of the head. Longer hair may take 1 to 2 hours.
- lift a 1- to 2-inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp.
- pin back each strand of hair after combing
- clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back.
- after combing, thoroughly recheck for lice/nits. Repeat combing if necessary.
- check daily for any lice/nits that you missed
- a second treatment must be done in 7 to 10 days to kill any newly hatched lice
- if infestation continues, see a doctor for other treatments
- Children under 2 years: ask a doctor

#### Other information

- store at 20°-25°C (68°-77°F)
- keep carton for important product information
- see package insert for additional information

#### Inactive ingredients

benzyl alcohol, C13-C14 isoparaffin, fragrance, polyoxyethylene (1,1,3,3-tetramethyl butyl) phenyl ether, water, xanthan gum

#### Questions or comments?

1-800-719-9260

#### Consumer Information

**LICE KILLING SHAMPOO**

## **For the treatment of Head Lice, Pubic (Crab) Lice and Body Lice**

### **SHAKE WELL BEFORE USING**

### **DIRECTIONS: IMPORTANT: READ WARNINGS BEFORE USING. A SECOND TREATMENT MUST BE DONE IN 7-10 DAYS TO KILL ANY NEWLY HATCHED LICE**

**Inspect** • Check each household member with a magnifying glass in bright light for lice/nits (eggs) • Look for tiny nits near scalp, beginning at back of neck and behind ears • Examine small sections of hair at a time • Unlike dandruff which moves when touched, nits stick to the hair • If either lice or nits are found, treat with this product

**Treat** • Apply thoroughly to dry hair or other affected area. For head lice, first apply behind ears and to back of neck • Allow product to remain for 10 minutes but no longer • Use warm water to form a lather, shampoo, then thoroughly rinse • For head lice, towel dry hair and comb out tangles

**Remove Lice and their eggs (nits)** • Use a fine-tooth or special lice/nit comb. Remove any remaining nits by hand (using a throw-away glove) • Hair should remain slightly damp while removing nits • If hair dries during combing, dampen slightly with water • For head lice, part hair into sections. Do one section at a time starting on top of the head. Longer hair may take 1 to 2 hours • Lift a 1-2 inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp • Pin back each strand of hair after combing • Clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back • After combing, thoroughly recheck for lice/nits. Repeat combing if necessary • Check daily for any lice/nits that you missed • A second treatment must be done in 7 to 10 days to kill any newly hatched lice • If infestation continues, see a doctor for other treatments

1. **HEAD LICE:** • Head lice lay small white eggs (nits) on the hair shaft close to the scalp • Nits are most easily found on the back of the neck or behind ears • Disinfect hats, hair ribbons, scarves, coats, towels and bed linens by machine washing in hot water (above 54° C (130° F)), then using hottest dryer cycle for at least 20 minutes • Items that cannot be washed (bedspreads, blankets, pillows, stuffed toys, etc.) should be dry-cleaned or sealed in a plastic bag for 4 weeks, then removed outdoors and shaken out very hard before using again • Items that cannot be washed, dry-cleaned, or stored may be sprayed with a product designed for this purpose • Soak all combs and brushes in hot water (above 54° C (130° F)) for at least 10 minutes • Vacuum all carpets, mattresses, upholstered furniture, and car seats that may have been used by affected people

2. **PUBIC (CRAB) LICE:** • Pubic lice may be transmitted by sexual contact. • Sexual partners should be treated simultaneously to avoid reinfestation • Lice are very small and look like brown or grey dots on the skin • Pubic lice usually cause intense itching and lay small white eggs (nits) on the hair shaft generally close to the skin surface • Pubic lice may be present on the short hairs of the groin, thighs, trunk and underarms, and occasionally on the beard and mustache • Disinfect underwear by machine washing in hot water (above 54° C (130° F)), then using hottest dryer cycle for at least 20 minutes

3. **BODY LICE:** • Body lice and their eggs (nits) are generally found in the seams of clothing, particularly in the waistline and armpit area • Body lice feed on skin then return to clothing to lay their eggs • Disinfect clothing by machine washing in hot water (above 54° C (130° F)), then using hottest dryer cycle for at least 20 minutes • Do not seal clothing in a plastic bag because nits can remain dormant for up to 30 days

**Save this insert for future reference.**

Packaged by Perrigo, Allegan, MI 49010 USA

### **CHAMPÚ PARA ELIMINAR PIOJOS**

**Para el tratamiento de piojos de la cabeza, piojos del pubis (ladillas) y piojos del cuerpo**

**AGITE MUY BIEN EL PRODUCTO ANTES DE USARLO**

**INSTRUCCIONES: ES IMPORTANTE QUE LEA LAS ADVERTENCIAS ANTES DE USAR**

## **EL PRODUCTO. DEBE REALIZAR UN SEGUNDO TRATAMIENTO 7 A 10 DÍAS DESPUÉS PARA ELIMINAR LA REPRODUCCIÓN DE NUEVOS PIOJOS**

**Inspección** • Revise a cada miembro de su familia con una lupa, a la luz del día, para encontrar piojos o huevecillos (liendres) • Busque las diminutas liendres cerca del cuero cabelludo, empezando por la nuca y detrás de las orejas • Examine pequeñas secciones de cabello una por una • Las liendres se adhieren al cuero cabelludo, lo cual las diferencia de la caspa, que se suelta al contacto • Si encuentra piojos o liendres, trátelos con este producto

**Tratamiento** • Aplique sobre todo el cabello seco u otras áreas afectadas. Para piojos de la cabeza, primero aplique detrás de las orejas y en la nuca • Deje el producto aplicado durante 10 minutos, sin exceder el tiempo sugerido • Use agua tibia para formar espuma, lave con champú y luego enjuague muy bien • para piojos de la cabeza, seque el cabello con una toalla y desenrede con un peine

**Remueva los piojos y sus liendres (huevecillos)** • Use un peine fino o un peine especial para piojos/liendres. Remueva todas las liendres restantes a mano (use guantes desechables) • El cabello debe permanecer apenas húmedo mientras remueve las liendres • Si el cabello se seca mientras lo peina, humidézcalo apenas con agua • Para piojos de la cabeza, divida el cabello en secciones. Revise una sección por vez, empezando por la parte de arriba de la cabeza. Si tiene el cabello largo, esto le puede llevar de 1 a 2 horas • Levante mechones de 2.5 a 5 cm (1 a 2 pulgadas) de anchura. Ponga el peine lo más cerca posible del cuero cabelludo y peine con movimientos firmes y uniformes • Sujete cada mechón de cabello después de peinarlo • Limpie el peine con frecuencia. Quite las liendres con un pañuelo de papel y arrójelas en una bolsa plástica. Cierre la bolsa y deséchela para evitar que los piojos vuelvan a aparecer • Después de peinar, vuelva a revisar exhaustivamente para ver si encuentra piojos o liendres. Si fuera necesario, repita el procedimiento • Revise a diario para ver si aparece algún piojo o liendre • Debe realizar un segundo tratamiento después de 7 a 10 días para matar piojos recién nacidos • Si la infestación continúa, debe consultar a su médico acerca de otros tratamientos

### **PACKAGE LEAFLET – SPANISH CONTINUED**

**1. PIOJOS DE LA CABEZA:** • Los piojos de la cabeza ponen pequeños huevecillos blancos (liendres) en los mechones de cabello, cerca del cuero cabelludo • Las liendres son fáciles de encontrar en la nuca o detrás de las orejas • Desinfecte sombreros, moños, bufandas, abrigos, toallas y sábanas; lávelos en la lavadora con agua caliente (a más de 54°C (130°F)); después use el ciclo de secado más caliente al menos durante 20 minutos • Los artículos que no se puedan lavar (cubrecamas, frazadas, almohadas, juguetes de peluche, etc.) se deben limpiar en seco o guardar en bolsas plásticas por cuatro semanas, después se los debe sacar a la intemperie y sacudir enérgicamente antes de ser usados otra vez • Los artículos que no se puedan lavar, limpiar en seco o guardar se deben rociar con productos designados para este fin • Sumerja todos los peines y cepillos en agua caliente (a más de 54°C (130°F)) al menos durante 10 minutos • Aspire las alfombras, colchones, muebles tapizados y asientos del automóvil que puedan haber sido usados por las personas afectadas

**2. PIOJOS DEL PUBIS (LADILLAS):** • Los piojos del pubis se pueden transmitir por contacto sexual • Los compañeros sexuales se deben recibir tratamiento simultáneo para evitar la reinfestación • Los piojos son muy pequeños y parecen puntos de color café o gris en la piel • Los piojos del pubis producen picazón muy intensa y ponen huevecillos muy pequeños (liendres) en los mechones de pelo, generalmente cerca de la superficie de la piel • Los piojos del pubis pueden estar presentes en los pelos cortos de la ingle, muslos, tronco, axilas y, ocasionalmente, en la barba y el bigote. Lave la ropa interior en la lavadora con agua caliente (a más de 54°C (130°F)) para desinfectarla y después use el ciclo de secado más caliente durante más de 20 minutos

**3. PIOJOS DEL CUERPO:** • Los piojos del cuerpo y sus huevos (liendres) generalmente se encuentran en las costuras de la ropa, particularmente en el elástico de la cintura o en la zona de la axila • Los piojos del cuerpo se alimentan de la piel y luego regresan a la ropa para depositar sus huevos • Lave la ropa en la lavadora con agua caliente (a más de 54°C (130°F)) para desinfectarla y después use el ciclo más caliente de la secadora, por lo menos durante 20 minutos • No guarde la ropa en bolsas de plástico, pues las liendres permanecen en estado latente por más de 30 días

**Guarde estas instrucciones para consulta futura.**

Empacado por Perrigo, Allegan, MI 49010 EE. UU.

**Principal Display Panel**

STOP LICE

COMPLETE LICE TREATMENT KIT

1-2-3 LICE CONTROL SYSTEM

STEP 1 LICE KILLING SHAMPOO

Lice Treatment

Works fast to kill lice and eggs

STEP 2 EGG & NIT COMBING GEL

Lice comb removes nits and eggs

Makes combing faster and easier

STEP 3 BEDDING SPRAY LICE TREATMENT

For bedding and furniture

Kills lice and their eggs on mattresses, furniture, car interiors and other non-washable items

Compare to RID® Lice Killing Shampoo active ingredients

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back and side panels for additional information.

Walgreens

**STOP LICE**

# COMPLETE LICE TREATMENT KIT

1-2-3 LICE CONTROL SYSTEM

INSTRUCCIONES EN ESPAÑOL INCLUIDAS

AVISO AL CONSUMIDOR: Si no sabe inglés, pídale a alguien que le lea toda la información que aparece en el empaque antes de utilizar este producto.

NDC 0363-0173-24

Walgreens

**STOP LICE**

# COMPLETE LICE TREATMENT KIT

1-2-3 LICE CONTROL SYSTEM

## STEP 1 LICE KILLING SHAMPOO

- Lice Treatment
- Works fast to kill lice and eggs

## STEP 2 EGG & NIT COMBING GEL

- Lice comb removes nits and eggs
- Makes combing faster and easier

## STEP 3 BEDDING SPRAY LICE TREATMENT

- For bedding and furniture
- Kills lice and their eggs on mattresses, furniture, car interiors and other non-washable items

Compare to RID® Lice Killing Shampoo active ingredients\*\*\*

Spray NET WT 5 OZ (142 g)

Gel NET WT 2 OZ (57 g)

Shampoo 4 FL OZ (118 mL)

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**  
See back and side panels for additional information.

ITEM 553625



### 1-2-3 LICE CONTROL SYSTEM

Complete lice control for your family and home.

- KILL LICE** STEP 1
- Apply Lice Killing Shampoo according to label directions.
  - Repeat this step 7 to 10 days later to help prevent reinfestation.
- COMB OUT EGGS & NITS** STEP 2
- After Step 1, apply Egg & Nit Combing Gel on damp hair to make removal faster and easier.
  - Comb out the eggs and nits in the hair with a fine-toothed comb or with the comb included.
- CLEAN HOME** STEP 3
- Use Lice Bedding Spray to kill lice and their eggs on mattresses, furniture, car interiors and other non-washable items.
  - Wash bed linens, clothing and other items in hot water and dry in high heat.

**IMPORTANT:** Read detailed directions before using

This kit contains 3 products for complete lice control for your family and home. The 4 oz bottle of shampoo and 2 oz bottle of gel should be enough product to treat one person. However, amount of product needed will vary by hair length. See below for guidelines.

| Lice Killing Shampoo          |                                       |
|-------------------------------|---------------------------------------|
| Hair Length                   | Approximate Amount For 1 Adult/Child: |
| Short (ear length or shorter) | First Application: 1 oz - 2 oz        |
|                               | Second Application: 1 oz - 2 oz       |
|                               | <b>Total: 2 oz - 4 oz</b>             |
| Medium (shoulder length)      | First Application: 2 oz - 3 oz        |
|                               | Second Application: 2 oz - 3 oz       |
|                               | <b>Total: 4 oz - 6 oz</b>             |
| Long (past shoulder length)   | First Application: 3 oz - 4 oz        |
|                               | Second Application: 3 oz - 4 oz       |
|                               | <b>Total: 6 oz - 8 oz</b>             |

  

| Egg & Nit Combing Gel                   |                                       |
|---|---------------------------------------|
| Hair Length                             | Approximate Amount For 1 Adult/Child: |
| Short (ear length or shorter)           | First Application: ½ bottle           |
|   | Second Application: ½ bottle          |
|   | <b>Total: 1 Bottle (2 oz)</b>         |
| Medium/Long (shoulder length or longer) | First Application: 1 bottle           |
|   | Second Application: 1 bottle          |
|   | <b>Total: 2 Bottles (4 oz)</b>        |



LOT NO.

EXP.

17324 94 C5

**LICE KILLING SHAMPOO**

**Drug Facts**

**Active Ingredients Purpose**  
 Piperonyl butoxide 4%.....Lice treatment  
 Pyrethrum extract (equivalent to 0.33% pyrethrins).....Lice treatment

**Uses** treats head, pubic (crab), and body lice

**Warnings**  
 For external use only

**Do not use**  
 ■ near eyes ■ inside nose, mouth, or vagina  
 ■ on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

**Ask a doctor before use if you are allergic to ragweed.** May cause breathing difficulty or an asthmatic attack.

**When using this product**  
 ■ keep eyes tightly closed and protect eyes with a washcloth or towel  
 ■ if product gets into eyes, flush with water right away  
 ■ scalp itching or redness may occur

**Stop use and ask a doctor if**  
 ■ breathing difficulty occurs ■ eye irritation occurs  
 ■ skin or scalp irritation continues or infection occurs

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**  
 Shake well before using  
 ■ **Important:** Read warnings before use  
**Adults and children 2 years and over:**  
**Inspect**  
 ■ check each household member with a magnifying glass in bright light for lice/nits (eggs)  
 ■ look for tiny nits near scalp, beginning at back of neck and behind ears  
 ■ examine small sections of hair at a time  
 ■ unlike dandruff which moves when touched, nits stick to the hair  
 ■ if either lice or nits are found, treat with this product  
**Treat**  
 ■ apply thoroughly to **DRY HAIR** or other affected area. For head lice, first apply behind ears and to back of neck.  
 ■ allow product to remain for 10 minutes, but no longer  
 ■ use warm water to form a lather, shampoo, then thoroughly rinse  
 ■ for head lice, towel dry hair and comb out tangles  
**Remove lice and their eggs (nits)**  
 ■ use a fine-tooth or special lice/nit comb. Remove any remaining nits by hand (using a throw-away glove).  
 ■ hair should remain slightly damp while removing nits  
 ■ if hair dries during combing, dampen slightly with water  
 ■ for head lice, part hair into sections. Do one section at a time starting on top of the head. Longer hair may take 1 to 2 hours.  
 ■ lift a 1- to 2-inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp.

**Drug Facts (continued)**

- pin back each strand of hair after combing
- clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back.
- after combing, thoroughly recheck for lice/nits. Repeat combing if necessary.
- check daily for any lice/nits that you missed
- a second treatment must be done in 7 to 10 days to kill any newly hatched lice
- if infestation continues, see a doctor for other treatments
- **Children under 2 years:** ask a doctor

**Other information**  
 ■ store at 20°-25°C (68°-77°F)  
 ■ keep carton for important product information  
 ■ see package insert for additional information

**Inactive ingredients**  
 benzyl alcohol, C13-C14 isoparaffin, fragrance, polyoxyethylene (1, 1, 3, 3-tetramethyl butyl) phenyl ether, water, xanthan gum

**Questions or comments?**  
 1-800-719-9260

**EGG & NIT COMBING GEL**

**Product Facts**

**Use**  
 to make egg and nit removal from the hair faster and easier

**Warnings**  
 For external use only. Keep out of reach of children.

**Directions**  
 ■ before using, read additional information on insert inside carton  
 ■ use AFTER a lice killing shampoo treatment  
 ■ towel dry hair and comb out tangles with regular comb  
 ■ apply this product to one section of damp hair at a time  
 ■ massage well to ensure product covers the entire section  
 ■ pay special attention to the areas behind the ears and nape of the neck where nits and eggs are more likely to be found  
 ■ comb out the dead lice, eggs and nits with the enclosed nit comb (see insert for additional information). **This step is very important.**  
 ■ if hair dries during combing, dampen slightly with water and re-apply gel as needed  
 ■ rinse thoroughly with warm water, after you have combed entire head  
 ■ disinfect combs with hot water (130°F)

**Other information**  
 ■ it is important to wash in hot water (130°F) all clothing, bedding, towels, and hair products (combs, brushes) used by affected persons  
 ■ dry clean non-washable fabrics

**Product Facts (continued)**

- to eliminate infestation of furniture and bedding that cannot be washed or dry cleaned, a multi-use lice spray may be used
- **this product does not kill lice or their eggs.** Use this product only after treating with a lice killing shampoo to kill lice and their eggs.

**Ingredients:**  
 water, glycolic acid, polyquaternium-10, hydroxyethylcellulose, potassium hydroxide, propylene glycol, diazolidinyl urea, methylparaben, propylparaben

Store at 20°-25°C (68°-77°F)

**LICE CONTROL SPRAY**

**Product Facts**

**ACTIVE INGREDIENT:**  
 \*Permethrin..... 0.50%

**\*\*OTHER INGREDIENTS:**..... 99.50%

100.00%

\*Cis/trans ratio: Min. 35% (+/-) cis and Max. 65% (+/-) trans.  
 \*\*Contains Petroleum distillate

**PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS**

**CAUTION:** Harmful if swallowed or absorbed through skin. Avoid inhalation of spray mist. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Avoid contamination of feed and foodstuffs.

**PHYSICAL OR CHEMICAL HAZARDS**  
 Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F (54°C) may cause bursting.

**FIRST AID**  
 Call a Poison Control Center or doctor immediately for treatment advice. Have product container or label with you when calling a Poison Control Center or doctor or going for treatment.

**If swallowed:**  
 ■ Immediately call a Poison Control Center or doctor.  
 ■ Do not induce vomiting unless told to do so by a Poison Control Center or doctor.  
 ■ Do not give any liquid to the person.  
 ■ Do not give anything by mouth to an unconscious person.

**If in eyes:**  
 ■ Hold eye open and rinse slowly and gently with water for 15-20 minutes.  
 ■ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

**If inhaled:**  
 ■ Move person to fresh air.  
 ■ If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

**Product Facts (continued)**

**If on skin or clothing:**  
 ■ Take off contaminated clothing.  
 ■ Rinse skin immediately with plenty of water for 15-20 minutes.

**NOTE TO PHYSICIAN**  
 Contains petroleum distillate - vomiting may cause aspiration pneumonia.

**DIRECTIONS FOR USE**  
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Remove pets and birds and cover fish aquaria before space spraying or surface applications. **This product is not for use on humans.** If lice infestation occurs on humans, use a product labeled for use on humans. Vacate room after treatment and ventilate before reoccupying. Do not allow children or pets to contact treated areas until surfaces are dry.

**SHAKE WELL BEFORE USING**  
 Remove protective cap. Hold container upright and spray from a distance of 8-10 inches. Individuals with asthma or severe allergies should consult their doctor before using this product and have someone else apply this product.

**To kill lice and louse eggs:** Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 8 to 10 inches. Spray each square foot for three seconds. Spray to the point of dampness but not run-off. Spray only those articles and parts of bedding including mattresses and furniture, that cannot be either laundered or dry-cleaned. Do not use on pillowcases, sheets or clothing. Allow all sprayed articles to dry thoroughly before each use.

**STORAGE AND DISPOSAL**  
**Storage:** Store in a cool, dry area away from children and pets and away from heat, sparks and open flame. Do not transport or store below 32°F (0°C).  
**Container disposal:** Do Not Puncture or Incinerate.  
 If empty: Place in trash or offer for recycling if available.  
 If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

**NOTICE: Purchaser assumes all responsibility for safety and use not in accordance with directions.**

EPA Reg. No. 71-1-43428  
 EPA Est. No. See Bottom or Top of Can  
 DOT 2P

\*\*\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of RID® Lice Killing Shampoo.

Distributed by: Walgreen Co.,  
 200 Wilmet Road  
 Deerfield, IL 60015-4616

 100% Satisfaction Guaranteed with all Walgreens Products or Your Money Back. www.walgreens.com

REV 1108

**Complete Lice Treatment Kit Carton Image 2**

piperonyl butoxide, pyrethrum extract kit

### Product Information

|                     |                |                           |               |
|---------------------|----------------|---------------------------|---------------|
| <b>Product Type</b> | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:0363-0173 |
|---------------------|----------------|---------------------------|---------------|

### Packaging

| # | Item Code        | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:0363-0173-24 | 1 in 1 CARTON       |                      |                    |

### Quantity of Parts

| Part # | Package Quantity | Total Product Quantity |
|--------|------------------|------------------------|
| Part 1 | 1 BOTTLE         | 118 mL                 |
| Part 2 | 1 BOTTLE         | 57 g                   |

### Part 1 of 2

#### LICE KILLING

piperonyl butoxide, pyrethrum extract shampoo

### Product Information

|                           |               |
|---------------------------|---------------|
| <b>Item Code (Source)</b> | NDC:0363-0866 |
|---------------------------|---------------|

|                                |         |
|--------------------------------|---------|
| <b>Route of Administration</b> | TOPICAL |
|--------------------------------|---------|

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength  | Strength          |
|---|--------------------|-------------------|
| <b>PIPERONYL BUTOXIDE</b> (UNII: LWK91TU9AH) (PIPERONYL BUTOXIDE - UNII:LWK91TU9AH) | PIPERONYL BUTOXIDE | 4 mL in 100 mL    |
| <b>PYRETHRUM EXTRACT</b> (UNII: ZUM06L90GV) (PYRETHRUM EXTRACT - UNII:ZUM06L90GV)   | PYRETHRUM EXTRACT  | 0.33 mL in 100 mL |

### Inactive Ingredients

| Ingredient Name                              | Strength |
|--|----------|
| <b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)     |          |
| <b>WATER</b> (UNII: 059QF0KO0R)              |          |
| <b>XANTHAN GUM</b> (UNII: TTV12P4NEE)        |          |
| <b>C13-14 ISOPARAFFIN</b> (UNII: E4F12ROE70) |          |

### Product Characteristics

|              |  |              |  |
|--------------|--|--------------|--|
| <b>Color</b> | YELLOW (cloudy to transparent, cream to straw) | <b>Score</b> |  |
| <b>Shape</b> |  | <b>Size</b>  |  |



|                 |  |                     |  |
|-----------------|--|---------------------|--|
| <b>Flavor</b>   |  | <b>Imprint Code</b> |  |
| <b>Contains</b> |  |                     |  |

| <b>Packaging</b> |                  |                     |                      |                    |
|------------------|------------------|---------------------|----------------------|--------------------|
| #                | Item Code        | Package Description | Marketing Start Date | Marketing End Date |
| 1                | NDC:0363-0866-26 | 1 in 1 CARTON       |                      |                    |
| 1                |                  | 118 mL in 1 BOTTLE  |                      |                    |

| <b>Marketing Information</b> |  |                      |                    |
|------------------------------|--|----------------------|--------------------|
| Marketing Category           | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| OTC monograph final          | part358G                                 | 09/10/2007           |                    |

**Part 2 of 2**

**EGG AND NIT**  
combing aid gel

| <b>Product Information</b>     |         |
|--------------------------------|---------|
| <b>Route of Administration</b> | TOPICAL |

| <b>Inactive Ingredients</b>            |          |
|--|----------|
| Ingredient Name                        | Strength |
| WATER (UNII: 059QF0KO0R)               |          |
| GLYCOLIC ACID (UNII: 0WT12SX38S)       |          |
| POTASSIUM HYDROXIDE (UNII: WZH3C48M4T) |          |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3)    |          |
| METHYL PARABEN (UNII: A2I8C7HI9T)      |          |
| PROPYL PARABEN (UNII: Z8IX2SC1OH)      |          |

| <b>Packaging</b> |           |                     |                      |                    |
|------------------|-----------|---------------------|----------------------|--------------------|
| #                | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1                |           | 57 g in 1 BOTTLE    |                      |                    |

| <b>Marketing Information</b> |  |                      |                    |
|------------------------------|--|----------------------|--------------------|
| Marketing Category           | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| Unapproved drug other        |  | 10/29/2007           |                    |

## Marketing Information

| Marketing Category  | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part358G                                 | 10/29/2007           |                    |

**Labeler** - Walgreen Company (008965063)

Revised: 6/2012

Walgreen Company