

ADVANCE TOPICAL ANESTHETIC GEL- benzocaine gel
The Atlanta Dental Supply Company DBA Advanced Dental Products

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active Ingredients

Benzocaine 200mg (in each g)

Purpose

Oral Anesthetic

Use

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use.

Keep Out of Reach of Children.

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Other Information

Store at 59° -86°F (15 - 30°C).

Protect from freezing.

Inactive ingredients

flavoring, PEG 3350, PEG 400, sodium saccharin, water. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

Questions or comments?

Call: 877-896-7341

Manufactured for:
Advanced Dental Products
Atlanta, GA 30359



GLUTEN FREE!
20% Benzocaine
1 fl. oz. (30 ml)

Advancing the Quality of
Dental Care Products

Topical Anesthetic Gel

Made in USA



directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.

- for teething
- in children under 2 years of age

When using this product

Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use.

Keep out of reach of children.

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Other information

Store at 59° - 86°F (15 - 30°C).
Protect from freezing.

Inactive ingredients

flavoring, PEG 3350, PEG 400, sodium saccharin, water. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

Questions or comments?
Call: 877-896-7341

4916467 Rev 06/2018

Drug Facts
Active ingredients Benzocaine 200 mg (in each g)
Purpose Oral Anesthetic
Use For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.
Warnings Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: <ul style="list-style-type: none"> • pale, gray or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics. Do not use <ul style="list-style-type: none"> • for more than 7 days unless

ADVANCE TOPICAL ANESTHETIC GEL

benzocaine gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71347-101
Route of Administration	DENTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

WATER (UNII: 059QF0KO0R)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
D&C GREEN NO. 5 (UNII: 8J6RDU8L9X)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
FD&C YELLOW NO. 5 (UNII: I753WB2F1M)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71347-101-30	30 g in 1 JAR; Type 0: Not a Combination Product	06/01/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	03/01/2017	

Labeler - The Atlanta Dental Supply Company DBA Advanced Dental Products (033505389)