

CETIRIZINE HYDROCHLORIDE HIVES RELIEF- cetirizine tablet
Amneal Pharmaceuticals LLC

CETIRIZINE HYDROCHLORIDE TABLETS

Drug Facts

ACTIVE INGREDIENT

(in each tablet)

Cetirizine HCl, USP 5 mg

PURPOSE

Antihistamine

INDICATIONS AND USAGE

Relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

WARNINGS

Severe Allergy Warning: Get emergency help **immediately** if you have hives along with any of the following symptoms:

- trouble swallowing
- dizziness or loss of consciousness
- swelling of tongue
- swelling in or around mouth
- trouble speaking
- drooling
- wheezing or problems breathing

These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional **immediately**. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.

Not a Substitute for Epinephrine . If your doctor has prescribed an epinephrine injector for “anaphylaxis” or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.

DO NOT USE

Do not use

- to **prevent** hives from any known cause such as:
- foods • insect stings • medicines • latex or rubber gloves

because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for medical exam. Your doctor may be able to help you find a cause.

- if you have ever had an allergic reaction to this product or its ingredients or to an antihistamine containing hydroxyzine.

ASK DOCTOR

Ask a doctor before use if you have

- liver or kidney disease. Your doctor should determine if you need a different dose.
- hives that are an unusual color, look bruised or blistered
- hives that do not itch

ASK DOCTOR/PHARMACIST

Ask a doctor of pharmacist before use if you are taking tranquilizers or sedatives.

WHEN USING

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

STOP USE

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve after 3 days of treatment
- the hives have lasted more than 6 weeks

PREGNANCY OR BREAST FEEDING

If pregnant or breast-feeding:

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

DIRECTIONS

Adults and children 6 years and over	1 tablet (5 mg) or 2 tablets (10 mg) once daily depending upon severity of symptoms; do not take more than 2 tablets (10 mg) in 24 hours.
Adults 65 years and over	1 tablet (5 mg) once daily; do not take more than 1 tablet (5 mg) in 24 hours.
Children under 6 years of age	ask a doctor
Consumers with liver or kidney disease	ask a doctor

OTHER INFORMATION***Other information***

- store between 20° to 25°C (68° to 77°F)

INACTIVE INGREDIENTS***Inactive Ingredients***

lactose monohydrate, magnesium stearate, polyvinyl alcohol, polyethylene glycol, povidone, starch, talc and titanium dioxide.

OTC - QUESTIONS***Questions or Comments?***

Call 1-877-835-5472

Monday through Friday 9AM - 5PM EST.

Distributed by:

Amneal Pharmaceuticals LLC
Bridgewater, NJ 08807

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC 65162-145-03

Cetirizine Hydrochloride Tablets, USP

5 mg

Antihistamine
HIVES RELIEF

30 Tablets



Drug Facts

Active ingredient (in each tablet) Purpose
Cetirizine HCl, USP 5 mg.....Antihistamine

Uses
relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

Warnings
Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms:
• trouble swallowing • dizziness or loss of consciousness • swelling of tongue • swelling in or around mouth • trouble speaking • drooling • wheezing or problems breathing

Do not use if imprinted foil inner seal on bottle is broken or missing.

Distributed by: Amneal Pharmaceuticals LLC
Bridgewater, NJ 08807

Rev. 08-2023-01


N 3 65162 14503 8

Lot No:
Exp. Date:

Non-Varnish Area

PEEL BACK FOR ADDITIONAL DRUG FACTS

Drug Facts (continued)

These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.

Not a Substitute for Epinephrine. If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.

Do not use
• to prevent hives from any known cause such as:
• foods • insect stings • medicines • latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for medical exam. Your doctor may be able to help you find a cause.
• if you have ever had an allergic reaction to this product or its ingredients or to an antihistamine containing hydroxyzine.

Ask a doctor before use if you have
• liver or kidney disease. Your doctor should determine if you need a different dose.
• hives that are an unusual color, look bruised or blistered • hives that do not itch

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product • drowsiness may occur • avoid alcoholic drinks • alcohol, sedatives, and tranquilizers may increase drowsiness • be careful when driving a motor vehicle or operating machinery

Drug Facts (continued)

Stop use and ask a doctor if • an allergic reaction to this product occurs. Seek medical help right away. • symptoms do not improve after 3 days of treatment • the hives have lasted more than 6 weeks

If pregnant or breast-feeding:
• If breast - feeding: not recommended
• If pregnant: ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

adults and children 6 years and over
1 tablet (5 mg) or 2 tablets (10 mg) once daily depending upon severity of symptoms; do not take more than 2 tablets (10 mg) in 24 hours.

adults 65 years and over
1 tablet (5 mg) once daily; do not take more than 1 tablet (5 mg) in 24 hours.

children under 6 years of age
ask a doctor

consumers with liver or kidney disease
ask a doctor

Other information

• store between 20° to 25°C (68° to 77°F)

Inactive ingredients

lactose monohydrate, magnesium stearate, polyvinyl alcohol, polyethylene glycol, povidone, starch, talc and titanium dioxide.

Questions or Comments?

Call 1-877-835-5472
Monday through Friday 9AM - 5PM EST.



CETIRIZINE HYDROCHLORIDE HIVES RELIEF

cetirizine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:65162-145
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CETIRIZINE HYDROCHLORIDE (UNII: 64O047KTOA) (CETIRIZINE - UNII:YO7261ME24)	CETIRIZINE HYDROCHLORIDE	5 mg

Inactive Ingredients				
Ingredient Name			Strength	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
POVIDONE (UNII: FZ989GH94E)				
STARCH, CORN (UNII: O8232NY3SJ)				
TALC (UNII: 7SEV7J4R1U)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
Product Characteristics				
Color	white	Score	no score	
Shape	OVAL	Size	6mm	
Flavor		Imprint Code	IP;45	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65162-145-03	1 in 1 CARTON	01/21/2010	
1		30 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:65162-145-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	01/21/2010	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA078780	01/21/2010	

Labeler - Amneal Pharmaceuticals LLC (123797875)

Establishment

Name	Address	ID/FEI	Business Operations
Amneal Pharmaceuticals of New York, LLC		831227801	analysis(65162-145) , label(65162-145) , manufacture(65162-145) , pack(65162-145)

Revised: 10/2024

Amneal Pharmaceuticals LLC