ASPEN IODINE CONC- iodine solution Aspen Veterinary Resources

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

FOR INDUSTRIAL USE ONLY KEEP OUT OF REACH OF CHILDREN NOT FOR HUMAN USE

Attention

For External Use Only

Wash hands thoroughly after handling.

Get medical advice/attention if you feel unwell.

DIRECTIONS:

IODINE CONCENTRATE MUST BE DILUTED BEFORE USE DO NOT MIX WITH ANYTHING BUT COOL, POTABLE WATER FOR USE ONLY IN A DISPENSING SYSTEM RECOMMENDED BY ECOLAB

Dilution Parameter: A dilute solution of a 0.5% iodine teat dip will consist of 87.5% potable water and 12.5% Iodine Concentrate by volume. A dilute solution of a 1% iodine teat dip will consist of 75% potable water and 25% Iodine Concentrates by volume.

IMPORTANT: Do not mix Iodine Concentrate or diluted Iodine Concentrate with any other products unless specified in the product use instructions. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for Iodine Concentrate. If product in dip cup becomes visibly dirty, discard contents and replenish with diluted, ready-to-use product. Do not reuse or return any unused product to the original container.

Before Milking: Wash teats thoroughly just prior to next milking with appropriate udder wash solution or pre-milking teat dip to avoid contamination of milk. Teats should then be dried with single-service towels. Use proper procedures for udder washing or pre-dipping.

Application of Diluted Iodine Concentrate

Directions for Teat Dipping

Pre-Milk Dipping: Fill teat dip cup with diluted Iodine Concentrate. Before each cow is milked, dip each teat full-length into the teat dip cup containing diluted Iodine Concentrate. Wipe teats dry after dipping using single-service towels to avoid contamination of milk.

Post-Milk Dipping: Immediately after each milking, dip each teat full-length into the teat dip cup containing diluted Iodine Concentrate. Allow to air dry. Do not wipe. Always use fresh, diluted Iodine Concentrate. Do not turn cows out in freezing weather until teat dip

is completely dry.

Directions for Teat Spraying

Pre-Milk Spraying: Before each cow is milked, spray entire teat with diluted Iodine Concentrate. Wipe teats after spraying using single-service towels to avoid contamination of milk.

Post-Milk Spraying: Immediately after each milking, spray entire teat with diluted Iodine Concentrate. Allow to air dry. Do not wipe. Always use fresh, diluted Iodine Concentrate. Do not turn cows out in freezing weather until diluted Iodine Concentrate is completely dry.

Expanded Usage: When freshening cows, begin dipping teats twice daily for about 10 days before calving.

PRECAUTION: Iodine Concentrate is not intended to cure or help the healing of chapped or irritated teats. As with any germicide, irritation or sensitization may occur in sensitive animals. In case of teat irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.

NOTE: Use blended Iodine Concentrate within six (6) months of blended date.

READ SAFETY DATA SHEET (SDS) BEFORE USING THIS PRODUCT

EMERGENCY HEALTH INFORMATION: Contact local poison control.

Representative label and principal display panel aspen DAIRY SOLUTIONS

IODINE CONCENTRATE

ACTIVE INGREDIENT

Titratable Iodine......3.6%

INERT INGREDIENTS:.....96.4%

(contains glycerin, propylene glycol)

TOTAL:.....100.0%

Iodine Teat Dip Concentrate

An aid in reducing the spread of organisms which may cause mastitis.

208 L (55 US GAL)

6302032

Manufactured for:

Aspen Veterinary Resources®, Ltd. 3155 Heartl1-and Drive, Liberty, MO 64068

1-800-203-5620

766884/5301/0217



IODINE CONCENTRATE



lodine Teat Dip Concentrate An aid in reducing the spread of organisms which may cause mastitis FOR INDUSTRIAL USE ONLY

IODINE CONCENTRATE MUST BE DILUTED BEFORE USE

100 ME CONCENTRATE MUST BE DILUTED BEFORE USE
DO NOT MIX WITH ANYTHING BUT CODE, POTRAGE WARTER
FOR USE ONLY IN A DISPENSING SYSTEM RECOMMENDED BY EACH DISPENSION OF THE PROPERTY OF THE PROP

any ounces provides to deep concession extraction.

Before Milliang: Wish that's throughly just prior to next milling with appropriate udder wash solution or pre-milling teat dip to avoid contamination of milli. Teats should then be deep with single-service towels. Use proper procedures for odder washing or pre-dipping. Application of Divided Indian Contamination of milling and the procedures of the procedures for odder washing or pre-dipping.

Application of Divided Indian Concession and the procedures for odder washing or pre-dipping.

Appreciation of unables forming consensations. Districtions for a first Rights and the second of the plant factor for the plant factor for the plant factor for the plant factor forming from the plant factor forming district forming consensations while test of yeller dispring using single-service towers to avoid contamination of milk. Post-Milk Dipping: Immediately after each milking, dip each text fall-length into the text dip cap containing district forming containing containing district forming containing containi

to just one wayer whelp is the first, unless domes continuous to not cause on our meeting resolute cause easily is conjected by J. Directediant for less favoring. Pre-Milit Sparying: Debre each come in milited, spray entire teat with diluted lodine Concentrate. Wipe leads after spraying using single-service towels to avoid contamination of milit.

avaid contamination of milk.

Pers-AHILS payang immediately after each milking, spray entire teal with diluted lodine Concentrate. Allow to air dry, Do not wipe. Always use fresh, diluted lodine Concentrate is completely dry.

Espanded Vlages, When notaving own, begin dipping test twick paid by red hours of days before calling.

PECAUTION: John Concentrate is not intended to our or help the healing of chapped or imitated tests. As with any germidde, initiation or secretization may contribute the secondary that the condition examined and, if necessary, treated by a veterinarian.

NOTE: Use blended lodine Concentrate within six (6) months or blended date.

KEEP OUT OF REACH OF CHILDREN

NOT FOR HUMAN USE

Attention

For External Use Only
Wash hands thoroughly after handling. Get medical advice/ attention if you feel unwell.

READ SAFETY DATA SHEET (SDS) BEFORE USING THIS PRODUCT

EMERGENCY HEALTH INFORMATION: Contact local poison control.

Baño antiséptico concentrado para pezones con iodo Ayuda a evitar la proliferación de organismos que pueden causar mastitis PARA USO INDUSTRIAL SOLAMENTE

INSTRUCCIONES:

IODINE CONCENTRATE DILUIR ANTES DE USAR

NOMEX CONCENTRATE DILUIR ANTES DE USAR

WAR OMEXCLAR CON NADA QUE NO SEA AGUA POTABLE FRÍA

USAR D'UNCAMENTE EN UN SISTEMA DE DOSIFICACIÓN RECOMENDADO POR ECOLAB

Parametro de diffusion: Una solución diluida de bala omitisopitio para pezones con indo al 0,5% contendrá 87,5% de agua potable y 12,5% de lodine Concentrate por
volumen. Una solución diluida de bala omitisopitio para pezones con indo al 1% contendrá 75% de agua potable y 25% de lodine Concentrate por
volumen. Una solución diluida de bala omitisopitio para pezones con indo al 1% contendrá 75% de agua potable y 25% de lodine Concentrate por
volumen. Una solución diluida de bala omitisopitio para pezones con indo al 1% contendrá 75% de agua potable y 25% de lodine Concentrate por volumen.
MPORTANTE: Nonezel de lorder Concentrate e olidine Concentrate diluido con tortos productos amones, que se indigue en las instrucciones de uso del producto. Se indigue en las instrucciones de sol del producto have bacarde indigue contraça el eliquetamiento del
servace de foldine Concentrate 3 el oproducto en el regispete para realizar el alho activa dividibientos sució secho el corcientos y reliene el regispete con producto dibid
y licito para usar. No ventra a utilizar el mismo producto, ni ventra a colocar el producto sin usar destro del envace originar parte para paramete paramete a consecuencia paramete sinte paramete paramete a producto paramete paramete

Antes de Ordeñar: Lave los pezones completamente justo antes de ordeñar con una solución de lavado apropiada para ubres o baño antiséptico para pezones previ al ordeño, o rocie el pezón para evitar la contaminación de la leche. Posteriormente, los pezones se deben secar con toallas de servicio único o descartables. Utilice el procedimiento adecuado para el lavado o el prebañado antiséptico de las ubres. Aplicación de dilución lodine Concentrate

representative de una constructiva en l'information de le partici de l'informationes para restaltar el stallado del partici previo al ordefeir. L'inne el ricipiente para el batho de parones con lodine Concentrate diluido. Antes de ordefair cada vara, bahe toda la superficie del partici del par

contaminación de la leche.

Sellado del pezón posterior al ordeño: Immediatamente después de ordeñar cada vaca, bañe toda la superficie del pezón en el recipiente que contiene lodine
Concentrate élinido, beje scar al aire. No utilite paños. Siempre utilize lodine Concentrate finas de liduido. Resta que lodine Concentrate no se haya secado
completamente, no saque las vacas al exterior en dimas gelidos.

Instrucciones para el Rociado de los Rezenes
Rociado previo al ordeño: Antes de ordeñar cada vaca, rocie toda la superficie del pezón con lodine Concentrate diluido. Seque los pezones después de motarlos
utilizando hazilas de un solo servición para evitar la contaminación con la leche.

Rociado posterior al ordeño: Immediatamente después de ordeñar cada vaca, nocie toda la superficie del pezón con lodine Concentrate diluido. Deje secar al aire.
No utiliza paños. Siempre utiliza lodine Concentrate después de ordeñar cada vaca, nocie toda la superficie del pezón con lodine Concentrate diluido. Deje secar al aire.
No utiliza paños. Siempre utiliza lodine Concentrate después de ordeñar cada vaca, nocie toda la superficie del pezón con lodine Concentrate diluido. Deje secar al aire.
No utiliza paños. Siempre utiliza lodine Concentrate después de ordeñar cada vaca, nocie toda la superficie del pezón con lodine Concentrate diluido. Deje secar al aire.

complete.

Web Expandidor. Al reflexcar a las vacas, comience con un baino antiséptico de sus pezones dos veces por dia por aproximadamente 10 dias antes de parir.

PRECAUCONE: odine Concentrate no está dischado para curar o ayudar con la cicalización de los pezones agrietados o initados. Como cualquier otro germició provocar rimicado en escribilización en animaise sentellas. En coso de initadino a qualquier administrate de los pecenoses, haga examinar dichas afecciones y, de ser nece háspabs tatatar con un veterinario.

WISSO: Utiliza cional concentrate mexidado dentro de los seis (6) meses a partir de la fecha de mexidado.

MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

NO ES PARA USA HUMANO

Atención

Solo de uso externo Lavarse las manos cuidadosamente después de la manipulación.

Consultar a un médico si la persona se encuentra mal.

LEA LA FICHA DE DATOS DE SEGURIDAD ANTES DE USAR EL PRODUCTO

208 L (55 US GAL) 6302032



Manufactured for: Aspen Veterinary Resources*, Ltd. 3155 Heartland Drive, Liberty, MO 64068

1-800-203-5620 766884/5301/0217

ASPEN IODINE CONC

iodine solution

Product	Inform	ation

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:46066-412
--------------	-----------------	--------------------	---------------

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	36 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46066-412-21	208000 mL in 1 DRUM		
2	NDC:46066-412-17	56800 mL in 1 DRUM		

Marketing Information			
Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
	08/17/2016		
	Application Number or Monograph	Application Number or Monograph Marketing Start Citation Date	

Labeler - Aspen Veterinary Resources (627265361)

Revised: 7/2025 Aspen Veterinary Resources