

ONLY FOR HEADACHE- acetaminophen, aspirin, caffeine tablet
O4 Global Trading Usa, Llc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Only For™ HEADACHE

Drug Facts

Active ingredient (in each geltab)

Acetaminophen 250mg

Aspirin 250mg (NSAID*)

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Pain reliever

Pain reliever aid

Uses:

Temporarily relieves minor aches and pains due to:

• headache • a cold • arthritis • muscular aches • toothache • premenstrual & menstrual cramps

Warnings:

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flue-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 8 geltabs in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others) • have 3 or more

alcoholic drinks every day while using this product • take more or for a longer time than directed

Caffeine warning: □The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally rapid heart beat.

Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if • you have liver disease • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma

Ask a doctor or pharmacist before use if you are taking

- a prescription drug for diabetes, gout, or arthritis
- any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if • an allergic reaction occurs. Seek medical help right away. • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • ringing in the ears or loss of hearing occurs • painful area is red or swollen • pain gets worse or lasts for more than 10 days • fever gets worse or lasts for more than 3 days • any new symptoms occur

These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

OVERDOSE WARNING: □Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

- do not use more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 gels every 6 hours; not more than 8 gels in 24 hours
- children under 12 years: ask a doctor

Other Information:

- store at room temperature between 20-25°C (68 to 77°F) • avoid excessive heat, cold and humidity • close cap tightly after use • read all product information before using.

Inactive Ingredients:

ammonium hydroxide, black iron oxide, brilliant blue FCF, corn starch, D&C yellow No.10 lake, ethyl cellulose, FD&C blue No.1 lake, FD&C red No.40 lake, gelatin, glycerin, hydroxypropyl methyl cellulose, isopropyl alcohol, isopropyl alcohol - butyl alcohol, lactose monohydrate, light liquid paraffin, microcrystalline cellulose, polysorb 85/70/00, povidone K30, pregelatinised starch, propylene glycol, purified water, shellac glaze, stearic acid, talc, titanium dioxide, triacetin

Questions or comments?

1-833-665-9444. (Mon-Fri 9am-5pm)

ADULT

PAIN RELIEVER

PAIN RELIEVER AID

TAMPER EVIDENT: Do not use if imprinted safety seal under cap is broken or missing

Distributed By: O4 GLOBAL TRADING USA, LLC.

12222 Merit Drive, Suite 1900

Dallas, TX 75251-3209

www.onlyfor.com

Packaged By: Strive Pharmaceuticals Inc.

East Brunswick, NJ 08816.

Packaging



25 GELTABS
NDC: 73006-132-95

ADULT

Only For™

**HEADACHE
PAIN RELIEVER
PAIN RELIEVER AID**

ACETAMINOPHEN 250mg
Aspirin (NSAID) 250mg / Caffeine 65mg
Pain reliever
Pain reliever aid

TAMPER EVIDENT Do not use if imprinted safety seal under cap is broken or missing



8 60003 57646 5

Drug Facts

Active ingredient (in each geltab) Purpose

Acetaminophen 250mg.....Pain reliever
Aspirin 250mg (NSAID)*.....Pain reliever
Caffeine 65 mg.....Pain reliever aid
*nonsteroidal anti-inflammatory drug

Uses:

Temporarily relieves minor aches and pains due to:
■ headache ■ a cold ■ arthritis ■ muscular aches ■ toothache
■ premenstrual & menstrual cramps

Warnings:

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

Lot No.
Exp. Date

LIFT HERE

Drug Facts

Active ingredient (in each geltab)	Purpose
Acetaminophen 250mg.....	Pain reliever
Aspirin 250mg (NSAID)*.....	Pain reliever
Caffeine 65 mg.....	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Uses:

Temporarily relieves minor aches and pains due to:
■ headache ■ a cold ■ arthritis ■ muscular aches ■ toothache
■ premenstrual & menstrual cramps

Warnings:

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 geltabs in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally rapid heart beat.

Do not use ■ if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking ■ a prescription drug for diabetes, gout, or arthritis ■ any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or loss of hearing occurs ■ painful area is red or swollen ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ any new symptoms occur

These could be signs of a serious condition

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

OVERDOSE WARNING: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

■ do not use more than directed
■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 2 geltabs every 6 hours; not more than 8 geltabs in 24 hours
■ children under 12 years: ask a doctor

Other Information:

■ store at room temperature between 20-25°C (68 to 77°F) ■ avoid excessive heat, cold and humidity ■ close cap tightly after use ■ read all product information before using.

Inactive Ingredients:

ammonium hydroxide, black iron oxide, brilliant blue FCF, corn starch, D&C yellow No.10 lake, ethyl cellulose, FD&C blue No.1 lake, FD&C red No.40 lake, gelatin, glycerin, hydroxypropyl methyl cellulose, isopropyl alcohol, isopropyl alcohol - butyl alcohol, lactose monohydrate, light liquid paraffin, microcrystalline cellulose, polysorb 85/70/00, povidone K30, pregelatinised starch, propylene glycol, purified water, shellac glaze, stearic acid, talc, titanium dioxide, triacetin

Questions or comments?

1-833-665-9444. (Mon-Fri 9am - 5pm)

Distributed By: O4 GLOBAL TRADING USA, LLC.
12222 Merit Drive, Suite 1900
Dallas, TX 75251-3209
www.onlyfor.com

Packaged By: Strive Pharmaceuticals Inc.
East Brunswick, NJ 08816.

3

LB7300613295-REV.00-072020

ONLY FOR HEADACHE

acetaminophen, aspirin, caffeine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73006-132
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
AMMONIA (UNII: 5138Q19F1X)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
BUTYL ALCOHOL (UNII: 8PJ61P6TS3)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SHELLAC (UNII: 46N107B71O)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	green (Dark Green - White)	Score	no score
Shape	ROUND	Size	13mm
Flavor		Imprint Code	S77
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73006-132-95	1 in 1 CARTON	11/15/2020	
1		25 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	11/15/2020	

Labeler - O4 Global Trading Usa, Llc (081226861)

Revised: 11/2020

O4 Global Trading Usa, Llc