

CLARINS EVERLASTING CUSHION LONG-WEARING AND HYDRATING FOUNDATION BROAD SPECTRUM SPF 50 TINT 103- titanium dioxide emulsion

Laboratoires Clarins

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Clarins Everlasting Cushion Long-Wearing and Hydrating Foundation Broad Spectrum SPF 50 Tint 103

Package Labeling:



Drug Facts

Active Ingredient:
TITANIUM DIOXIDE: 6.2%

Purpose:
Sunscreen

Uses:

- helps prevent sunburn.
- if used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings:

For external use only.

Do not use

- on damaged or broken skin.

When using this product

- keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if

- rash occurs.

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions:

- apply liberally 15 minutes before sun exposure.
- apply to all skin exposed to the sun.
- children under 6 months: Ask a doctor.
- **Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. – 2 p.m.
- wear long-sleeved shirts, pants, hats and sunglasses.
- reapply at least every 2 hours.
- use a water resistant sunscreen if swimming or sweating.

Inactive Ingredients: AQUA/WATER/EAU · CYCLOPENTASILOXANE · PHENYL TRIMETHICONE · CYCLOHEXASILOXANE · BUTYLENE GLYCOL · DIPENTAERYTHRITYL HEXA C5-9 ACID ESTERS · TRIETHYLHEXANOIN · CETYL PEG/PPG-10/1 DIMETHICONE · GLYCERIN · PHENYLBENZIMIDAZOLE SULFONIC ACID ▶

www.clarins.com

Drug Facts (continued)

DIMETHICONE · NIACINAMIDE · TROMETHAMINE · 1,2-HEXANEDIOL · SORBITAN SESQUIOLEATE · STEARIC ACID · DIMETHICONE/PEG-10/15 CROSSPOLYMER · NYLON-12 · PEG-30 DIPOLYHYDROXYSTEARATE · SODIUM CHLORIDE · ZINC STEARATE · ALUMINUM HYDROXIDE · DISTEARYL DIMONIUM CHLORIDE · PARFUM/FRAGRANCE · TOCOPHERYL ACETATE · BIS-VINYL DIMETHICONE/DIMETHICONE COPOLYMER · DISTEARDIMONIUM HECTORITE · TRIETHOXYCAPRYLYLSILANE · ADENOSINE · PALMITOYL PROLINE · TRIETHOXSILYLETHYL POLYDIMETHYLSILOXYETHYL HEXYL DIMETHICONE · CHE-NOPODIUM QUINOA SEED EXTRACT · MAGNESIUM PALMITOYL GLUTAMATE · SODIUM PALMITOYL SARCOINATE · CERAMIDE 3 · PALMITIC ACID · HYDROGENATED LECITHIN · LAPSANA COMMUNIS FLOWER/LEAF/STEM EXTRACT · DIPROPYLENE GLYCOL · GLYCERETH-20 · CAMELLIA SINENSIS LEAF EXTRACT · PEG-10 RAPESEED STEROL · GLYCERYL STEARATE · SODIUM BENZOATE · PHENOXYETHANOL.
+/- MAY CONTAIN/PEUT CONTENIR : MICA · SILICA · CI 77891/TITANIUM DIOXIDE · CI 77491/CI 77492/CI 77499/IRON OXIDES. [MS2256A]

Other information:

- protect this product from excessive heat and direct sun.

Questions or comments?

Call 1-866-252-7467 (In the US)
Monday-Friday 8 a.m. to 5 p.m.

Everlasting Cushion

Teint Haute Tenue & Hydratation
Long-Wearing & Hydrating Foundation

Broad Spectrum SPF 50
Titanium Dioxide - Sunscreen
Large Spectre FPS 50
Filtre Solaire - Dioxyde de Titane

103
ivory

13 mL - 0.5 Oz. Net Wt.

Directions for use:

apply liberally 15 minutes before sun exposure. Use a water resistant sunscreen if swimming or sweating. Children under 6 months: ask a doctor.

Cautions and warnings: do not use on damaged and broken skin. Discontinue use if skin irritation develops or increases. If irritation persists, consult a health care practitioner. Avoid contact with eyes. Rinse with water to remove. If product is swallowed, get medical help or contact a Poison Control Center right away.

**ACTIVE INGREDIENT/MEDICINAL INGREDIENT/
INGREDIENT MEDICINAL :**

TITANIUM DIOXIDE/ DIOXYDE DE TITANE : 6.2 %

Questions or comments?
Call 1-866-252-7467 (in the US)
Monday-Friday,
8 a.m. to 5 p.m.

Conseils d'utilisation:

appliquer généreusement 15 minutes avant l'exposition solaire. Utilisez un écran solaire résistant à l'eau si vous nagez ou transpirez. Enfants de moins de 6 mois : demander à un médecin.

Précautions et avertissements : ne pas utiliser sur les peaux abîmées. Cesser l'utilisation si une irritation cutanée se développe ou devient plus sévère. Si l'irritation persiste, consulter un praticien de soins de santé. Éviter tout contact avec les yeux. Rincer avec de l'eau pour l'enlever. En cas d'ingestion du produit, consultez immédiatement un praticien de soins de santé ou contactez un centre anti-poison.

Distrib. Canada: CLARINS CANADA Inc. 815
Desserte est Autoroute Chomedey - LAVAL
(QUEBEC) H7W 5N4.

MFD FOR CLARINS - 92200 Neuilly-sur-Seine
95300 Pontoise - France
Made in Korea



CLARINS EVERLASTING CUSHION LONG-WEARING AND HYDRATING FOUNDATION BROAD SPECTRUM SPF 50 TINT 103

titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58668-9900
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	62 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
CYCLOMETHICONE 6 (UNII: XHK3U310BA)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
DIPENTAERYTHRITYL HEXA C5-9 ACID ESTERS (UNII: D818OLQ51Z)	
TRIETHYLHEXANOIN (UNII: 7K3W1BIU6K)	
GLYCERIN (UNII: PDC6A3C0OX)	
ENSULIZOLE (UNII: 9YQ9DI1W42)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
NIACINAMIDE (UNII: 25X51I8RD4)	
TROMETHAMINE (UNII: 023C2WHX2V)	
1,2-HEXANEDIOL (UNII: TR046Y3K1G)	
SORBITAN SESQUIOLEATE (UNII: 0W8RRI5W5A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
NYLON-12 (UNII: 446U8J075B)	
PEG-30 DIPOLYHYDROXYSTEARATE (UNII: 9713Q0S7FO)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
ZINC STEARATE (UNII: H92E6QA4FV)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
DISTEARYLDIMONIUM CHLORIDE (UNII: OM9573ZX3X)	
.ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
TRIETHOXYCAPRYLSILANE (UNII: LDC331P08E)	
ADENOSINE (UNII: K72T3FS567)	
PALMITOYL PROLINE (UNII: I49727TDYF)	
CHENOPODIUM QUINOA SEED (UNII: OO4K72NZ2X)	
MAGNESIUM PALMITOYL GLUTAMATE (UNII: DH37YM1F48)	
SODIUM PALMITOYL SARCO SINATE (UNII: 7297LY09YF)	
CERAMIDE NP (UNII: 4370DF050B)	
PALMITIC ACID (UNII: 2V16EO95H1)	
HYDROGENATED SOYBEAN LECITHIN (UNII: H1109Z9J4N)	
LAPSANA COMMUNIS FLOWERING TOP (UNII: T9B77O7PZA)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
GLYCERETH-20 (UNII: GJ57297I4F)	
GREEN TEA LEAF (UNII: W2ZU1RY8B0)	
PEG-10 RAPESEED STEROL (UNII: 258O76T85M)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
PHENOXYETHANOL (UNII: HIE49ZZZ3T)	
MICA (UNII: V8A1AW0880)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

FERRIC OXIDE YELLOW (UNII: EX438O2MRT)

FERROSFERRIC OXIDE (UNII: XM0M87F357)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58668-9900-1	13 mL in 1 CASE; Type 0: Not a Combination Product	03/08/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	03/08/2017	

Labeler - Laboratoires Clarins (266317555)

Revised: 3/2019

Laboratoires Clarins