

SULFASOL - sulfadimethoxine sodium powder, for solution
Med-Pharmex, Inc

SULFASOL SOLUBLE POWDER

(sulfadimethoxine)

Antibacterial

CAUTION:Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Approved by FDA under ANADA #200-238

Each packet contains 3.34 oz. (94.6 g) sulfadimethoxine in the form of the soluble sodium salt and disodium edetate.

Restricted Drug (California) - Use Only as Directed. Not for Human Use.

For Broiler and Replacement Chickens Only - Use for the treatment of disease outbreaks of coccidiosis, fowl cholera, and infectious coryza.

For Meat-producing Turkeys Only - Use for the treatment of disease outbreaks of coccidiosis, and fowl cholera.

For Dairy Calves, Dairy Heifers, and Beef Cattle - Use for the treatment of shipping fever complex and bacterial pneumonia associated with *Pasteurella* spp. sensitive to sulfadimethoxine; and calf diphtheria and foot rot associated with *Sphaerophorus necrophorus* sensitive to sulfadimethoxine.

Net Wt.: 3.77 oz (107g)

		SULFADIMETHOXINE	
		SOLUBLE POWDER	
Dosage and Administration:			
Species	Concentration	Use of Directions	
Chickens	0.05%	Contents of packet to 50 gal of water	
Turkeys	0.025%	Contents of packet to 100 gal of water	
Automatic Proportioners: To make stock solution, add contents of 5 packets to 2 gal of water for chickens and to 4 gal of water for turkeys. Set proportioner to feed at a rate of 1 fl oz stock solution per gal of water.			
Treatment Period: 6 consecutive days			
Dairy Calves, Dairy Heifers, and Beef Cattle			
Dosage: 25 mg/lb first day followed by 12.5 mg/lb/day for 4 days			
		Sulfadimethoxine in Water	
		Water Consumption	
	Amount of Stock Solution for Cattle*	(Summer)* 1 gal/100 lb body weight**	(Winter)* 1 gal/150 lb body weight**
First Day Add	1 qt	10 gal	7 gal
	2 qt	20 gal	14 gal

	1 gal	40 gal	28 gal
Next 4 Days Add	1 qt	20 gal	14 gal
	2 qt	40 gal	28 gal
	1 gal	80 gal	56 gal
<p>*Note: Make a cattle stock solution by adding 1 packet of Sulfadimethoxine Soluble Powder to 1 gal of water.</p> <p>*Twenty fl oz of cattle stock solution will medicate 1-600 lb animal initially or 2-600 lb animals on maintenance dose. Contents of packet will medicate 6-600 lb animals initially or 12-600 lb animals on maintenance dose.</p> <p>**This dosage recommendation is based on a water consumption of 1 gal per 100 lb of body weight per day, the expected water consumption rate for summer. Water consumption during cold months (winter) may drop markedly (30%-40%). Accordingly, adjustments must be made in the dilution rates to compensate for this and insure proper drug intake.</p> <p>For treatment of individual cattle, Sulfadimethoxine Soluble Powder stock solution for cattle may be given as a drench.</p> <p>Treatment Period: 5 consecutive days</p>			

Caution: *Chickens and Turkeys* - If animals show no improvement within 5 days, discontinue treatment and reevaluate diagnosis. Prepare a fresh stock solution daily. Handle the recommended dilutions (chickens 0.05% and turkeys 0.025%) as regular drinking water. Administer as sole source of drinking water and sulfonamide medication. Chickens and turkeys that have survived fowl cholera outbreaks should not be kept for replacements or breeders.

Cattle - During treatment period, make certain that animals maintain adequate water intake. If animals show no improvement within 2 or 3 days, reevaluate diagnosis. Treatment should not be continued beyond 5 days.

Warning:

Chickens and Turkeys - Withdraw 5 days before slaughter. Do not administer to chickens over 16 weeks (112 days) of age or to turkeys over 24 weeks (168 days) of age.

Cattle - Withdraw 7 days before slaughter. For dairy calves, dairy heifers and beef cattle only.

A withdrawal period has not been established for this product in pre-ruminating calves.

Do Not Use in Calves to be Processed For Veal

CONTACT INFORMATION: To report adverse drug events, for technical assistance, or to obtain a copy of the Safety Data Sheet (SDS), please contact MED-PHARMEX at (800) 587-4306.

For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at <http://www.fda.gov/reportanimalae>

MANUFACTURED BY: Med-Pharmex, Inc., Pomona CA 91767

Rev. 06/2023

NDC 54925-035-37

SULFASOL (sulfadimethoxine)

Antibacterial

SOLUBLE POWDER

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Each packet contains 3.34 oz. (94.6 g) sulfadimethoxine in the form of the soluble sodium salt and disodium edetate.

Restricted Drug (California) - Use Only as Directed.

Not for use in Humans.

For Broiler and Replacement Chickens Only - Use for the treatment of disease outbreaks of coccidiosis, fowl cholera, and infectious coryza.

For Meat-producing Turkeys Only - Use for the treatment of disease outbreaks of coccidiosis, and fowl cholera.

For Dairy Calves, Dairy Heifers, and Beef Cattle - Use for the treatment of shipping fever complex and bacterial pneumonia associated with *Pasteurella* spp. sensitive to sulfadimethoxine; and calf diphtheria and foot rot associated with *Sphaerophorus necrophorus* sensitive to sulfadimethoxine.

Net Wt.: 3.77 oz (107g)



Approved by FDA under ANADA # 200-238

Rev. March 2023

SULFADIMETHOXINE SOLUBLE POWDER

Dosage and Administration:

Species	Concentration	Use of Directions
Chickens	0.05%	Contents of packet to 50 gal of water
Turkeys	0.025%	Contents of packet to 100 gal of water

Automatic Proportioners: To make stock solution, add contents of 5 packets to 2 gal of water for chickens and to 4 gal of water for turkeys. Set proportioner to feed at a rate of 1 fl oz stock solution per gal of water.

Treatment Period: 6 consecutive days

Dairy Calves, Dairy Heifers, and Beef Cattle

Dosage: 25 mg/lb first day followed by 12.5 mg/lb/day for 4 days

Sulfadimethoxine in Water

	Amount of Stock Solution for Cattle*	Water Consumption	
		(Summer)* 1 gal/100 lb body weight**	(Winter)* 1 gal/150 lb body weight**
First Day Add	1 qt	10 gal	7 gal
	2 qt	20 gal	14 gal
	1 gal	40 gal	28 gal
Next 4 Days Add	1 qt	20 gal	14 gal
	2 qt	40 gal	28 gal
	1 gal	80 gal	56 gal

***Note:** Make a cattle stock solution by adding 1 packet of Sulfadimethoxine Soluble Powder to 1 gal of water. *Twenty fl oz of cattle stock solution will medicate 1-600 lb animal initially or 2-600 lb animals on maintenance dose. Contents of packet will medicate 6-600 lb animals initially or 12-600 lb animals on maintenance dose.

**This dosage recommendation is based on a water consumption of 1 gal per 100 lb of body weight per day, the expected water consumption rate for summer. Water consumption during cold months (winter) may drop markedly (30%-40%). Accordingly, adjustments must be made in the dilution rates to compensate for this and insure proper drug intake.

For treatment of individual cattle, Sulfadimethoxine Soluble Powder stock solution for cattle may be given as a drench.

Treatment Period: 5 consecutive days

Caution: Chickens and Turkeys - If animals show no improvement within 5 days, discontinue treatment and reevaluate diagnosis. Prepare a fresh stock solution daily. Handle the recommended dilutions (chickens 0.05% and turkeys 0.025%) as regular drinking water. Administer as sole source of drinking water and sulfonamide medication. Chickens and turkeys that have survived fowl cholera outbreaks should not be kept for replacements or breeders.

Cattle - During treatment period, make certain that animals maintain adequate water intake. If animals show no improvement within 2 or 3 days, reevaluate diagnosis. Treatment should not be continued beyond 5 days.

Warning:

Chickens and Turkeys - Withdraw 5 days before slaughter. Do not administer to chickens over 16 weeks (112 days) of age or to turkeys over 24 weeks (168 days) of age.

Cattle - Withdraw 7 days before slaughter. For dairy calves, dairy heifers and beef cattle only. A withdrawal period has not been established for this product in pre-ruminating calves.

Do Not Use in Calves to be Processed For Veal

CONTACT INFORMATION:

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact MED-PHARMEX at (800) 587-4306.

For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at <http://www.fda.gov/reportanimalae>

MANUFACTURED BY: Med-Pharmex, Inc., Pomona, CA 91767

SULFASOL

sulfadimethoxine sodium powder, for solution

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:54925-035
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SULFADIMETHOXINE SODIUM (UNII: 49DG2B481W) (SULFADIMETHOXINE - UNII:30CPC5LDEX)	SULFADIMETHOXINE SODIUM	94.6 g in 107 g

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54925-035-37	107 g in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

ANADA

ANADA200238

07/28/1998

Labeler - Med-Pharmex, Inc (025353699)

Revised: 6/2023

Med-Pharmex, Inc