

**ACETAMINOPHEN- acetaminophen tablet**  
**Akron Pharma Inc.**

-----  
**Acetaminophen 500 mg Tablets**  
**Extra Strength**

**Active ingredient (in each tablet)**

Acetaminophen 500 mg

**Purpose**

Pain Reliever/Fever Reducer

**Uses**

To reduce fever and for the temporary relief of minor aches and pains due to:

- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps.
- Temporarily reduces fever.

**Warnings**

**Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if the user has ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if the user has**

- has liver disease
- is a child with pain of arthritis

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin**

**Stop use and ask a doctor if:**

- pain gets worse or lasts more than 10 days (for adults) or 5 days (for children)
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present.

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose Warning:**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions - do not take more than directed (see overdose warning)**

---

adults & children 12 years and over	<ul style="list-style-type: none"><li>• take 2 tablets every 6 hours while symptoms last</li><li>• do not take more than 6 tablets in 24 hours, unless directed by a doctor</li><li>• do not use for more than 10 days unless directed by a doctor</li></ul>
children under 12 years	<ul style="list-style-type: none"><li>• ask a doctor</li></ul>

---

**Other information**

- store at 20° to 25°C (68° to 77°F)

## Inactive Ingredients:

Gelatinized starch, magnesium stearate, povidone

## Questions or Comments?

Call toll-free 1-877-225-6999

Manufactured for  
Akron Pharma, Inc.,

373 RT US46 W Building E,

Suite 117, Fairfield, NJ - 07004

\*This product is not manufactured or distributed by Johnson and Johnson, consumer inc., distributor of regular Tylenol Tablets.

<b>Drug Facts</b>
<b>Active ingredient (in each tablet)</b> Acetaminophen 500 mg ..... Pain reliever/ Fever reducer
<b>Uses</b> ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscle aches ■ backache ■ minor pain of arthritis ■ the common cold ■ toothache ■ premenstrual and menstrual cramps ■ temporarily reduces fever
<b>Warnings</b> <b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product <b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.
<b>Do not use</b> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if the user has ever had an allergic reaction to this product or any of its ingredients.
<b>Ask a doctor before use if the user</b> ■ has liver disease
<b>Ask a doctor or pharmacist before use if the user is</b> ▲ taking the blood thinning drug warfarin

**Child-Resistant Packaging**

**Drug Facts (Continued under label)**

**Do not use if imprinted safety seal under cap is broken or missing**

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Extra Strength**

KTK251666/2017  
 Akron Pharma, Inc.  
 Akron, Ohio 44316  
 373 RT US 46 W Building E  
 Fairfield, NJ 07004  
 www.akronpharma.com  
 REC 18/24

No Tamish

PEEL HERE FOR MORE DRUG FACTS

**Drug Facts (continued)**

**Stop use and ask a doctor if** ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present.  
 These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.


**Directions** ■ do not take more than directed (see overdose warning)  
**adults & children**  
 12 years and over:  
 ■ take 2 tablets every 6 hours while symptoms last  
 ■ do not take more than 6 tablets in 24 hours, unless directed by a doctor  
 ■ do not use for more than 10 days unless directed by a doctor

**Children under 12 years:**  
 ask a doctor

**Other information:** store at 20° to 25°C. (68° to 77°F)

**Inactive ingredients:** Gelatinized starch, magnesium stearate, povidone

**Questions or comments?** Call toll-free 1-877-225-6999



**Acetaminophen**  
**500MG TABLETS**  
**EXTRA STRENGTH**

1000 TABLETS

**Drug Facts**

**Active ingredient (in each tablet)**  
Acetaminophen 500 mg

**Purpose**  
Pain reliever/ Fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - backache
  - minor pain of arthritis
  - the common cold
  - toothache
  - premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if the user has ever had an allergic reaction to this product or any of the ingredients.

**Ask a doctor before use if the user**

- has liver disease

**Ask a doctor or pharmacist before use if the user is** taking the blood thinning drug warfarin

Child-Resistant Packaging

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present.

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** do not take more than directed (see overdose warning)

<b>adults &amp; children 12 years and over</b>	take 2 tablets every 6 hours while symptoms last
	do not take more than 6 tablets in 24 hours, unless directed by a doctor
	do not use for more than 10 days unless directed by a doctor
<b>child under 12 years</b>	ask a doctor

**Other information:** store at 20° to 25° C (68° to 77° F)

**Inactive ingredients:** gelatinized starch, magnesium stearate, povidone.

**Questions or comments?** Call toll-free 1-877-225-6999

Do not use if imprinted safety seal under cap is broken or missing

\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Extra Strength

KTK25/69/2017

**AKRON Pharma**

Manufactured for:  
Akron Pharma, Inc.  
10000 Building E  
Suite 117, Fairfield, NJ 07004  
www.akronpharma.com  
Rev. 06/24

No Varnish

<b>ACETAMINOPHEN</b> acetaminophen tablet				
<b>Product Information</b>				
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:71399-1026	
<b>Route of Administration</b>	ORAL			
<b>Active Ingredient/Active Moiety</b>				
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	500 mg	
<b>Inactive Ingredients</b>				
<b>Ingredient Name</b>			<b>Strength</b>	
STARCH, CORN (UNII: O8232NY3S)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
POVIDONE (UNII: FZ989GH94E)				
<b>Product Characteristics</b>				
<b>Color</b>	white	<b>Score</b>	2 pieces	
<b>Shape</b>	ROUND	<b>Size</b>	5mm	
<b>Flavor</b>		<b>Imprint Code</b>	A500	
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:71399-1026-1	100 in 1 BOTTLE; Type 0: Not a Combination Product	12/06/2024	
2	NDC:71399-1026-2	1000 in 1 BOTTLE; Type 0: Not a Combination Product	12/06/2024	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/06/2024	

**Labeler** - Akron Pharma Inc. (067878881)

Revised: 1/2026

Akron Pharma Inc.