

LORATADINE AND PSEUDOEPHEDRINE- loratadine and pseudoephedrine tablet, film coated, extended release
Walgreen Company

Loratadine and Pseudoephedrine

Drug Facts

Active ingredients (in each tablet)	Purpose
Loratadine, USP 10 mg	Antihistamine
Pseudoephedrine sulfate, USP 240 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - sneezing
 - itchy, watery eyes
 - runny nose
 - itching of the nose or throat
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not divide, crush, chew or dissolve the tablet

adults and children 12 years and over	1 tablet daily with a full glass of water; not more than 1 tablet in 24 hours
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- **sodium:** contains 10 mg/tablet
- **calcium:** contains 25 mg/tablet
- **TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.**
- store between 20° C to 25° C (68° F to 77° F)
- protect from light and store in a dry place

Inactive ingredients

calcium carbonate, colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, iron oxide black, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glaze, sodium alginate, sodium citrate, talc and titanium dioxide

Questions?

call **1-800-406-7984**

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015

PRINCIPAL DISPLAY PANEL - 240 mg/10 mg Tablet Blister Pack Carton

ORIGINAL PRESCRIPTION STRENGTH

NDC 0363-0725-15

Walgreens

Compare to the active ingredients in

Claritin-D[®] 24 Hour^{††}

•WALGREENS•

PHARMACIST RECOMMENDED[†]

NON-DROWSY*

Allergy Relief D

Nasal Decongestant

LORATADINE, USP 10 mg / ANTIHISTAMINE

PSEUDOEPHEDRINE SULFATE, USP 240 mg / NASAL DECONGESTANT

24 Hour

Indoor & Outdoor Allergies

- Relief of nasal & sinus congestion due to colds or allergies
- Relief of sneezing, runny nose, itchy, watery eyes & itchy throat or nose due to allergies

15

EXTENDED-RELEASE

TABLETS

*WHEN TAKEN AS DIRECTED.

SEE DRUG FACTS PANEL.

ACTUAL SIZE

Drug Facts (continued)

Other information

- sodium: contains 10 mg/tablet
- calcium: contains 25 mg/tablet
- **TAMPER EVIDENT: DO NOT USE IF PLUSTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.**
- store between 20° C (68° F) to 25° C (77° F).
- protect from light and store in a dry place.

Inactive ingredients calcium carbonate, colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, iron oxide black, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shelec glaze, sodium alginate, sodium citrate, sodium chloride and titanium dioxide

Questions? Call 1-800-406-7984

Drug Facts

Active ingredients (in each tablet)

- Loratadine, USP 10 mg.....Antihistamine
- Pseudoephedrine sulfate, USP 240 mg.....Nasal decongestant

Purpose

Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: sneezing, itchy, watery eyes; itchy, watery nose; runny nose; reduces swelling of nasal passages; temporarily relieves sinus congestion and pressure; temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies; temporarily restores freer breathing through the nose.

Warnings

- Do not use if you have ever had an allergic reaction to this product or any of its ingredients.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if you are taking a prescription drug containing an MAOI, ask a doctor or pharmacist before taking this product.

Uses

Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: sneezing, itchy, watery eyes; itchy, watery nose; runny nose; reduces swelling of nasal passages; temporarily relieves sinus congestion and pressure; temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies; temporarily restores freer breathing through the nose.

Directions

- do not divide, crush, chew or dissolve the tablet.
- adults and children 12 years and over: 1 tablet daily with a full glass of water; not more than 1 tablet in 24 hours.
- children under 12 years of age: ask a doctor.
- consumers with liver or kidney disease: ask a doctor.

Drug Facts (continued)

Ask a doctor before use if you have heart disease, high blood pressure, diabetes, thyroid disease, liver or kidney disease, your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if:

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever.
- nervousness, dizziness or sleeplessness occurs before use.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION. SEE END PANEL FOR EXPIRATION DATE.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
walgreens.com
©2021 Walgreen Co.

ITEM 500830 W00000-0000-0

3 11917 06238 9

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Claritin-D.

Batch No. _____

Expiration Date: _____

Walgreens 

NON-DROWSY*

Allergy Relief D Nasal Decongestant

W3ORG0821-F2

ORIGINAL PRESCRIPTION STRENGTH NDC 0363-0725-15

Walgreens 

Compare to the active ingredients in Claritin-D * 24 Hour™

NON-DROWSY*

Allergy Relief D Nasal Decongestant

LORATADINE, USP 10 mg / ANTIHISTAMINE
PSEUDOEPHEDRINE SULFATE, USP 240 mg / NASAL DECONGESTANT

24 Hour Indoor & Outdoor Allergies

- Relief of nasal & sinus congestion due to colds or allergies
- Relief of sneezing, runny nose, itchy, watery eyes & itchy throat or nose due to allergies

15 EXTENDED-RELEASE TABLETS

***WHEN TAKEN AS DIRECTED. SEE DRUG FACTS PANEL.**

 ACTUAL SIZE

5219620

5219620



LORATADINE AND PSEUDOEPHEDRINE

loratadine and pseudoephedrine tablet, film coated, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0725
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LORATADINE (UNII: 7AJ03BO7QN) (LORATADINE - UNII:7AJ03BO7QN)	LORATADINE	10 mg
PSEUDOEPHEDRINE SULFATE (UNII: Y9DL7QPE6B) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE SULFATE	240 mg

Inactive Ingredients

Ingredient Name	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE (160000 WAMW) (UNII: RFW2ET671P)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
SODIUM ALGINATE (UNII: C269C4G2ZQ)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white (White to Off-White)	Score	no score
Shape	CAPSULE	Size	17mm
Flavor		Imprint Code	RX724
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC 0363			

1	NDC:0363-0725-56	1 in 1 CARTON	11/07/2013	
1		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0363-0725-69	1 in 1 CARTON	11/07/2013	
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:0363-0725-15	1 in 1 CARTON	11/07/2013	
3		15 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076557	11/07/2013	

Labeler - Walgreen Company (008965063)

Establishment

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		051565745	manufacture(0363-0725)

Revised: 8/2021

Walgreen Company