

**GOOD SENSE CHILDRENS PAIN AND FEVER- acetaminophen suspension**  
**L. Perrigo Company**

-----  
**Perrigo Children's Pain & Fever Drug Facts**

**Active ingredient (in each 5 mL)**

Acetaminophen 160 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if your child has**

liver disease

**Ask a doctor or pharmacist before use if your child is**

taking the blood thinning drug warfarin

**When using this product**

**do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

<b>Weight (lb)</b>	<b>Age (yr)</b>	<b>Dose (mL)*</b>
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL

72-95	11 years	15 mL
-------	----------	-------

\*or as directed by a doctor

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

### **Other information**

- **each 5 mL contains:** sodium 3 mg
- store at 20-25°C (68-77°F)
- **do not use if printed neckband is broken or missing**

### **Inactive ingredients**

anhydrous citric acid, calcium sulfate, carrageenan, FD&C red #40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

### **Questions or comments?**

**1-800-719-9260**

### **Principal Display Panel**

GOOD SENSE®

Ages 2 - 11 Years

Children's Pain & Fever

Pain Reliever/Fever Reducer

Acetaminophen 160 mg per 5 mL

Oral Suspension

Fast, effective relief of children's pain and fever

Alcohol Free

Aspirin Free

Ibuprofen Free

Paraben Free

Cherry Flavor

Compare to active ingredient of Children's Tylenol® Oral Suspension

4 FL OZ (118 mL)



GOODSENSE.

Children's  
**Pain & Fever**  
Cherry Flavor

NDC 0113-0608-26

GOODSENSE®

Ages 2 - 11 Years

Children's  
**Pain & Fever**

Pain Reliever/Fever Reducer  
Acetaminophen 160 mg per 5 mL

Oral Suspension

Fast, effective relief of  
children's pain and fever



Compare to active ingredient of  
Children's Tylenol® Oral Suspension

4 FL OZ (118 mL)



17520 C2 C11

GOODSENSE®

Ages 2 - 11 Years

Children's  
**Pain & Fever**

DIRECTIONS:  
STEP 1.  
Shake well  
before using



STEP 2.  
Remove child  
protective  
cap



STEP 3.  
Squeeze your  
child's dose  
into the  
dosing cup



IMPORTANT:  
Please see dosing chart or ask  
a doctor for your child's correct dose.

**Drug Facts**

**Active ingredient** Purpose  
(in each 5 mL)

Acetaminophen 160 mg.....Pain reliever/fever reducer

**Uses temporarily:**

- reduces fever
- relieves minor aches and pains due to:
  - the common cold
  - flu
  - headache
  - sore throat
  - toothache

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount.

■ with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 5 days
  - fever gets worse or lasts more than 3 days
  - new symptoms occur
  - redness or swelling is present
- These could be signs of a serious condition.

Keep out of reach of children.  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

\*or as directed by a doctor

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

**Other information**

- each 5 mL contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- do not use if printed neckband is broken or missing

**Inactive ingredients**

anhydrous citric acid, calcium sulfate, carrageenan, FD&C red #40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments? 1-800-719-9260

Gluten Free

Distributed By  
**Perrigo®**  
Allegan, MI 49010



**GOOD SENSE CHILDRENS PAIN AND FEVER**

acetaminophen suspension

**Product Information**

**Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:0113-0608

Route of Administration ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WAT0DDB505)	
CARRAGEENAN (UNII: 5C69YCD2YJ)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)	

### Product Characteristics

Color	RED (opaque)	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-0608-26	1 in 1 CARTON	02/15/2016	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0113-0608-28	1 in 1 CARTON	05/28/2013	05/28/2013
2		148 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/28/2013	

**Labeler** - L. Perrigo Company (006013346)

Revised: 5/2024

L. Perrigo Company