# APAP- acetaminophen tablet Advance Pharmaceutical Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

## ACETAMINOPHEN USP 500 mg

# **Active Ingredient**

(in each tablet)

Acetaminophen 500 mg

# Purpose

Pain Reliever / Fever Reducer

#### Uses

temporarily relieves minor aches and pains due to

- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

# Warnings

**Liver warning:** this product contains acetaminophen. The maximum daily dose of this product is 6 tablets (3,000 mg) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

#### do not use

- with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are allergic to acetaminophen or any of the inactive ingredients in this product

## Ask a doctor before use if the you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

## Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

• these could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

## Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

- Do not take more than directed (see over dose warning) adults & children 12 years and over:
- take 2 tablets every 6 hours while symptoms last
- do not take more than 6 tablets in 24 hours, unless directed by a doctor
- do not take for more than 10 days unless directed by a doctor
- children under 12 years : ask a doctor

#### Other Information

store at 15-30 °C (59-86 °F)

For Bulk package: This is a bulk package, dispense contents with a child-resistant closure in a tight, light resistant container as defined in the USP.

## **Inactive Ingredients**

polyvinylpyrrolidone, pregelatinized corn starch, sodium starch glycolate, stearic acid


## **Questions or Comments**

**Call 631-981-4600** 8.30 am- 4.30 pm ET, Monday-Friday

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Manufactured by: Advance Pharmaceutical, Inc. Holtsville, NY 11742

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL



# **Drug Facts**

Active ingredient (in each tablet)

**Purpose** 

Acetaminophen 500 mg ...... Pain reliever/Fever reducer

## Uses

- temporarily relieves minor aches and pains due to:
  - headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold toothache
- premenstrual and menstrual cramps
- m temporarily reduces fever

# Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 6 tablets (3,000 mg) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Drug Facts continued on back of label

#### TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

\*Advance Pharmaceutical, Inc. is not affiliated with the owner of the trademark Extra Strength Tylenol®.

Manufactured by: Advance Pharmaceutical Inc. Holtsville, NY 11742



Lot No.: Exp. Date:

# Drug Facts (continued)

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

# Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- m new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

# Directions on do not take more than directed (see overdose warning)

adults & children 12 years and over:

- take 2 tablets every 6 hours while symptoms last
- do not take more than 6 tablets in 24 hours, unless directed by a doctor
- do not take for more than 10 days unless directed by a doctor

children under 12 years: ask a doctor

# Other information

■ store at 15°to 30°C (59° to 86°F)

Inactive ingredients polyvinylpyrrolidone, pregelatinized corn starch, sodium starch glycolate, stearic acid

Questions or comments? call 631-981-4600, 8:30 am - 4:30 pm ET, Monday - Friday



NDC 17714-013-10

Compare to active ingredient in Extra Strength TYLENOL®

See New Warmings Information

Acetaminophen USP, 500 mg

Pain Reliever - Fever Reducer

# Drug Facts

Active ingredient (in each tablet)

Purpose

Acetaminophen 500 mg ..

Pain reliever/Fever reducer

#### Uses

■ temporarily relieves minor aches and pains due to:

- m headache
- muscular aches
- backache
   toothache
- minor pain of arthritis the common cold
- m premenstrual and menstrual cramps
- temporarily reduces fever

#### Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 6 tablets (3,000 mg) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

mif you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if pain gets worse or lasts more than 10 days

- fever gets worse or lasts more than 3 days
- new symptoms occur redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any sions or symptoms.

Directions m do not take more than directed (see overdose warning)

- adults & children 12 years and over: # take 2 tablets every 6 hours while symptoms last
- do not take more than 6 tablets in 24 hours, unless directed by a doctor
- onot take for more than 10 days unless directed by a doctor

children under 12 years: mask a doctor

Other information store at 15° to 30°C (59° to 86°F)

This is a bulk package. Dispense contents with a child-resistant closure in a tight, light-resistant container as defined in the USP.

Inactive ingredients polyvinylpyrrolidone, pregelatinized corn starch, sodium starch glycolate, stearic acid

Questions or comments?

call 631-981-4600, 8:30 am - 4:30 pm ET, Monday - Friday

#### THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

# TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

\*Advance Pharmaceutical, Inc. is not affiliated with the owner of the trademark Extra Strength Tylenol®,

Lot No .:

Exp. Date:



Manufactured by: Advance Pharmaceutical, Inc. Holtsville, NY 11742

AIII

NDC: 17714-013-01 - 100 COUNT

NDC: 17714-013-10 - 1000 COUNT

## **APAP**

acetaminophen tablet

-				T (			. •	
v	חשי	<i>a</i> 11	Ct	Int	nγ	ma	tio	m
1	ΙU	uu	LLL	Inf	UΙ	ша	uu	ш

Product Type HUMAN OTC DRUG Item Code (Source) NDC:17714-013

Route of Administration ORAL

## **Active Ingredient/Active Moiety**

I	Ingredient Name	Basis of Strength	Strength
l	ACETAMINO PHEN (UNII: 36209 ITL9D) (ACETAMINO PHEN - UNII: 36209 ITL9D)	ACETAMINOPHEN	500 mg

## **Inactive Ingredients**

Ingredient Name	Strength	
PO VIDO NE (UNII: FZ989 GH9 4E)		
STARCH, CORN (UNII: O8232NY3SJ)		
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)		
STEARIC ACID (UNII: 4ELV7Z65AP)		

#### **Product Characteristics**

Color	white	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	AP;013
Contains			

Packaging

	Τ,	1 uchuş ing					
	#	# Item Code Package Description		<b>Marketing Start Date</b>	<b>Marketing End Date</b>		
	1	NDC:17714-013-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/05/1989			
1	2	NDC:17714-013-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	02/05/1989			

# **Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	02/05/1989	

# Labeler - Advance Pharmaceutical Inc. (078301063)

# **Registrant -** Advance Pharmaceutical Inc. (078301063)

Establishment					
Name	Address	ID/FEI	Business Operations		
Advance Pharmaceutical Inc.		078301063	manufacture(17714-013)		

Revised: 12/2017 Advance Pharmaceutical Inc.