

**EQUATE IBUPROFEN- ibuprofen tablet, chewable**  
**Wal-Mart Stores Inc**

-----  
**Wal-Mart Children's Ibuprofen Tablets, 100 mg Drug Facts**

**Active ingredient (in each chewable tablet)**

Ibuprofen 100 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more

than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

### **Do not use**

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

### **Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- mouth or throat burning may occur; give with food or water
- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
  - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days

- redness or swelling is present in the painful area
- any new symptoms appear

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- **chew or crush tablets completely before swallowing**
- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**

<b>Dosing Chart</b>		
<b>Weight (lb)</b>	<b>Age (yr)</b>	<b>Tablets</b>
under 24	under 2	ask a doctor
24-35	2-3	1
36-47	4-5	1 ½
48-59	6-8	2
60-71	9-10	2 ½
72-95	11	3

**Other information**

- phenylketonurics: contains phenylalanine 6 mg per tablet
- store between 20-25°C (68-77°F)
- **do not use if printed seal under cap is broken or missing**

**Inactive ingredients**

acesulfame potassium, ammonium glycyrrhizin, aspartame, carnauba wax, croscarmellose sodium, D&C red no. 27 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, mannitol, natural and artificial flavors, silicon dioxide, sodium lauryl sulfate, soybean oil, succinic acid

**Questions or comments?**

**1-888-287-1915**

**Principal Display Panel**

equate™

Compare to Children's Motrin® active ingredient

FOR AGES 2 TO 11 YEARS

children's

IBUPROFEN

Chewable Tablets, 100 mg

LASTS UP TO 8 HOURS

Pain Reliever/Fever Reducer (NSAID)

Actual Size

Chewables

Grape Flavor

Chew or crush tablets completely before swallowing

24 CHEWABLE TABLETS

how2recycle.info  
 BOTTLE  
 PAPER PLASTIC  
 KEEP CARTON FOR REFERENCE  
 Satisfaction guaranteed - Or we'll replace it or give you your money back. For questions or comments please call 1-888-287-1915.  
 DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716  
 \*\*\*This product is not manufactured or distributed by Kenvee Inc., owner of the registered trademark Motrin®.

**Drug Facts (continued)**  
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions** ■ chew or crush tablets completely before swallowing ■ this product does not contain directions or complete warnings for adult use ■ do not give more than directed ■ find right dose on chart below. If possible, use weight to dose, otherwise use age. ■ if needed, repeat dose every 6-8 hours ■ do not use more than 4 times a day

Weight (lb)	Age (yr)	Tablets
under 24	under 2	ask a doctor
24-35	2-3	1
36-47	4-5	1½
48-59	6-8	2
60-71	9-10	2½
72-85	11	3

**Other information** ■ phenylethanamine: contains phenylethanamine 6 mg per tablet ■ store between 20-25°C (68-77°F) ■ do not use if printed seal under cap is broken or missing

**Inactive ingredients** acacia, caffeine potassium, ammonium glycyrrhizate, aspartame, carnauba wax, croscarmellose sodium, D&C red no. 27 aluminum lake, FD&C blue no. 1, aluminum lake, hypromellose, magnesium stearate, mannitol, natural and artificial flavors, silicon dioxide, sodium lauryl sulfate, soybean oil, succinic acid

**Questions or comments? 1-888-287-1915**

**Drug Facts (continued)**  
 Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Do not use**  
 ■ if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer  
 ■ right before or after heart surgery

**Ask a doctor before use if**  
 ■ stomach bleeding  
 ■ warning applies to your child ■ child has a history of stomach problems, such as heartburn ■ child has problems or serious side effects from taking pain relievers or fever reducers ■ child has not been drinking fluids ■ child has lost a lot of fluid due to vomiting or diarrhea ■ child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke ■ child has asthma ■ child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**  
 ■ under a doctor's care for any serious condition ■ taking any other drug

**When using this product**  
 ■ mouth or throat burning may occur; give with food or water ■ take with food or milk, if stomach upset occurs

**Stop use and ask a doctor if**  
 ■ child experiences any of the following signs of stomach bleeding:  
 ■ feels faint ■ vomits blood ■ has bloody or black stools ■ has stomach pain that does not get better  
 ■ child has symptoms of heart problems or stroke:  
 ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ the child does not get any relief within first day (24 hours) of treatment ■ fever or pain gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

164152  
 \* 6 81131 01659 9  
 52 162 2E C6

**Important: Read all product information before using. Keep this box for important information. This product is for children only.**

**Drug Facts**  
**Active ingredient**  
 (in each chewable tablet)  
 ibuprofen 100 mg (NSAID) ... pain reliever/fever reducer \*nonsteroidal anti-inflammatory drug

**Purposes**  
 ■ reduces fever  
 ■ relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

**Uses temporarily:**  
 ■ reduces fever  
 ■ relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

**Warnings**  
**Allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin redness ■ rash ■ blisters If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:  
 ■ has had stomach ulcers or bleeding problems ■ takes a blood thinning (anticoagulant) or steroid drug ■ takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, malaise, and vomiting may be serious. Consult doctor promptly.

NDC 49035-521-62

**equate™**

FOR AGES 2 to 11 YEARS

**children's**  
**IBUPROFEN**  
 Chewable Tablets, 100 mg

☐ Pain Reliever/Fever Reducer (NSAID) Actual Size

Chewables Grape Flavor Chew or crush tablets completely before swallowing

**24 CHEWABLE TABLETS**

Compare to Children's Motrin® active ingredients  
 LASTS UP TO 8 HOURS

EQUATE IBUPROFEN			
ibuprofen tablet, chewable			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49035-521
Route of Administration	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>AMMONIUM GLYCYRRHIZATE</b> (UNII: 3VRD35U26C)	
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>D&amp;C RED NO. 27 ALUMINUM LAKE</b> (UNII: ZK64F7XSTX)	
<b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>SOYBEAN OIL</b> (UNII: 241ATL177A)	
<b>SUCCINIC ACID</b> (UNII: AB6MNQ6J6L)	

**Product Characteristics**

<b>Color</b>	PURPLE (Lavender)	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	12mm
<b>Flavor</b>	GRAPE	<b>Imprint Code</b>	L521
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49035-521-62	1 in 1 CARTON	10/25/2011	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:49035-521-67	2 in 1 CARTON	10/25/2011	
2		24 in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076359	10/25/2011	

