## DOUBLE DINE BARRIER NPE FREE- iodine and chlorhexidine gluconate liquid IBA

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

-----

### **Double Dine Barrier NPE Free**

### **ACTIVE INGREDIENTS**

**ACTIVE INGREDIENT:** 

CHLORHEXIDINE GLUCONATE	0.50%
IODINE	0.50%
(PROVIDES A Full .5% AVAILABLE IODINE AS USED)	

### **INERT INGREDIENTS**

GLYCERIN (AS A HUMECTANT AND SKIN CONDITION	NER)2.00%
OTHER INERTS	97.00%
(INCLUDES IODINE STABILIZERS, BUFFERING AGEN	ITS AND SURFACTANTS)
TOTAL:	100.00%

THIS PRODUCT CONTAINS HIGHLY EFFECTIVE LEVELS OF FREE IODINE FOR FAST KILL OF MASTITIS CAUSING ORGANISMS.

#### INDICATIONS AND USAGE

### GENERAL RECOMMENDATIONS

- 1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
- 2. When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
- 3. Do not use this product for cleaning or sanitizing equipment
- 4. Wash Teat Dip cups after each milking.

# RECOMMENDED PROCEDURES FOR TEAT SANITATION POST-MILKING (POST-DIPPING)

Immediately after milking, dip or spray all teats full length with DOUBLE DINE NPE FREE BARRIER TEAT DIP. Ensure good coverage of all sides of the teats and teat orifices.

### WARNING

DO NOT DILUTE THIS PRODUCT.

DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.

ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO

WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS. PROTECT THIS PRODUCT FROM FREEZING.

CAUTION: KEEP OUT OF REACH OF CHILDREN.
AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

### FIRST AID PROCEDURES

**EYE CONTACT:** IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

**SKIN CONTACT:** FOR READY TO USE PRODUCTS, WASH ANY SENSITIVE AREAS. **INGESTION:**DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION FOR MEDICAL EMERGENCIES CALL 1-800-424-9300

**Double Dine Barrier - NPE Free** 



### **DOUBLE DINE BARRIER NPE FREE**

iodine and chlorhexidine gluconate liquid

Product Information				
Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:29983-1073	
Route of Administration	TOPICAL			

Active Ingredient/Active Moiety				
Ingredient Name	<b>Basis of Strength</b>	Strength		
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	0.5 kg in 100 kg		
CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII: R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	0.5 kg in 100 kg		

P	ackaging			
#	Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:29983-1073-9	1064.03 kg in 1 CONTAINER		

2 NDC:29983-1073-8	212.81 kg in 1 DRUM		
Marketing In	formation		
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug		03/31/2023	

## **Labeler -** IBA (019494160)

## **Registrant -** IBA (019494160)

Establishment			
Name	Address	ID/FEI	<b>Business Operations</b>
Morgan Gallacher Inc. DBA, Custom Chemical Formulators Inc.		028311595	manufacture, api manufacture, pack

Revised: 2/2024 IBA