

**RI MOX- aluminum hydroxide, magnesium hydroxide, simethicone liquid**  
**Rij Pharmaceutical Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active ingredients (in each 5 mL) (One teaspoonful)**

Aluminum Hydroxide (Equi. to Dried Gel, USP) 200 mg

Magnesium Hydroxide 200 mg

Simethicone 20 mg

**Purpose**

Antacid

Antigas

**Uses**

relieves

- heartburn
- acid indigestion
- sour stomach
- associated symptoms of gas and upset stomach
- overindulgence of food and drink

**Warnings**

**Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Antacids may interact with certain prescription drugs.

**Stop use and ask a doctor if**

symptoms last for more than 2 weeks

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center immediately.

**Directions**

- **shake well before each use**

- Adults and children 12 years and older, take 2 to 4 teaspoons four times a day or as directed by a physician
- do not take more than 16 teaspoonsfuls in 24 or use the maximum dosage for more than 2 weeks
- Children under 12 years: consult a physician

### **Other information**

- TAMPER EVIDENT: DO NOT USE IF BREAKAWAY BAND ON CAP IS BROKEN OR MISSING
- Store at room temperature 15° - 30°C (59° - 86°F)
- keep tightly closed
- protect from freezing
- each teaspoon (5 mL) contains: magnesium 83 mg
- each teaspoon (5 mL) contains: sodium 1 mg
- does not meet USP requirements for preservative effectiveness

### **Inactive ingredients**

flavors, hypromellose, parabens, saccharin, sodium hypochlorite, sorbitol, water and xanthan gum

### **Principal Display Panel**



NDC 53807-502-12



# RI-MOX

**ANTACID / ANTIGAS**

**Fast Relief of**

- Heartburn
- Acid Indigestion

**MINT FLAVOR**



**12 FL. OZ. (355 ML)**

# RI-MOX

## Drug Facts

### Active Ingredients

(In each 5 ml) (One teaspoonful)

Purpose

Aluminum Hydroxide

(Equi. to Dried Gel, USP) 200 mg .....Antacid

Magnesium Hydroxide 200 mg .....Antacid

Simethicone 20 mg .....Antigas

**Uses** relieves ■ heartburn ■ acid indigestion ■ sour stomach  
■ associated symptoms of gas and upset stomach  
■ overindulgence of food and drink

## Warnings

**Ask a doctor before use if you have**

■ kidney disease ■ a magnesium-restricted diet

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Antacids may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ symptoms last more than 2 weeks

**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center immediately.

## Directions ■ shake well before each use

■ Adults and children 12 years and older, take 2 to 4 teaspoons four times a day or as directed by a physician

■ do not take more than 16 teaspoonfuls in 24-hours or use the maximum dosage for more than 2 weeks

■ Children under 12 years: consult a physician

## Other information

■ **TAMPER EVIDENT: DO NOT USE IF BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.**

■ store at room temperature 15°-30°C (59°-86°F) ■ keep tightly closed  
■ protect from freezing

■ each teaspoon (5 mL) contains: magnesium 83 mg

■ each teaspoon (5 mL) contains: sodium 1 mg

■ does not meet USP requirements for preservative effectiveness

**Inactive ingredients:** flavors, hypromellose, parabens, saccharin, sodium hypochlorite, sorbitol, water and xanthan gum.

\*\*\*This product is not manufactured or distributed by the owner of the registered trademark Maalox®.

RIJ PHARMACEUTICAL CORP.  
40 Commercial Avenue  
Middletown, NY 10941

Rev.06/12



N 3 53807-502-12 4

aluminum hydroxide, magnesium hydroxide, simethicone liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:53807-502
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
ALUMINUM HYDRO XIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	200 mg in 5 mL
MAGNESIUM HYDRO XIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	200 mg in 5 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	20 mg in 5 mL

### Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
HYPROMELLOSES (UNII: 3NXW29V3WO)	
BUTYL PARABEN (UNII: 3QP1I03FV8)	
SACCHARIN (UNII: FST467XS7D)	
SODIUM HYPOCHLORITE (UNII: DY38VHM5OD)	
SORBITOL (UNII: 506T60A25R)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0K00R)	
XANTHAN GUM (UNII: TTV12P4NEE)	

### Product Characteristics

<b>Color</b>	WHITE (opaque)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	PEPPERMINT	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53807-502-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/16/1999	
2	NDC:53807-502-28	3785 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/16/1999	

### Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph final	part332	03/16/1999	

**Labeler** - Rij Pharmaceutical Corporation (144679156)

**Establishment**

Name	Address	ID/FEI	Business Operations
Rij Pharmaceutical Corporation		144679156	manufacture(53807-502)

Revised: 4/2018

Rij Pharmaceutical Corporation