

NIGHT TIME SLEEP-AID- diphenhydramine hcl liquid
Discount Drug Mart

Drug Facts

Active ingredients (in each 30 mL dose cup or 2 tablespoons)

Diphenhydramine HCL 50 mg

Purpose

Nighttime sleep-aid

Keep out of reach of children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Uses

- reduces time to fall asleep if you have difficulty falling asleep
- for relief of occasional sleeplessness

Warnings

DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN

Failure to follow these warnings could result in serious consequences.

Do not use

- for children under 12 years of age.
- with any other product containing diphenhydramine, even one used on skin
- with any other drugs that cause drowsiness such as antihistamines and nighttime cough, cold/flu products

Ask a doctor before use if you have

- heart disease
- trouble urinating due to an enlarged prostate gland
- glaucoma
- a breathing problem such as asthma, emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers or any other sleep aid

When using this product

- **do not use more than directed**
- avoid alcoholic beverages and other drugs that cause drowsiness
- drowsiness will occur
- be careful driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.

If pregnant or breast feeding,

ask a health professional before use.

Directions

- take only as recommended
- use dosage cup or tablespoon

Age	Dose
Adults and children 12 years and over	One dose = 2 tablespoons (30 mL) at bedtime if needed or as directed by a doctor
Children under 12 years	Do not use

Other information

- **each 30 mL dose (2 tablespoons) contains:** sodium 23 mg
- dosage cup provided
- store at room temperature

Inactive ingredients

citric acid, FD and C Blue # 1, FD and C Red # 40, flavor, high fructose corn syrup, poloxamer 407, proplene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin

Questions?

Call weekdays from 9:30 AM to 4:30 PM EST at **1-877-798-5944**

DISCOUNT drug mart FOOD FAIR NIGHT TIME SLEEP-AID product label

* COMPARE TO THE ACTIVE INGREDIENT IN ZzzQuil™ NIGHTTIME SLEEP-AID

DISCOUNT

drug mart

FOOD FAIR

NIGHT TIME SLEEP-AID

Diphenhydramine HCL

Non-Habit Forming

Warming Berry Flavored Liquid

Not for treating Cold or Flu

12 FL OZ (354 mL)

* This product is not manufactured or distributed by Proctor and Gamble owner of the registered trademark ZzzQuil™

LR-055

SATISFACTION GUARANTEED

IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO THE STORE WHERE PURCHASED. IF UNABLE TO RETURN TO THE STORE, SEND REASON FOR DISSATISFACTION , NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE MEDINNA, OHIO 44256

Drug Facts (continued)

Directions

- take only as recommended
- use dosage cup or tablespoon

Age	Dose
Adults & children 12 years and over	One dose = 2 tablespoons (30 mL) at bed time if needed or as directed by a doctor
Children under 12 years	Do not use

Other information

- each 30 mL dose (2 tablespoons) contains: sodium 23 mg
- dosage cup provided
- store at room temperature

Inactive ingredients

citric acid, FD&C Blue #1, FD&C Red #40, flavor, high fructose corn syrup, poloxamer 407, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

*This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark ZzzQuil™.

LR-055

SATISFACTION GUARANTEED
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256



* COMPARE TO THE ACTIVE INGREDIENT IN ZzzQuil™ NIGHTTIME SLEEP-AID

NIGHT TIME SLEEP-AID

Diphenhydramine HCl

Non-Habit Forming

Warming Berry Flavored Liquid
Not for treating Cold or Flu

12 FL OZ (354 mL)

DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN
Failure to follow these warnings could result in serious consequences

Drug Facts

Active ingredients	Purpose
Diphenhydramine HCl 50 mg	Nighttime sleep-aid

Uses

- reduces time to fall asleep if you have difficulty falling asleep
- for relief of occasional sleeplessness

Warnings

Do not use

- for children under 12 years of age.
- with any other product containing diphenhydramine, even one used on skin
- with any other drugs that cause drowsiness such as antihistamines and nighttime cough, cold/flu products

Ask a doctor before use if you have

- heart disease
- trouble urinating due to an enlarged prostate gland
- glaucoma
- a breathing problem such as asthma, emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers or any other sleep aid

When using this product

- do not use more than directed
- avoid alcoholic beverages and other drugs that cause drowsiness
- drowsiness will occur
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts Failure to follow these warnings could result in serious consequences

Active ingredients	Purpose
Diphenhydramine HCl 50 mg	Nighttime sleep-aid

Uses

- reduces time to fall asleep if you have difficulty falling asleep
- for relief of occasional sleeplessness

Warnings

Do not use

- for children under 12 years of age.
- with any other product containing diphenhydramine, even one used on skin
- with any other drugs that cause drowsiness such as antihistamines and nighttime cough, cold/flu products

Ask a doctor before use if you have

- heart disease
- trouble urinating due to an enlarged prostate gland
- glaucoma
- a breathing problem such as asthma, emphysema or chronic bronchitis

Peel Here



* COMPARE TO THE ACTIVE INGREDIENT IN ZzzQuil™ NIGHTTIME SLEEP-AID

NIGHT TIME SLEEP-AID

Diphenhydramine HCl

Non-Habit Forming

Warming Berry Flavored Liquid
Not for treating Cold or Flu

6 FL OZ (177 mL)

DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN



*This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark ZzzQuil™.

LR-054

SATISFACTION GUARANTEED
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

EXP:

LOT:

NIGHT TIME SLEEP-AID

Diphenhydramine HCl

Non-Habit Forming

Warming Berry Flavored Liquid
Not for treating Cold or Flu

<p>Drug Facts (continued)</p> <p>Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers or any other sleep aid</p> <p>When using this product</p> <ul style="list-style-type: none"> ■ do not use more than directed ■ avoid alcoholic beverages and other drugs that cause drowsiness ■ drowsiness will occur ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness. <p>If pregnant or breast feeding, ask a health professional before use.</p>

<p>Drug Facts (continued)</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ take only as recommended ■ use dosage cup or tablespoon <table border="1"> <thead> <tr> <th>Age</th> <th>Dose</th> </tr> </thead> <tbody> <tr> <td>Adults & children 12 years and over</td> <td>One dose = 2 tablespoons (30 mL) at bed time if needed or as directed by a doctor</td> </tr> <tr> <td>Children under 12 years</td> <td>Do not use</td> </tr> </tbody> </table>	Age	Dose	Adults & children 12 years and over	One dose = 2 tablespoons (30 mL) at bed time if needed or as directed by a doctor	Children under 12 years	Do not use
Age	Dose					
Adults & children 12 years and over	One dose = 2 tablespoons (30 mL) at bed time if needed or as directed by a doctor					
Children under 12 years	Do not use					

<p>Drug Facts (continued)</p> <p>Other information</p> <ul style="list-style-type: none"> ■ each 30 mL dose (2 tablespoons) contains: sodium 23 mg ■ dosage cup provided ■ store at room temperature <p>Inactive ingredients</p> <p>citric acid, FD&C Blue #1, FD&C Red #40, flavor, high fructose corn syrup, poloxamer 407, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin</p> <p>Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944</p>
--

res

Drug Facts (continued)

Directions

- take only as recommended
- use dosage cup or tablespoon

Age	Dose
Adults & children 12 years and over	One dose = 2 tablespoons (30 mL) at bed time if needed or as directed by a doctor
Children under 12 years	Do not use

Other information

- each 30 mL dose (2 tablespoons) contains: sodium 23 mg
- dosage cup provided
- store at room temperature

Inactive ingredients

citric acid, FD&C Blue #1, FD&C Red #40, flavor, high fructose corn syrup, poloxamer 407, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

*This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark ZzzQuil™.

LR-055

SATISFACTION GUARANTEED
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256



* COMPARE TO THE ACTIVE INGREDIENT IN ZzzQuil™ NIGHTTIME SLEEP-AID

NIGHT TIME SLEEP-AID

Diphenhydramine HCl

Non-Habit Forming

Warming Berry Flavored Liquid
Not for treating Cold or Flu

12 FL OZ (354 mL)

DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN
Failure to follow these warnings could result in serious consequences

Drug Facts

Active ingredients	Purpose
(in each 30 mL dose cup or 2 tablespoons) Diphenhydramine HCl 50 mg	Nighttime sleep-aid

Uses

- reduces time to fall asleep if you have difficulty falling asleep
- for relief of occasional sleeplessness

Warnings

Do not use

- for children under 12 years of age.
- with any other product containing diphenhydramine, even one used on skin
- with any other drugs that cause drowsiness such as antihistamines and nighttime cough, cold/flu products

Ask a doctor before use if you have

- heart disease
- trouble urinating due to an enlarged prostate gland
- glaucoma
- a breathing problem such as asthma, emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers or any other sleep aid

When using this product

- do not use more than directed
- avoid alcoholic beverages and other drugs that cause drowsiness
- drowsiness will occur
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts Failure to follow these warnings could result in serious consequences

Active ingredients	Purpose
(in each 30 mL dose cup or 2 tablespoons) Diphenhydramine HCl 50 mg	Nighttime sleep-aid

Uses

- reduces time to fall asleep if you have difficulty falling asleep
- for relief of occasional sleeplessness

Warnings

Do not use

- for children under 12 years of age.
- with any other product containing diphenhydramine, even one used on skin
- with any other drugs that cause drowsiness such as antihistamines and nighttime cough, cold/flu products

Ask a doctor before use if you have

- heart disease
- trouble urinating due to an enlarged prostate gland
- glaucoma
- a breathing problem such as asthma, emphysema or chronic bronchitis

Peel Here



* COMPARE TO THE ACTIVE INGREDIENT IN ZzzQuil™ NIGHTTIME SLEEP-AID

NIGHT TIME SLEEP-AID

Diphenhydramine HCl

Non-Habit Forming

Warming Berry Flavored Liquid
Not for treating Cold or Flu

6 FL OZ (177 mL)

DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN



*This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark ZzzQuil™.

LR-054

SATISFACTION GUARANTEED
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

EXP:

LOT:

NIGHT TIME SLEEP-AID

Diphenhydramine HCl
Non-Habit Forming

Warming Berry Flavored Liquid
Not for treating Cold or Flu

<p>Drug Facts (continued)</p> <p>Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers or any other sleep aid</p> <p>When using this product</p> <ul style="list-style-type: none"> do not use more than directed avoid alcoholic beverages and other drugs that cause drowsiness drowsiness will occur be careful when driving a motor vehicle or operating machinery alcohol, sedatives, and tranquilizers may increase drowsiness <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness. <p>If pregnant or breast feeding, ask a health professional before use.</p>

<p>Drug Facts (continued)</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> take only as recommended use dosage cup or tablespoon <table border="1"> <thead> <tr> <th>Age</th> <th>Dose</th> </tr> </thead> <tbody> <tr> <td>Adults & children 12 years and over</td> <td>One dose = 2 tablespoons (30 mL) at bed time if needed or as directed by a doctor</td> </tr> <tr> <td>Children under 12 years</td> <td>Do not use</td> </tr> </tbody> </table>	Age	Dose	Adults & children 12 years and over	One dose = 2 tablespoons (30 mL) at bed time if needed or as directed by a doctor	Children under 12 years	Do not use
Age	Dose					
Adults & children 12 years and over	One dose = 2 tablespoons (30 mL) at bed time if needed or as directed by a doctor					
Children under 12 years	Do not use					

<p>Drug Facts (continued)</p> <p>Other information</p> <ul style="list-style-type: none"> each 30 mL dose (2 tablespoons) contains: sodium 23 mg dosage cup provided store at room temperature <p>Inactive ingredients</p> <p>citric acid, FD&C Blue #1, FD&C Red #40, flavor, high fructose corn syrup, poloxamer 407, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin</p> <p>Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944</p>
--

NIGHT TIME SLEEP-AID

diphenhydramine hcl liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53943-522
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	50 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLOXAMER 407 (UNII: TUF2IVW3M2)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	BERRY (BERRY)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-522-25	117 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/10/2013	
2	NDC:53943-522-28	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/10/2013	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M010	12/10/2013	

Labeler - Discount Drug Mart (047741335)

Revised: 12/2023

Discount Drug Mart