

**TUSSIN LONG ACTING COUGH READYINCASE- dextromethorphan hbr liquid
P & L Development, LLC**

Drug Facts

Active ingredient (in each 10 mL)

Dextromethorphan HBr, 30 mg

Purpose

Cough suppressant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema

Stop use and ask a doctor if

cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.

- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age
- adults and children 12 years and over: 10 mL every 6 to 8 hours
- children under 12 years: do not use

Other information

- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, dextrose, ethyl alcohol, FD&C red 40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate

Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal Display Panel

Compare to active ingredient in **Robitussin® Lingerin Cold Long-Acting Cough***

Tussin

Long-acting

Cough

Dextromethorphan HBr, 30 mg

cough suppressant

Relieves cough for up to 8 hours

For ages 12 & over

alcohol 1.4%

non-drowsy

Dosing Cup Included

FL OZ (mL)

*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributors of Robitussin® Lingerin Cold Long-Acting Cough.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

Manufactured by:

PL Developments

11865 S. Alamea St

Package Label

<p>Drug Facts (continued) saccharin sodium, sodium benzoate</p> <p>Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST</p>	 <p>Compare to active ingredient in Robitussin® Lingerin Cold Long-Acting Cough*</p> <p>NDC 49580-0383-4</p>	 <p>Compare to active ingredient in Robitussin® Lingerin Cold Long-Acting Cough*</p> <p>NDC 49580-0383-4</p>	<p>Drug Facts</p> <table border="1"> <thead> <tr> <th>Active ingredient (in each 10 mL)</th> <th>Purpose</th> </tr> </thead> <tbody> <tr> <td>Dextromethorphan HBr 30 mg</td> <td>Cough suppressant</td> </tr> </tbody> </table> <p>Use</p> <ul style="list-style-type: none"> temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold <p>Warnings</p> <p>Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> cough that occurs with too much phlegm (mucus) cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema <p>Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p> <p>Directions</p> <ul style="list-style-type: none"> do not take more than 4 doses in any 24-hour period measure only with dosing cup provided. Do not use any other dosing device. Keep dosing cup with product mL = milliliter this adult product is not intended for use in children under 12 years of age adults and children 12 years and over: 10 mL every 6 to 8 hours children under 12 years of age: do not use <p>Other information</p> <ul style="list-style-type: none"> store between 20-25°C (68-77°F). Do not refrigerate. <p>Inactive ingredients citric acid, dextrose, ethyl alcohol, FD&C red #40, flavor, glycerin, high fructose corn syrup, menthol, purified water, ▶</p>	Active ingredient (in each 10 mL)	Purpose	Dextromethorphan HBr 30 mg	Cough suppressant
Active ingredient (in each 10 mL)	Purpose						
Dextromethorphan HBr 30 mg	Cough suppressant						
<p>TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.</p>	 <p>dextromethorphan HBr, 30 mg cough suppressant</p>						
 <p>Dosing Cup Included</p>	<p>relieves cough for up to 8 hours</p> <p>for ages 12 & over alcohol 1.4% non-drowsy</p> <p>4 fl oz (118 mL)</p>						
 <p>Manufactured by: PL Developments 11865 S. Alameda St Lynwood, CA 90262</p>	 <p>7 15256 71104 3</p> <p>PLD-H323C FC004060</p> <p>Lot No: Exp. Date:</p>						
<p>PARENTS: Learn about teen medicine abuse www.StopMedicineAbuse.org</p>							

READYINCASE Tussin long-acting cough

TUSSIN LONG ACTING COUGH READYINCASE			
dextromethorphan hbr liquid			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49580-0383
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 10 mL	
Inactive Ingredients				
	Ingredient Name	Strength		
	ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
	FD&C RED NO. 40 (UNII: WZB9127XOA)			
	GLYCERIN (UNII: PDC6A3C0OX)			
	WATER (UNII: 059QF0KO0R)			
	SODIUM BENZOATE (UNII: OJ245FE5EU)			
	MENTHOL (UNII: L7T10EIP3A)			
	ALCOHOL (UNII: 3K9958V90M)			
	DEXTROSE (UNII: IY9XDZ35W2)			
	HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)			
	SACCHARIN SODIUM (UNII: SB8ZUX40TY)			
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49580-0383-4	1 in 1 BOX	08/31/2014	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing Information				
	Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
	OTC Monograph Drug	M012	08/31/2014	

Labeler - P & L Development, LLC (101896231)

Revised: 12/2024

P & L Development, LLC