

NICOTINE- nicotine polacrilex gum, chewing
P & L Development, LLC

Nicotine Polacrilex Gum 2 mg Coated Fruit

Drug Facts

Active ingredient (in each chewing piece)

Nicotine Polacrilex (equal to 2 mg nicotine)

Purpose

Stop smoking aid

Use

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions - 2 mg

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information - 2 mg

- **each piece contains:** calcium 115 mg, sodium 8 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Inactive ingredients - 2 mg

acacia, acesulfame potassium, carnauba wax, FD&C blue #2 lake, FD&C red #40 lake, FD&C yellow #6 lake, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, sucralose, talc, titanium dioxide, xylitol.

Questions or comments?

Call **1-877-753--3935** Monday-Friday 9AM-5PM EST

Principal display panel - 2 mg

*Compare to the active ingredient of Nicorette® Gum

Nicotine Gum

nicotine polacrilex gum USP, 2 mg (nicotine)

stop smoking aid

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE **MORE THAN 30 MINUTES** AFTER WAKING UP

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicotine Polacrilex Gum USP, 4 mg

2 mg

sugar-free

coated fruit flavor

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew **at least 9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use with a Support Program** as directed in the enclosed User's Guide.

- **not for sale to those under 18 years of age**
- **proof of age required**
- **not for sale in vending machines or from any source where proof of age cannot be verified**

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

*This product is not manufactured or distributed by Haleon Group of Companies, distributor of Nicorette® Gum.

Distributed by: **PL Developments**

200 Hicks Street, Westbury, NY 11590

Package label 2 mg

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use With a Support Program** as directed in the enclosed User's Guide.



To remove the gum, tear off single unit.



Peel off backing starting at corner with loose edge.



Push gum through foil.



nicotine gum
 nicotine polacrilex gum
 USP, 2 mg (nicotine)
 stop smoking aid

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE **MORE THAN 30 MINUTES** AFTER WAKING UP.

If you smoke your first cigarette **WITHIN 30 MINUTES** of waking up, use Nicotine Polacrilex Gum USP, 4 mg

2 mg

sugar-free

coated
 fruit
 flavor

160 pieces, 2 mg each

*Compare to the active ingredient in Nicorette® Gum



nicotine gum
 nicotine polacrilex gum USP, 2 mg (nicotine)
 stop smoking aid

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE **MORE THAN 30 MINUTES** AFTER WAKING UP.
 If you smoke your first cigarette **WITHIN 30 MINUTES** of waking up, use Nicotine Polacrilex Gum USP, 4 mg

Wellness Basics - 2mg Nicotine Strengths - 160 Count
 New
 X0045A838X

Keep out of reach of children and pets. Pieces of nicotine gum may have bitter taste. If you have difficulty swallowing or have difficulty chewing, use a chewing gum. Do not use if you are under 18 years of age, use a doctor before use if you are under 18 years of age, or if you are pregnant, nursing, or taking any medication. Do not use if you are taking any medication for depression, anxiety, or other mental health conditions. Do not use if you are taking any medication for high blood pressure, heart disease, or other cardiovascular conditions. Do not use if you are taking any medication for diabetes, or if you have diabetes. Do not use if you are taking any medication for asthma, or if you have asthma. Do not use if you are taking any medication for epilepsy, or if you have epilepsy. Do not use if you are taking any medication for glaucoma, or if you have glaucoma. Do not use if you are taking any medication for high cholesterol, or if you have high cholesterol. Do not use if you are taking any medication for kidney disease, or if you have kidney disease. Do not use if you are taking any medication for liver disease, or if you have liver disease. Do not use if you are taking any medication for thyroid disease, or if you have thyroid disease. Do not use if you are taking any medication for ulcers, or if you have ulcers. Do not use if you are taking any medication for stomach acid, or if you have stomach acid. Do not use if you are taking any medication for allergies, or if you have allergies. Do not use if you are taking any medication for autoimmune disease, or if you have autoimmune disease. Do not use if you are taking any medication for cancer, or if you have cancer. Do not use if you are taking any medication for HIV/AIDS, or if you have HIV/AIDS. Do not use if you are taking any medication for hepatitis, or if you have hepatitis. Do not use if you are taking any medication for tuberculosis, or if you have tuberculosis. Do not use if you are taking any medication for malaria, or if you have malaria. Do not use if you are taking any medication for parasites, or if you have parasites. Do not use if you are taking any medication for HIV/AIDS, or if you have HIV/AIDS. Do not use if you are taking any medication for hepatitis, or if you have hepatitis. Do not use if you are taking any medication for tuberculosis, or if you have tuberculosis. Do not use if you are taking any medication for malaria, or if you have malaria. Do not use if you are taking any medication for parasites, or if you have parasites.

Directions
 If you are under 18 years of age, use a doctor before use. Do not use if you are pregnant, nursing, or taking any medication. Do not use if you are taking any medication for depression, anxiety, or other mental health conditions. Do not use if you are taking any medication for high blood pressure, heart disease, or other cardiovascular conditions. Do not use if you are taking any medication for diabetes, or if you have diabetes. Do not use if you are taking any medication for asthma, or if you have asthma. Do not use if you are taking any medication for epilepsy, or if you have epilepsy. Do not use if you are taking any medication for glaucoma, or if you have glaucoma. Do not use if you are taking any medication for high cholesterol, or if you have high cholesterol. Do not use if you are taking any medication for kidney disease, or if you have kidney disease. Do not use if you are taking any medication for liver disease, or if you have liver disease. Do not use if you are taking any medication for thyroid disease, or if you have thyroid disease. Do not use if you are taking any medication for ulcers, or if you have ulcers. Do not use if you are taking any medication for stomach acid, or if you have stomach acid. Do not use if you are taking any medication for allergies, or if you have allergies. Do not use if you are taking any medication for autoimmune disease, or if you have autoimmune disease. Do not use if you are taking any medication for cancer, or if you have cancer. Do not use if you are taking any medication for HIV/AIDS, or if you have HIV/AIDS. Do not use if you are taking any medication for hepatitis, or if you have hepatitis. Do not use if you are taking any medication for tuberculosis, or if you have tuberculosis. Do not use if you are taking any medication for malaria, or if you have malaria. Do not use if you are taking any medication for parasites, or if you have parasites.

Keep out of reach of children and pets. Pieces of nicotine gum may have bitter taste. If you have difficulty swallowing or have difficulty chewing, use a chewing gum. Do not use if you are under 18 years of age, use a doctor before use if you are under 18 years of age, or if you are pregnant, nursing, or taking any medication. Do not use if you are taking any medication for depression, anxiety, or other mental health conditions. Do not use if you are taking any medication for high blood pressure, heart disease, or other cardiovascular conditions. Do not use if you are taking any medication for diabetes, or if you have diabetes. Do not use if you are taking any medication for asthma, or if you have asthma. Do not use if you are taking any medication for epilepsy, or if you have epilepsy. Do not use if you are taking any medication for glaucoma, or if you have glaucoma. Do not use if you are taking any medication for high cholesterol, or if you have high cholesterol. Do not use if you are taking any medication for kidney disease, or if you have kidney disease. Do not use if you are taking any medication for liver disease, or if you have liver disease. Do not use if you are taking any medication for thyroid disease, or if you have thyroid disease. Do not use if you are taking any medication for ulcers, or if you have ulcers. Do not use if you are taking any medication for stomach acid, or if you have stomach acid. Do not use if you are taking any medication for allergies, or if you have allergies. Do not use if you are taking any medication for autoimmune disease, or if you have autoimmune disease. Do not use if you are taking any medication for cancer, or if you have cancer. Do not use if you are taking any medication for HIV/AIDS, or if you have HIV/AIDS. Do not use if you are taking any medication for hepatitis, or if you have hepatitis. Do not use if you are taking any medication for tuberculosis, or if you have tuberculosis. Do not use if you are taking any medication for malaria, or if you have malaria. Do not use if you are taking any medication for parasites, or if you have parasites.

Warnings
 Do not use if you are pregnant, nursing, or taking any medication. Do not use if you are taking any medication for depression, anxiety, or other mental health conditions. Do not use if you are taking any medication for high blood pressure, heart disease, or other cardiovascular conditions. Do not use if you are taking any medication for diabetes, or if you have diabetes. Do not use if you are taking any medication for asthma, or if you have asthma. Do not use if you are taking any medication for epilepsy, or if you have epilepsy. Do not use if you are taking any medication for glaucoma, or if you have glaucoma. Do not use if you are taking any medication for high cholesterol, or if you have high cholesterol. Do not use if you are taking any medication for kidney disease, or if you have kidney disease. Do not use if you are taking any medication for liver disease, or if you have liver disease. Do not use if you are taking any medication for thyroid disease, or if you have thyroid disease. Do not use if you are taking any medication for ulcers, or if you have ulcers. Do not use if you are taking any medication for stomach acid, or if you have stomach acid. Do not use if you are taking any medication for allergies, or if you have allergies. Do not use if you are taking any medication for autoimmune disease, or if you have autoimmune disease. Do not use if you are taking any medication for cancer, or if you have cancer. Do not use if you are taking any medication for HIV/AIDS, or if you have HIV/AIDS. Do not use if you are taking any medication for hepatitis, or if you have hepatitis. Do not use if you are taking any medication for tuberculosis, or if you have tuberculosis. Do not use if you are taking any medication for malaria, or if you have malaria. Do not use if you are taking any medication for parasites, or if you have parasites.

Questions or comments?
 Call 1-877-750-3888 Monday-Friday 9AM-5PM EST. The product is not manufactured or distributed by the above group of companies. Distributed by: PL Developments, 200 Hicks Street, Westbury, NY 11590. Product of United Kingdom. Actual Size

See inside for more details or visit www.wellnessbasics.com.

not for sale to those under 18 years of age
 not for sale in vending machines or other any
 source where proof of age cannot be withheld

WELLNESS BASICS NICOTINE GUM COATED FRUIT FLAVOR

<p>Drug Facts (continued)</p> <p>Warnings</p> <p>If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can harm your fetus. Try to stop if you are pregnant or breast-feeding. However, the harm to your child from this medicine is believed to be less than smoking. However, the harm to your child from this medicine is believed to be less than smoking. However, the harm to your child from this medicine is believed to be less than smoking.</p> <p>Use</p> <p>Nicotine patches (used by 2 mg nicotine) Stop smoking and</p> <p>Other Information</p> <p>Keep this and all other medicines out of the reach of children. This medicine contains 2 mg nicotine (tar and nicotine) per piece.</p>	<p>Drug Facts</p> <p>Active Ingredient (in each chewing piece)</p> <p>Nicotine polacrilex (used by 2 mg nicotine)</p> <p>Purpose</p> <p>Stop smoking and</p> <p>Use</p> <p>Nicotine patches (used by 2 mg nicotine) Stop smoking and</p> <p>Warnings</p> <p>If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can harm your fetus. Try to stop if you are pregnant or breast-feeding. However, the harm to your child from this medicine is believed to be less than smoking. However, the harm to your child from this medicine is believed to be less than smoking.</p>
---	--



WELLNESS BASICS Nicotine Gum Coated Fruit Flavor

NICOTINE

nicotine polacrilex gum, chewing

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59726-592
Route of Administration	BUCCAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)	
FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T)	
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
MAGNESIUM OXIDE (UNII: 3A3U0G171G)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
SUCRALOSE (UNII: 96K6UQ3Z D4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XYLITOL (UNII: VCQ006KQ1E)	

Product Characteristics

Color	brown (light brown)	Score	no score
Shape	SQUARE	Size	14mm
Flavor	FRUIT	Imprint Code	

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59726-592-16	160 in 1 CARTON	03/01/2024	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079044	03/01/2024	

Labeler - P & L Development, LLC (800014821)

Revised: 2/2026

P & L Development, LLC