

**PARICALCITOL- paricalcitol capsule, liquid filled**  
**Banner Pharmacaps**

-----

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use paricalcitol capsules safely and effectively. See full prescribing information for paricalcitol capsules.**

**PARICALCITOL Capsules**

**Initial U.S. Approval: 1998**

----- **RECENT MAJOR CHANGES** -----

Warnings and Precautions, Laboratory tests (5.3) 10/2014

----- **INDICATIONS AND USAGE** -----

Paricalcitol capsules is a vitamin D analog indicated for the prevention and treatment of secondary hyperparathyroidism associated with

- Chronic kidney disease (CKD) Stages 3 and 4 (1.1).
- CKD Stage 5 in patients on hemodialysis (HD) or peritoneal dialysis (PD) (1.2).

----- **DOSAGE AND ADMINISTRATION** -----

- CKD Stages 3 and 4: Paricalcitol capsules may be administered once daily or every other day, three times a week (2.1).
- CKD Stage 5: Paricalcitol capsules are dosed every other day, three times a week (2.2). To minimize the risk of hypercalcemia patients should be treated only after their baseline serum calcium has been reduced to 9.5 mg/dL or lower.

Initial Dosage		
CKD Stages 3, 4		CKD Stage 5
Baseline intact parathyroid (iPTH) Level	Starting Dose	
≤ 500 pg/mL	1 mcg daily or 2 mcg three times a week (e.g. every other day)	Dose in micrograms is based on baseline iPTH level (pg/mL)/80. Dose three times a week (e.g. every other day).
> 500 pg/mL	2 mcg daily or 4 mcg three times a week (e.g. every other day)	

Dose Titration			
CKD Stages 3, 4		CKD Stage 5	
iPTH Level Relative to Baseline	Dosing Recommendation		
Decreased by < 30%	Increase dose by 1 mcg daily or 2 mcg three times a week (e.g. every other day)		Dose in micrograms is based on most recent iPTH level (pg/mL)/80 with adjustments based on serum calcium and phosphorous levels. Dose three times a week (e.g. every other day).
Decreased by ≥ 30% and ≤ 60%	Maintain dose		
Decreased by > 60% or iPTH < 60 pg/mL	Decrease dose by 1 mcg daily or 2 mcg three times a week (e.g. every other day)		

----- **DOSAGE FORMS AND STRENGTHS** -----

Capsules: 1 mcg, 2 mcg, and 4 mcg (3).

----- **CONTRAINDICATIONS** -----

- Evidence of hypercalcemia or vitamin D toxicity (4).

----- **WARNINGS AND PRECAUTIONS** -----

- Hypercalcemia: Excessive administration of paricalcitol capsules can cause over suppression of PTH, hypercalcemia, hypercalciuria, hyperphosphatemia, and adynamic bone disease. Prescription-based doses of vitamin D and its derivatives should be withheld during paricalcitol treatment (5.1).
- Digitalis toxicity: Potentiated by hypercalcemia of any cause. Use caution when paricalcitol capsules are prescribed concomitantly with digitalis compounds (5.2).
- Laboratory tests: Monitor serum calcium, serum phosphorus, and serum or plasma iPTH during initial dosing or following any dose adjustment. Paricalcitol capsules may increase serum creatinine and therefore decrease the

estimated GFR (eGFR) (5.3).

- Aluminum overload and toxicity: Avoid excessive use of aluminum containing compounds (5.4).

----- **ADVERSE REACTIONS** -----

The most common adverse reactions (> 5% and more frequent than placebo) include diarrhea, hypertension, dizziness and vomiting. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Banner Pharmacaps at 1-800-526-6993 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch) (6)

----- **DRUG INTERACTIONS** -----

- Strong CYP3A inhibitors (e.g. ketoconazole) will increase the exposure of paricalcitol. Use with caution (7.1).
- Cholestyramine, Mineral Oil: Intestinal absorption of paricalcitol may be reduced if administered simultaneously with mineral oil or cholestyramine (7.2,7.3).

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 12/2014

---

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1 INDICATIONS AND USAGE**

- 1.1 Chronic Kidney Disease Stages 3 and 4
- 1.2 Chronic Kidney Disease Stage 5

**2 DOSAGE AND ADMINISTRATION**

- 2.1 Chronic Kidney Disease Stages 3 and 4
- 2.2 Chronic Kidney Disease Stage 5

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

- 5.1 Hypercalcemia
- 5.2 Digitalis Toxicity
- 5.3 Laboratory Tests
- 5.4 Aluminum Overload and Toxicity

**6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

**7 DRUG INTERACTIONS**

- 7.1 CYP3A Inhibitors
- 7.2 Cholestyramine
- 7.3 Mineral Oil

**8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use

**10 OVERDOSAGE**

**11 DESCRIPTION**

**12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

**13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis and Impairment of Fertility

**14 CLINICAL STUDIES**

- 14.1 Chronic Kidney Disease Stages 3 and 4

## 14.2 Chronic Kidney Disease Stage 5

### 16 HOW SUPPLIED/STORAGE AND HANDLING

### 17 PATIENT COUNSELING INFORMATION

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

#### 1.1 Chronic Kidney Disease Stages 3 and 4

Paricalcitol capsules are indicated for the prevention and treatment of secondary hyperparathyroidism associated with Chronic Kidney Disease (CKD) Stages 3 and 4.

#### 1.2 Chronic Kidney Disease Stage 5

Paricalcitol capsules are indicated for the prevention and treatment of secondary hyperparathyroidism associated with CKD Stage 5 in patients on hemodialysis (HD) or peritoneal dialysis (PD).

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Chronic Kidney Disease Stages 3 and 4

Paricalcitol capsules may be administered daily or three times a week. When dosing three times weekly, the dose should be administered not more frequently than every other day. The total weekly doses for both daily and three times a week dosage regimens are similar [see *Clinical Studies (14.1)*].

Paricalcitol capsules may be taken without regard to food. No dosing adjustment is required in patients with mild and moderate hepatic impairment.

#### Initial Dose

The initial dose of paricalcitol capsules for CKD Stages 3 and 4 patients is based on baseline intact parathyroid hormone (iPTH) levels.

Baseline iPTH Level	Daily Dose	Three Times a Week Dose*
≤ 500 pg/mL	1 mcg	2 mcg
> 500 pg/mL	2 mcg	4 mcg

\* To be administered not more often than every other day

#### Dose Titration

Dosing must be individualized and based on serum or plasma iPTH levels, with monitoring of serum calcium and serum phosphorus. The following is a suggested approach to dose titration.

iPTH Level Relative to Baseline	Paricalcitol Capsule Dose	Dose Adjustment at 2 to 4 Week Intervals	
		Daily Dosage	Three Times a Week Dosage*
The same, increased or decreased by < 30%	Increase dose by	1 mcg	2 mcg

Decreased by $\geq 30\%$ and $\leq 60\%$	Maintain dose	-	-
Decreased by $> 60\%$ or iPTH $< 60$ pg/mL	Decrease dose by	1 mcg	2 mcg
* To be administered not more often than every other day			

If a patient is taking the lowest dose, 1 mcg, on the daily regimen and a dose reduction is needed, the dose can be decreased to 1 mcg three times a week. If a further dose reduction is required, the drug should be withheld as needed and restarted at a lower dosing frequency. If a patient is on a calcium-based phosphate binder, the phosphate-binder dose may be decreased or withheld, or the patient may be switched to a non-calcium-based phosphate binder. If hypercalcemia or an elevated Ca x P is observed, the dose of paricalcitol should be reduced or withheld until these parameters are normalized.

Serum calcium and phosphorus levels should be closely monitored after initiation of paricalcitol capsules, during dose titration periods and during co-administration with strong CYP3A inhibitors [see *Warnings and Precautions (5.3)*, *Drug Interactions (7)* and *Clinical Pharmacology (12.3)*].

## 2.2 Chronic Kidney Disease Stage 5

Paricalcitol capsules are to be administered three times a week, not more frequently than every other day.

Paricalcitol capsules may be taken without regard to food. No dosing adjustment is required in patients with mild and moderate hepatic impairment.

### Initial Dose

The initial dose of paricalcitol capsules in micrograms is based on a baseline iPTH level (pg/mL)/80. To minimize the risk of hypercalcemia patients should be treated only after their baseline serum calcium has been adjusted to 9.5 mg/dL or lower [see *Clinical Pharmacology (12.2)* and *Clinical Studies (14.2)*].

### Dose Titration

Subsequent dosing should be individualized and based on iPTH, serum calcium and phosphorus levels. A suggested dose titration of paricalcitol capsules is based on the following formula:

Titration dose (micrograms) = most recent iPTH level (pg/ml)/80

Serum calcium and phosphorus levels should be closely monitored after initiation, during dose titration periods, and with co-administration of strong P450 3A inhibitors. If an elevated serum calcium or elevated Ca x P is observed and the patient is on a calcium-based phosphate binder, the binder dose may be decreased or withheld, or the patient may be switched to a non-calcium-based phosphate binder. If serum calcium or Ca x P are elevated, the dose should be decreased by 2 to 4 micrograms lower than that calculated by the most recent iPTH/80. If further adjustment is required, the dose of paricalcitol capsules should be reduced or withheld until these parameters are normalized.

As iPTH approaches the target range, small, individualized dose adjustments may be necessary in order to achieve a stable iPTH. In situations where monitoring of iPTH, Ca or P occurs less frequently than once per week, a more modest initial and dose titration ratio (e.g., iPTH/100) may be warranted.

## 3 DOSAGE FORMS AND STRENGTHS

Paricalcitol capsules are available as 1 mcg, 2 mcg, and 4 mcg soft gelatin capsules.

- 1 mcg: oval, gray capsule imprinted with "PA1"
- 2 mcg: oval, orange-brown capsule imprinted with "PA2"
- 4 mcg: oval, yellow capsule imprinted with "PA4"

## 4 CONTRAINDICATIONS

Paricalcitol capsules should not be given to patients with evidence of

- hypercalcemia or
- vitamin D toxicity [see *Warnings and Precautions (5.1)* ].

## **5 WARNINGS AND PRECAUTIONS**

Excessive administration of vitamin D compounds, including paricalcitol capsules, can cause over suppression of PTH, hypercalcemia, hypercalciuria, hyperphosphatemia, and adynamic bone disease.

### **5.1 Hypercalcemia**

Progressive hypercalcemia due to overdosage of vitamin D and its metabolites may be so severe as to require emergency attention [see *Overdosage (10)* ]. Acute hypercalcemia may exacerbate tendencies for cardiac arrhythmias and seizures and may potentiate the action of digitalis. Chronic hypercalcemia can lead to generalized vascular calcification and other soft-tissue calcification. Concomitant administration of high doses of calcium-containing preparations or thiazide diuretics with paricalcitol may increase the risk of hypercalcemia. High intake of calcium and phosphate concomitant with vitamin D compounds may lead to serum abnormalities requiring more frequent patient monitoring and individualized dose titration. Patients also should be informed about the symptoms of elevated calcium, which include feeling tired, difficulty thinking clearly, loss of appetite, nausea, vomiting, constipation, increased thirst, increased urination and weight loss.

Prescription-based doses of vitamin D and its derivatives should be withheld during paricalcitol treatment to avoid hypercalcemia.

### **5.2 Digitalis Toxicity**

Digitalis toxicity is potentiated by hypercalcemia of any cause. Use caution when paricalcitol capsules are prescribed concomitantly with digitalis compounds.

### **5.3 Laboratory Tests**

During the initial dosing or following any dose adjustment of medication, serum calcium, serum phosphorus, and serum or plasma iPTH should be monitored at least every two weeks for 3 months, then monthly for 3 months, and every 3 months thereafter.

In pre-dialysis patients, paricalcitol capsules may increase serum creatinine and therefore decrease the estimated GFR (eGFR). Similar effects have also been seen with calcitriol.

### **5.4 Aluminum Overload and Toxicity**

Aluminum-containing preparations (e.g., antacids, phosphate binders) should not be administered chronically with paricalcitol, as increased blood levels of aluminum and aluminum bone toxicity may occur.

## **6 ADVERSE REACTIONS**

Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical studies of a drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in practice.

### **6.1 Clinical Trials Experience**

#### **CKD Stages 3 and 4**

The safety of paricalcitol capsules has been evaluated in three 24-week (approximately six-month), double-blind, placebo-controlled, multicenter clinical studies involving 220 CKD Stages 3 and 4 patients. Six percent (6%) of paricalcitol capsules treated patients and 4% of placebo treated patients

discontinued from clinical studies due to an adverse event. Adverse events occurring in the paricalcitol capsules group at a frequency of 2% or greater and more frequently than in the placebo group are presented in Table 1:

**Table 1. Treatment-Emergent Adverse Events by Body System Occurring in  $\geq 2\%$  of Subjects in the Paricalcitol-Treated Group of Three, Double-Blind, Placebo-Controlled, Phase 3, CKD Stages 3 and 4 Studies; All Treated Patients**

Adverse Event <sup>a</sup>	Number (%) of Subjects			
	Paricalcitol Capsules (n = 107)		Placebo (n = 113)	
<b>Overall</b>	<b>88</b>	<b>(82%)</b>	<b>86</b>	<b>(76%)</b>
<b>Ear and Labyrinth Disorders</b>				
Vertigo	5	(4.7%)	0	(0.0%)
<b>Gastrointestinal Disorders</b>				
Abdominal Discomfort	4	(3.7%)	1	(0.9%)
Constipation	4	(3.7%)	4	(3.5%)
Diarrhea	7	(6.5%)	5	(4.4%)
Nausea	6	(5.6%)	4	(3.5%)
Vomiting	5	(4.7%)	5	(4.4%)
<b>General Disorders and Administration Site Conditions</b>				
Chest Pain	3	(2.8%)	1	(0.9%)
Edema	6	(5.6%)	5	(4.4%)
Pain	4	(3.7%)	4	(3.5%)
<b>Immune System Disorders</b>				
Hypersensitivity	6	(5.6%)	2	(1.8%)
<b>Infections and Infestations</b>				
Fungal Infection	3	(2.8%)	0	(0.0%)
Gastroenteritis	3	(2.8%)	3	(2.7%)
Infection	3	(2.8%)	3	(2.7%)
Sinusitis	3	(2.8%)	1	(0.9%)
Urinary Tract Infection	3	(2.8%)	1	(0.9%)
Viral Infection	8	(7.5%)	8	(7.1%)
<b>Metabolism and Nutrition Disorders</b>				
Dehydration	3	(2.8%)	1	(0.9%)
<b>Musculoskeletal and Connective Tissue Disorders</b>				
Arthritis	5	(4.7%)	0	(0.0%)
Back Pain	3	(2.8%)	1	(0.9%)
Muscle Spasms	3	(2.8%)	0	(0.0%)
<b>Nervous System Disorders</b>				
Dizziness	5	(4.7%)	5	(4.4%)
Headache	5	(4.7%)	5	(4.4%)
Syncope	3	(2.8%)	1	(0.9%)
<b>Psychiatric Disorders</b>				
Depression	3	(2.8%)	0	(0.0%)
<b>Respiratory, Thoracic and Mediastinal Disorders</b>				

Cough	3	(2.8%)	2	(1.8%)
Oropharyngeal Pain	4	(3.7%)	0	(0.0%)
<b>Skin and Subcutaneous Tissue Disorders</b>				
Pruritus	3	(2.8%)	3	(2.7%)
Rash	4	(3.7%)	1	(0.9%)
Skin Ulcer	3	(2.8%)	0	(0.0%)
<b>Vascular Disorders</b>				
Hypertension	7	(6.5%)	4	(3.5%)
Hypotension	5	(4.7%)	3	(2.7%)
a. Includes only events more common in the paricalcitol treatment group.				

The following adverse reactions, with a causal relationship to paricalcitol, occurred in <2% of the paricalcitol treated patients in the above double-blind, placebo-controlled clinical trial data set.

*Gastrointestinal Disorders:* Dry mouth

*Investigations:* Hepatic enzyme abnormal

*Nervous System Disorders:* Dysgeusia

*Skin and Subcutaneous Tissue Disorders:* Urticaria

### CKD Stage 5

The safety of paricalcitol capsules has been evaluated in one 12-week, double-blind, placebo-controlled, multicenter clinical study involving 88 CKD Stage 5 patients. Sixty-one patients received paricalcitol capsules and 27 patients received placebo.

The proportion of patients who terminated prematurely from the study due to adverse events was 7% for paricalcitol capsules treated patients and 7% for placebo patients.

Adverse events occurring in the paricalcitol capsules group at a frequency of 2% or greater and more frequently than in the placebo group are as follows:

**Table 2. Treatment-Emergent Adverse Events by Body System Occurring in  $\geq 2\%$  of Subjects in the Paricalcitol-Treated Group, Double-Blind, Placebo-Controlled, Phase 3, CKD Stage 5 Study; All Treated Patients**

Adverse Events <sup>a</sup>	Number (%) of Subjects			
	Paricalcitol Capsules (n=61)		Placebo (n = 27)	
<b>Overall</b>	<b>43</b>	<b>(70%)</b>	<b>19</b>	<b>(70%)</b>
<b>Gastrointestinal Disorders</b>				
Constipation	3	(4.9%)	0	(0.0%)
Diarrhea	7	(11.5%)	3	(11.1%)
Vomiting	4	(6.6%)	0	(0.0%)
<b>General Disorders and Administration Site Conditions</b>				
Fatigue	2	(3.3%)	0	(0.0%)
Edema Peripheral	2	(3.3%)	0	(0.0%)
<b>Infections and Infestations</b>				
Nasopharyngitis	5	(8.2%)	2	(7.4%)
Peritonitis	3	(4.9%)	0	(0.0%)

Sinusitis	2	(3.3%)	0	(0.0%)
Urinary Tract Infection	2	(3.3%)	0	(0.0%)
<b>Metabolism and Nutrition Disorders</b>				
Fluid Overload	3	(4.9%)	0	(0.0%)
Hypoglycemia	2	(3.3%)	0	(0.0%)
<b>Nervous System Disorders</b>				
Dizziness	4	(6.6%)	0	(0.0%)
Headache	2	(3.3%)	0	(0.0%)
<b>Psychiatric Disorders</b>				
Anxiety	2	(3.3%)	0	(0.0%)
Insomnia	3	(4.9%)	0	(0.0%)
<b>Renal and Urinary Disorders</b>				
Renal Failure Chronic	2	(3.3%)	0	(0.0%)
a. Includes only events more common in the paricalcitol treatment group.				

The following adverse reactions, with a causal relationship to paricalcitol, occurred in <2% of the paricalcitol treated patients in the above double-blind, placebo-controlled clinical trial data set.

*Gastrointestinal Disorders:* Gastroesophageal reflux disease

*Metabolism and Nutrition Disorders:* Decreased appetite, hypercalcemia, hypocalcemia

*Reproductive System and Breast Disorders:* Breast tenderness

*Skin and Subcutaneous Tissue Disorders:* Acne

## 6.2 Postmarketing Experience

The following additional adverse reactions have been reported during post-approval use and post-approval clinical trials with the active ingredient in paricalcitol capsules:

*Immune System Disorders:* Angioedema (including laryngeal edema)

*Metabolism and Nutrition Disorders:* Hypercalcemia

*Investigations:* Blood creatinine increased

## 7 DRUG INTERACTIONS

### 7.1 CYP3A Inhibitors

Since paricalcitol is partially metabolized by CYP3A, exposure of paricalcitol will be increased while paricalcitol is co-administered with strong CYP3A inhibitors including the following drugs but not limited to: ketoconazole, atazanavir, clarithromycin, indinavir, itraconazole, nefazodone, nelfinavir, ritonavir, saquinavir, telithromycin or voriconazole. Dose adjustment of paricalcitol capsules may be required, and iPTH and serum calcium concentrations should be closely monitored if a patient initiates or discontinues therapy with a strong CYP3A4 inhibitor [see *Clinical Pharmacology (12.3)*].

### 7.2 Cholestyramine

Drugs that impair intestinal absorption of fat-soluble vitamins, such as cholestyramine, may interfere with the absorption of paricalcitol capsules.

### 7.3 Mineral Oil

The use of mineral oil or other substances that may affect absorption of fat may influence the absorption of paricalcitol capsules.



## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Pregnancy Category C.

Paricalcitol has been shown to cause minimal decreases in fetal viability (5%) when administered daily to rabbits at a dose 0.5 times a human dose of 14 mcg or 0.24 mcg/kg (based on body surface area, mcg/m<sup>2</sup>), and when administered to rats at a dose two times the 0.24 mcg/kg human dose (based on body surface area, mcg/m<sup>2</sup>). At the highest dose tested, 20 mcg/kg administered three times per week in rats (13 times the 14 mcg human dose based on surface area, mcg/m<sup>2</sup>), there was a significant increase in the mortality of newborn rats at doses that were maternally toxic and are known to produce hypercalcemia in rats. No other effects on offspring development were observed.

Paricalcitol was not teratogenic at the doses tested.

Paricalcitol (20 mcg/kg) has been shown to cross the placental barrier in rats. There are no adequate and well-controlled clinical studies in pregnant women. Paricalcitol capsules should be used during pregnancy only if the potential benefit to the mother justifies the potential risk to the fetus.

### 8.3 Nursing Mothers

Studies in rats have shown that paricalcitol is present in the milk. It is not known whether paricalcitol is excreted in human milk. In the nursing patient, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

### 8.4 Pediatric Use

Safety and efficacy of paricalcitol capsules in pediatric patients have not been established.

### 8.5 Geriatric Use

Of the total number (n = 220) of CKD Stages 3 and 4 patients in clinical studies of paricalcitol capsules, 49% were age 65 and over, while 17% were age 75 and over. Of the total number (n = 88) of CKD Stage 5 patients in the pivotal study of paricalcitol capsules, 28% were age 65 and over, while 6% were age 75 and over. No overall differences in safety and effectiveness were observed between these patients and younger patients, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

## 10 OVERDOSAGE

Excessive administration of paricalcitol capsules can cause hypercalcemia, hypercalciuria, and hyperphosphatemia, and over suppression of PTH [see *Warnings and Precautions (5.1)*].

### Treatment of Overdosage

The treatment of acute overdosage of paricalcitol capsules should consist of general supportive measures. If drug ingestion is discovered within a relatively short time, induction of emesis or gastric lavage may be of benefit in preventing further absorption. If the drug has passed through the stomach, the administration of mineral oil may promote its fecal elimination. Serial serum electrolyte determinations (especially calcium), rate of urinary calcium excretion, and assessment of electrocardiographic abnormalities due to hypercalcemia should be obtained. Such monitoring is critical in patients receiving digitalis. Discontinuation of supplemental calcium and institution of a low-calcium diet are also indicated in accidental overdosage. Due to the relatively short duration of the pharmacological action of paricalcitol, further measures are probably unnecessary. If persistent and markedly elevated serum calcium levels occur, there are a variety of therapeutic alternatives that may be

considered depending on the patient's underlying condition. These include the use of drugs such as phosphates and corticosteroids, as well as measures to induce an appropriate forced diuresis.

Paricalcitol is not significantly removed by dialysis.

## 11 DESCRIPTION

Paricalcitol, USP, the active ingredient in paricalcitol capsules, is a synthetically manufactured, metabolically active vitamin D analog of calcitriol with modifications to the side chain (D<sub>2</sub>) and the A (19-nor) ring. Paricalcitol is indicated for the prevention and treatment of secondary hyperparathyroidism in chronic kidney disease. Paricalcitol is available as soft gelatin capsules for oral administration containing 1 microgram, 2 micrograms or 4 micrograms of paricalcitol. Each capsule also contains medium chain triglycerides, alcohol, and butylated hydroxytoluene. The medium chain triglycerides are fractionated from coconut oil or palm kernel oil. The capsule shell is composed of gelatin, glycerin, titanium dioxide, iron oxide red (2 microgram capsules only), iron oxide yellow (2 microgram and 4 microgram capsules), iron oxide black (1 microgram capsules only), and water. The capsules are printed with black ink composed of purified water, black iron oxide, isopropyl alcohol, propylene glycol and hypromellose.

Paricalcitol is a white, crystalline powder with the empirical formula of C<sub>27</sub>H<sub>44</sub>O<sub>3</sub>, which corresponds to a molecular weight of 416.64. Paricalcitol is chemically designated as 19-nor-1 $\alpha$ ,3 $\beta$ ,25-trihydroxy-9,10-secoergosta-5(Z),7(E),22(E)-triene and has the following structural formula:

## 12 CLINICAL PHARMACOLOGY

Secondary hyperparathyroidism is characterized by an elevation in parathyroid hormone (PTH) associated with inadequate levels of active vitamin D hormone. The source of vitamin D in the body is from synthesis in the skin as vitamin D<sub>3</sub> and from dietary intake as either vitamin D<sub>2</sub> or D<sub>3</sub>. Both vitamin D<sub>2</sub> and D<sub>3</sub> require two sequential hydroxylations in the liver and the kidney to bind to and to activate the vitamin D receptor (VDR). The endogenous VDR activator, calcitriol [1,25(OH)<sub>2</sub>D<sub>3</sub>], is a hormone that binds to VDRs that are present in the parathyroid gland, intestine, kidney, and bone to maintain parathyroid function and calcium and phosphorus homeostasis, and to VDRs found in many other tissues, including prostate, endothelium and immune cells. VDR activation is essential for the proper formation and maintenance of normal bone. In the diseased kidney, the activation of vitamin D is diminished,

resulting in a rise of PTH, subsequently leading to secondary hyperparathyroidism and disturbances in the calcium and phosphorus homeostasis. Decreased levels of  $1,25(\text{OH})_2\text{D}_3$  have been observed in early stages of chronic kidney disease. The decreased levels of  $1,25(\text{OH})_2\text{D}_3$  and resultant elevated PTH levels, both of which often precede abnormalities in serum calcium and phosphorus, affect bone turnover rate and may result in renal osteodystrophy.

### 12.1 Mechanism of Action

Paricalcitol is a synthetic, biologically active vitamin D<sub>2</sub> analog of calcitriol. Preclinical and *in vitro* studies have demonstrated that paricalcitol's biological actions are mediated through binding of the VDR, which results in the selective activation of vitamin D responsive pathways. Vitamin D and paricalcitol have been shown to reduce parathyroid hormone levels by inhibiting PTH synthesis and secretion.

### 12.2 Pharmacodynamics

Paricalcitol decreases serum intact parathyroid hormone (iPTH) and increases serum calcium and serum phosphorous in both HD and PD patients. This observed relationship was quantified using a mathematical model for HD and PD patient populations separately. Computer-based simulations of 100 trials in HD or PD patients (N = 100) using these relationships predict slightly lower efficacy (at least two consecutive  $\geq 30\%$  reductions from baseline iPTH) with lower hypercalcemia rates (at least two consecutive serum calcium  $\geq 10.5$  mg/dL) for lower iPTH-based dosing regimens. Further lowering of hypercalcemia rates was predicted if the treatment with paricalcitol is initiated in patients with lower serum calcium levels at screening.

Based on these simulations, a dosing regimen of iPTH/80 with a screening serum calcium  $\leq 9.5$  mg/dL, approximately 76.5% (95% CI: 75.6% – 77.3%) of HD patients are predicted to achieve at least two consecutive weekly  $\geq 30\%$  reductions from baseline iPTH over a duration of 12 weeks. The predicted incidence of hypercalcemia is 0.8% (95% CI: 0.7% – 1.0%). In PD patients, with this dosing regimen, approximately 83.3% (95% CI: 82.6% – 84.0%) of patients are predicted to achieve at least two consecutive weekly  $\geq 30\%$  reductions from baseline iPTH. The predicted incidence of hypercalcemia is 12.4% (95% CI: 11.7% - 13.0%) [see *Clinical Studies (14.2) and Dosage and Administration (2.2)*].

### 12.3 Pharmacokinetics

#### Absorption

The mean absolute bioavailability of paricalcitol capsules under low-fat fed condition ranged from 72% to 86% in healthy subjects, CKD Stage 5 patients on HD, and CKD Stage 5 patients on PD. A food effect study in healthy subjects indicated that the  $C_{\text{max}}$  and  $\text{AUC}_{0-\infty}$  were unchanged when paricalcitol was administered with a high fat meal compared to fasting. Food delayed  $T_{\text{max}}$  by about 2 hours. The  $\text{AUC}_{0-\infty}$  of paricalcitol increased proportionally over the dose range of 0.06 to 0.48 mcg/kg in healthy subjects.

#### Distribution

Paricalcitol is extensively bound to plasma proteins ( $\geq 99.8\%$ ). The mean apparent volume of distribution following a 0.24 mcg/kg dose of paricalcitol in healthy subjects was 34 L. The mean apparent volume of distribution following a 4 mcg dose of paricalcitol in CKD Stage 3 and a 3 mcg dose in CKD Stage 4 patients is between 44 and 46 L.

#### Metabolism

After oral administration of a 0.48 mcg/kg dose of <sup>3</sup>H-paricalcitol, parent drug was extensively metabolized, with only about 2% of the dose eliminated unchanged in the feces, and no parent drug was found in the urine. Several metabolites were detected in both the urine and feces. Most of the systemic exposure was from the parent drug. Two minor metabolites, relative to paricalcitol, were detected in human plasma. One metabolite was identified as 24(R)-hydroxy paricalcitol, while the other metabolite

was unidentified. The 24(R)-hydroxy paricalcitol is less active than paricalcitol in an *in vivo* rat model of PTH suppression.

*In vitro* data suggest that paricalcitol is metabolized by multiple hepatic and non-hepatic enzymes, including mitochondrial CYP24, as well as CYP3A4 and UGT1A4. The identified metabolites include the product of 24(R)-hydroxylation, 24,26- and 24,28-dihydroxylation and direct glucuronidation.

#### Elimination

Paricalcitol is eliminated primarily via hepatobiliary excretion; approximately 70% of the radiolabeled dose is recovered in the feces and 18% is recovered in the urine. While the mean elimination half-life of paricalcitol is 4 to 6 hours in healthy subjects, the mean elimination half-life of paricalcitol in CKD Stages 3, 4, and 5 (on HD and PD) patients ranged from 14 to 20 hours.

**Table 3. Paricalcitol Capsule Pharmacokinetic Characteristics in CKD Stages 3, 4, and 5 Patients**

Pharmacokinetic Parameters	CKD Stage 3 n = 15*	CKD Stage 4 n = 14*	CKD Stage 5 HD** n = 14	CKD Stage 5 PD** n = 8
C <sub>max</sub> (ng/mL)	0.11 ± 0.04	0.06 ± 0.01	0.575 ± 0.17	0.413 ± 0.06
AUC <sub>0-∞</sub> (ng•h/mL)	2.42 ± 0.61	2.13 ± 0.73	11.67 ± 3.23	13.41 ± 5.48
CL/F (L/h)	1.77 ± 0.50	1.52 ± 0.36	1.82 ± 0.75	1.76 ± 0.77
V/F (L)	43.7 ± 14.4	46.4 ± 12.4	38 ± 16.4	48.7 ± 15.6
t <sub>1/2</sub>	16.8 ± 2.65	19.7 ± 7.2	13.9 ± 5.1	17.7 ± 9.6

\* Four mcg paricalcitol capsules were given to CKD Stage 3 patients; three mcg paricalcitol capsules were given to CKD Stage 4 patients.  
 \*\* CKD Stage 5 HD and PD patients received a 0.24 mcg/kg dose of paricalcitol as capsules.

#### Specific Populations

##### *Geriatric*

The pharmacokinetics of paricalcitol has not been investigated in geriatric patients greater than 65 years [see *Use in Specific Populations (8.5)*].

##### *Pediatric*

The pharmacokinetics of paricalcitol has not been investigated in patients less than 18 years of age.

##### *Gender*

The pharmacokinetics of paricalcitol following single doses over the 0.06 to 0.48 mcg/kg dose range was gender independent.

##### *Hepatic Impairment*

The disposition of paricalcitol (0.24 mcg/kg) was compared in patients with mild (n = 5) and moderate (n = 5) hepatic impairment (as indicated by the Child-Pugh method) and subjects with normal hepatic function (n = 10). The pharmacokinetics of unbound paricalcitol was similar across the range of hepatic function evaluated in this study. No dose adjustment is required in patients with mild and moderate hepatic impairment. The influence of severe hepatic impairment on the pharmacokinetics of paricalcitol has not been evaluated.

##### *Renal Impairment*

Following administration of paricalcitol capsules, the pharmacokinetic profile of paricalcitol for CKD Stage 5 on HD or PD was comparable to that in CKD 3 or 4 patients. Therefore, no special dose adjustments are required other than those recommended in the Dosage and Administration section [see

*Dosage and Administration (2)*].

### *Drug Interactions*

An *in vitro* study indicates that paricalcitol is neither an inhibitor of CYP1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP2E1 or CYP3A nor an inducer of CYP2B6, CYP2C9 or CYP3A. Hence, paricalcitol is neither expected to inhibit nor induce the clearance of drugs metabolized by these enzymes.

### Omeprazole

The effect of omeprazole (40 mg capsule), a strong inhibitor of CYP2C19, on paricalcitol (four 4 mcg capsules) pharmacokinetics was investigated in a single dose, crossover study in healthy subjects. The pharmacokinetics of paricalcitol was not affected when omeprazole was administered approximately 2 hours prior to the paricalcitol dose.

### Ketoconazole

The effect of multiple doses of ketoconazole, a strong inhibitor of CYP3A, administered as 200 mg BID for 5 days on the pharmacokinetics of paricalcitol (4 mcg capsule) has been studied in healthy subjects. The  $C_{max}$  of paricalcitol was minimally affected, but  $AUC_{0-\infty}$  approximately doubled in the presence of ketoconazole. The mean half-life of paricalcitol was 17.0 hours in the presence of ketoconazole as compared to 9.8 hours, when paricalcitol was administered alone [see *Drug Interactions (7)*].

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis and Impairment of Fertility**

In a 104-week carcinogenicity study in CD-1 mice, an increased incidence of uterine leiomyoma and leiomyosarcoma was observed at subcutaneous doses of 1, 3, 10 mcg/kg given three times weekly (2 to 15 times the AUC at a human dose of 14 mcg, equivalent to 0.24 mcg/kg based on AUC). The incidence rate of uterine leiomyoma was significantly different than the control group at the highest dose of 10 mcg/kg. In a 104-week carcinogenicity study in rats, there was an increased incidence of benign adrenal pheochromocytoma at subcutaneous doses of 0.15, 0.5, 1.5 mcg/kg (< 1 to 7 times the exposure following a human dose of 14 mcg, equivalent to 0.24 mcg/kg based on AUC). The increased incidence of pheochromocytomas in rats may be related to the alteration of calcium homeostasis by paricalcitol. Paricalcitol did not exhibit genetic toxicity *in vitro* with or without metabolic activation in the microbial mutagenesis assay (Ames Assay), mouse lymphoma mutagenesis assay (L5178Y), or a human lymphocyte cell chromosomal aberration assay. There was also no evidence of genetic toxicity in an *in vivo* mouse micronucleus assay. Paricalcitol had no effect on fertility (male or female) in rats at intravenous doses up to 20 mcg/kg/dose (equivalent to 13 times a human dose of 14 mcg based on surface area, mcg/m<sup>2</sup>).

## **14 CLINICAL STUDIES**

### **14.1 Chronic Kidney Disease Stages 3 and 4**

The safety and efficacy of paricalcitol capsules were evaluated in three, 24-week, double blind, placebo-controlled, randomized, multicenter, Phase 3 clinical studies in CKD Stages 3 and 4 patients. Two studies used an identical three times a week dosing design, and one study used a daily dosing design. A total of 107 patients received paricalcitol capsules and 113 patients received placebo. The mean age of the patients was 63 years, 68% were male, 71% were Caucasian, and 26% were African-American. The average baseline iPTH was 274 pg/mL (range: 145-856 pg/mL). The average duration of CKD prior to study entry was 5.7 years. At study entry 22% were receiving calcium based phosphate binders and/or calcium supplements. Baseline 25-hydroxyvitamin D levels were not measured.


The initial dose of paricalcitol capsules was based on baseline iPTH. If iPTH was  $\leq 500$  pg/mL, paricalcitol capsules were administered 1 mcg daily or 2 mcg three times a week, not more than every other day. If iPTH was  $> 500$  pg/mL, paricalcitol capsules were administered 2 mcg daily or 4 mcg three times a week, not more than every other day. The dose was increased by 1 mcg daily or 2 mcg three times a week every 2 to 4 weeks until iPTH levels were reduced by at least 30% from baseline. The overall average weekly dose of paricalcitol capsules was 9.6 mcg/week in the daily regimen and 9.5 mcg/week in the three times a week regimen.

In the clinical studies, doses were titrated for any of the following reasons: if iPTH fell to  $< 60$  pg/mL, or decreased  $> 60\%$  from baseline, the dose was reduced or temporarily withheld; if iPTH decreased  $< 30\%$  from baseline and serum calcium was  $\leq 10.3$  mg/dL and serum phosphorus was  $\leq 5.5$  mg/dL, the dose was increased; and if iPTH decreased between 30 to 60% from baseline and serum calcium and phosphorus were  $\leq 10.3$  mg/dL and  $\leq 5.5$  mg/dL, respectively, the dose was maintained. Additionally, if serum calcium was between 10.4 to 11.0 mg/dL, the dose was reduced irrespective of iPTH, and the dose was withheld if serum calcium was  $> 11.0$  mg/dL. If serum phosphorus was  $> 5.5$  mg/dL, dietary counseling was provided, and phosphate binders could have been initiated or increased. If the elevation persisted, the paricalcitol capsules dose was decreased. Seventy-seven percent (77%) of the paricalcitol capsules treated patients and 82% of the placebo treated patients completed the 24-week treatment. The primary efficacy endpoint of at least two consecutive  $\geq 30\%$  reductions from baseline iPTH was achieved by 91% of paricalcitol capsules treated patients and 13% of the placebo treated patients ( $p < 0.001$ ). The proportion of paricalcitol capsules treated patients achieving two consecutive  $\geq 30\%$  reductions was similar between the daily and the three times a week regimens (daily: 30/33, 91%; three times a week: 62/68, 91%).

The incidence of hypercalcemia (defined as two consecutive serum calcium values  $> 10.5$  mg/dL), hyperphosphatemia and elevated Ca x P product in paricalcitol capsules treated patients was similar to placebo. There were no treatment related adverse events associated with hypercalcemia or hyperphosphatemia in the paricalcitol capsules group. No increases in urinary calcium or phosphorus were detected in paricalcitol capsules treated patients compared to placebo.

The pattern of change in the mean values for serum iPTH during the studies is shown in Figure 1.

**Figure 1. Mean Values for Serum iPTH Over Time in the Three Double-Blind, Placebo-Controlled, Phase 3, CKD Stages 3 and 4 Studies Combined**



The mean changes from baseline to final treatment visit in serum iPTH, calcium, phosphorus, calcium-phosphorus product (Ca x P), and bone-specific alkaline phosphatase are shown in Table 4.

**Table 4. Mean Changes from Baseline to Final Treatment Visit in Serum iPTH, Bone Specific Alkaline Phosphatase, Calcium, Phosphorus, and Calcium x Phosphorus Product in Three Combined Double-Blind, Placebo-Controlled, Phase 3, CKD Stages 3 and 4 Studies**

	<b>Paricalcitol Capsules</b>	<b>Placebo</b>
<b>iPTH (pg/mL)</b>	n = 104	n = 110
Mean Baseline Value	266	279
Mean Final Treatment Value	162	315
Mean Change from Baseline (SE)	-104 (9.2)	+35 (9.0)
<b>Bone Specific Alkaline Phosphatase (mcg/L)</b>	n = 101	n = 107
Mean Baseline	17.1	18.8
Mean Final Treatment Value	9.2	17.4
Mean Change from Baseline (SE)	-7.9 (0.76)	-1.4 (0.74)
<b>Calcium (mg/dL)</b>	n = 104	n = 110
Mean Baseline	9.3	9.4
Mean Final Treatment Value	9.5	9.3
Mean Change from Baseline (SE)	+0.2 (0.04)	-0.1 (0.04)
<b>Phosphorus (mg/dL)</b>	n = 104	n = 110
Mean Baseline	4.0	4.0
Mean Final Treatment Value	4.3	4.3
Mean Change from Baseline (SE)	+0.3 (0.08)	+0.3 (0.08)
<b>Calcium x Phosphorus Product (mg<sup>2</sup>/dL<sup>2</sup>)</b>	n = 104	n = 110
Mean Baseline	36.7	36.9
Mean Final Treatment Value	40.7	39.7
Mean Change from Baseline (SE)	+4.0 (0.74)	+2.9 (0.72)

#### 14.2 Chronic Kidney Disease Stage 5

The safety and efficacy of paricalcitol capsules were evaluated in a Phase 3, 12-week, double blind, placebo-controlled, randomized, multicenter study in patients with CKD Stage 5 on HD or PD. The study used a three times a week dosing design. A total of 61 patients received paricalcitol capsules and 27 patients received placebo. The mean age of the patients was 57 years, 67% were male, 50% were Caucasian, 45% were African- American, and 53% were diabetic. The average baseline iPTH was 701 pg/mL (range: 216-1933 pg/mL). The average time since first dialysis across all subjects was 3.3 years.


The initial dose of paricalcitol capsules was based on baseline iPTH/60. Subsequent dose adjustments were based on iPTH/60 as well as primary chemistry results that were measured once a week. Starting at Treatment Week 2, study drug was maintained, increased or decreased weekly based on the results of the previous week's calculation of iPTH/60. Paricalcitol capsules were administered three times a week, not more than every other day.

The proportion of patients achieving at least two consecutive weekly  $\geq 30\%$  reductions from baseline iPTH was 88% of paricalcitol capsules treated patients and 13% of the placebo treated patients. The proportion of patients achieving at least two consecutive weekly  $\geq 30\%$  reductions from baseline iPTH was similar for HD and PD patients.

The incidence of hypercalcemia (defined as two consecutive serum calcium values  $> 10.5$  mg/dL) in patients treated with paricalcitol capsules was 6.6% as compared to 0% for patients given placebo. In PD patients the incidence of hypercalcemia in patients treated with paricalcitol capsules was 21% as compared to 0% for patients given placebo. The patterns of change in the mean values for serum iPTH

are shown in Figure 2. The rate of hypercalcemia with paricalcitol capsules may be reduced with a lower dosing regimen based on the iPTH/80 formula as shown by computer simulations. The hypercalcemia rate can be further predicted to decrease, if the treatment is initiated in only those with baseline serum calcium  $\leq 9.5$  mg/dL [see *Clinical Pharmacology (12.2) and Dosage and Administration (2.2)*].

**Figure 2. Mean Values for Serum iPTH Over Time in a Phase 3, Double-Blind, Placebo-Controlled CKD Stage 5 Study**



## 16 HOW SUPPLIED/STORAGE AND HANDLING

Paricalcitol capsules are available as 1 mcg, 2 mcg, and 4 mcg capsules.

The 1 mcg capsule is an oval, gray, soft gelatin capsule imprinted with "PA1", and is available in the following package size:

Bottles of 30 (NDC 10888-5001-2)

The 2 mcg capsule is an oval, orange-brown, soft gelatin capsule imprinted with "PA2", and is available in the following package size:

Bottles of 30 (NDC 10888-5002-2)

The 4 mcg capsule is an oval, yellow soft gelatin capsule imprinted with "PA4", and is available in the following package size:

Bottles of 30 (NDC 10888-5003-2)

### Storage

Store paricalcitol capsules at 25°C (77°F). Excursions permitted between 15°- 30°C (59°- 86°F). See USP Controlled Room Temperature.

## 17 PATIENT COUNSELING INFORMATION

Patients should be advised:

- of the most common adverse reactions with use of paricalcitol capsules, which include diarrhea,



hypertension, dizziness and vomiting.

- to adhere to instructions regarding diet and phosphorus restriction.
- to contact a health care provider if you develop symptoms of elevated calcium, (e.g. feeling tired, difficulty thinking clearly, loss of appetite, nausea, vomiting, constipation, increased thirst, increased urination and weight loss).
- to return to the physician's office for routine monitoring. More frequent monitoring is necessary during the initiation of therapy, following dose changes or when potentially interacting medications are started or discontinued.
- to inform their physician of all medications, including prescription and nonprescription drugs, supplements, and herbal preparations they are taking and any change to their medical condition. Patients should also be advised to inform their physicians prescribing a new medication that they are taking paricalcitol capsules.

Banner Pharmacaps Inc.

4125 Premier Drive

High Point, NC 27265

10/2014

### PRINCIPAL DISPLAY PANEL - 1 mcg Capsule Bottle Label

**Banner Pharmacaps Inc.**

NDC 10888-5001-2

**Paricalcitol  
Capsules**

**1 mcg**

30 Capsules

Rx only

The image shows a rectangular label with rounded corners and a red border. On the left side, there is vertical text: "Store at 25°C (77°F). Excursions permitted between 15°-30°C (59°-86°F). See USP Controlled Room Temperature." In the center, there is a large green box containing "1 mcg" in white. Above this box, the text reads "Each capsule contains: 1 mcg Paricalcitol, USP, and 1.2% v/v (1.0% w/w) alcohol, USP dehydrated". Below the green box, it says "30 Capsules" and "Rx only". To the right of the green box, the text reads "Banner Pharmacaps Inc. NDC 10888-5001-2 Paricalcitol Capsules". Further right, it says "Manufactured by: Banner Pharmacaps Inc. High Point, NC 27265" and "Rev. 06/12". At the bottom right, there are fields for "Lot. No.:" and "Exp. Date:". On the far right, there is a barcode with the number "N 3 10888-5001-2 8" below it.

### PRINCIPAL DISPLAY PANEL - 2 mcg Capsule Bottle Label

**Banner Pharmacaps Inc.**

NDC 10888-5002-2

**Paricalcitol  
Capsules**

**2 mcg**

30 Capsules

Rx only

**Banner Pharmacaps Inc.**  
NDC 10888-5002-2  
**Paricalcitol Capsules**  
**2 mcg**  
30 Capsules Rx only

Each capsule contains:  
2 mcg Paricalcitol, USP, and  
1.2% v/v (1.0% w/w) alcohol,  
USP dehydrated

See package insert for full  
prescribing information.  
Dispense in a USP tight  
container. Do not accept if seal  
over bottle opening is broken  
or missing.

Store at 25°C (77°F).  
Excursions permitted between  
15°-30°C (59°-86°F). See USP  
Controlled Room Temperature.

Manufactured by:  
Banner Pharmacaps Inc.  
High Point, NC 27265  
Rev. 06/12

Lot. No.:  
Exp. Date:



**PRINCIPAL DISPLAY PANEL - 4 mcg Capsule Bottle Label**

**Banner Pharmacaps Inc.**

NDC 10888-5003-2

**Paricalcitol Capsules**

**4 mcg**

30 Capsules

Rx only

**Banner Pharmacaps Inc.**  
NDC 10888-5003-2  
**Paricalcitol Capsules**  
**4 mcg**  
30 Capsules Rx only

Each capsule contains:  
4 mcg Paricalcitol, USP, and  
1.2% v/v (1.0% w/w) alcohol,  
USP dehydrated

See package insert for full  
prescribing information.  
Dispense in a USP tight  
container. Do not accept if seal  
over bottle opening is broken  
or missing.

Store at 25°C (77°F).  
Excursions permitted between  
15°-30°C (59°-86°F). See USP  
Controlled Room Temperature.

Manufactured by:  
Banner Pharmacaps Inc.  
High Point, NC 27265  
Rev. 06/12

Lot. No.:  
Exp. Date:



**PARICALCITOL**

paricalcitol capsule, liquid filled

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:10888-5001
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>Paricalcitol</b> (UNII: 6702D36OG5) (Paricalcitol - UNII:6702D36OG5)	Paricalcitol	1 ug

**Inactive Ingredients**

Ingredient Name	Strength
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	

WATER (UNII: 059QF0KO0R)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALCOHOL (UNII: 3K9958V90M)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	

### Product Characteristics

Color	GRAY	Score	no score
Shape	OVAL	Size	10mm
Flavor		Imprint Code	PA1
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10888-5001-2	30 in 1 BOTTLE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202539	03/27/2014	

## PARICALCITOL

paricalcitol capsule, liquid filled

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:10888-5002
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PARICALCITOL (UNII: 6702D36OG5) (PARICALCITOL - UNII:6702D36OG5)	PARICALCITOL	2 ug

### Inactive Ingredients

Ingredient Name	Strength
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALCOHOL (UNII: 3K9958V90M)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	

### Product Characteristics

Color	ORANGE (brown)	Score	no score
Shape	OVAL	Size	10mm
Flavor		Imprint Code	PA2
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10888-5002-2	30 in 1 BOTTLE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202539	03/27/2014	

## PARICALCITOL

paricalcitol capsule, liquid filled

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:10888-5003
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PARICALCITOL (UNII: 6702D36OG5) (PARICALCITOL - UNII:6702D36OG5)	PARICALCITOL	4 ug

### Inactive Ingredients

Ingredient Name	Strength
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALCOHOL (UNII: 3K9958V90M)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	

### Product Characteristics

<b>Color</b>	YELLOW	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	PA4
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10888-5003-2	30 in 1 BOTTLE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202539	03/27/2014	

**Labeler** - Banner Pharmacaps (002193829)

### Establishment

Name	Address	ID/FEI	Business Operations
Formosa Laboratories, Inc.		657846858	API MANUFACTURE(10888-5001, 10888-5002, 10888-5003)

Revised: 12/2014

Banner Pharmacaps