

**FEXOFENADINE HCL AND PSEUDOEPHEDRINE HCL- fexofenadine hcl and pseudoephedrine hcl tablet, extended release  
Walgreens company**

-----  
**Fexofenadine HCl 180 mg and Pseudoephedrine HCl 240 mg ER Tablets, USP**

**Active ingredient(s)**

Fexofenadine HCl USP, 180 mg

Pseudoephedrine HCl USP, 240 mg

**Purpose**

Antihistamine

Nasal decongestant

**Use(s)**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

**Warnings**

**Do not use**

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have difficulty swallowing

**Ask a doctor before use if you have**

- heart disease
- thyroid disease
- glaucoma
- high blood pressure

- diabetes
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.

**When using this product**

- **do not take more than directed**
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)
- the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).

**Stop use and ask doctor if**

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- you get nervous, dizzy, or sleepless

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not divide, crush, chew or dissolve the tablet; swallow tablet whole

adults and children 12 years of age and over	take 1 tablet with a glass of water every 24 hours on an empty stomach; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

**Other information**

- each tablet contains: **28 mg sodium**
- safety sealed: do not use if carton is opened or if individual blister units are torn or opened
- store between 20° - 25°C (68° - 77°F)
- FDA approved dissolution test specifications differ from USP

**Inactive ingredients**

acetone, black iron oxide, cellulose acetate, colloidal silicon dioxide, copovidone, croscarmellose sodium, FD&C blue #1 aluminum lake, hypromellose, isopropyl alcohol,

magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, propylene glycol, red iron oxide, sodium chloride, talc, titanium dioxide, water

**Questions?**

**Questions?** Call 1-888-375-3784 Weekdays (9am - 8pm EST)

Distributed by:

Dr. Reddy's Laboratories, Inc.

Princeton, NJ 08540

**Carton Label**



**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>FEXOFENADINE HYDROCHLORIDE</b> (UNII: 2S068B75ZU) (FEXOFENADINE - UNII:E6582LOH6V)	FEXOFENADINE HYDROCHLORIDE	180 mg
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b> (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	240 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>ACETONE</b> (UNII: 1364PS73AF)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>CELLULOSE ACETATE</b> (UNII: 3J2P07GVB6)	
<b>COPOVIDONE</b> (UNII: D9C330MD8B)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>Polyethylene Glycol, Unspecified</b> (UNII: 3WJQ0SDW1A)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302)	
<b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)	

**Product Characteristics**

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	892
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0049-10	2 in 1 CARTON	07/13/2022	08/30/2026
1		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0363-0049-15	3 in 1 CARTON	07/13/2022	07/30/2026
2		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079043	07/13/2022	08/30/2026

**Labeler** - Walgreens company (008965063)

## Establishment

Name	Address	ID/FEI	Business Operations
Dr.Reddy's Laboratories Limited (FTO III)		918608162	analysis(0363-0049) , manufacture(0363-0049)

Revised: 7/2025

Walgreens company