

**LANSOPRAZOLE- lansoprazole capsule, delayed release**  
**AMERISOURCEBERGEN DRUG CORPORATION**

-----  
**Lansoprazole Delayed Release Capsules USP**

**Active ingredient (in each capsule)**

Lansoprazole USP, 15 mg

**Purpose**

Acid reducer

**Use**

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**

**Allergy alert:**

- do not use if you are allergic to lansoprazole
- lansoprazole may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have**

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor

**Ask a doctor before use if you have**

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

**If pregnant or breast-feeding,**

ask a healthcare professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (**1-800-222-1222**).

**Directions**

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

**Repeated 14-Day Courses (if needed)**

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

**Other information**

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20 - 25°C (68° - 77° F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

## **Inactive ingredients**

D&C Red 28, FD&C Blue No.1, FD&C Green 3, FD&C Red 40, gelatin, hydroxypropyl cellulose, iron oxide black, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer, polyethylene glycol 6000, polysorbate 80, sodium lauryl sulphate, starch (corn), sucrose, sugar spheres, talc, titanium dioxide

## **Questions or comments?**

call **1-888-375-3784**

## **Tips For Managing Heartburn**

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

## **Distributed By:**

AmerisourceBergen

1 West First Avenue

Conshohocken PA 19428

Made in India

## **Principal Display Panel**



NDC 46122-744-74

# Lansoprazole <sup>24 HR</sup>

delayed-release capsules, USP  
15 mg

## Acid Reducer

**Treats Frequent Heartburn**  
• May take 1 to 4 days for full effect

14 Capsules  
One 14-day Course of Treatment

**IMPORTANT:** This label does not contain full product information. See carton for complete information. Read the directions and warnings before use. Retain carton for reference.

**TAMPER EVIDENT:** Do not use if foil seal under cap printed with "sealed for your protection" or black band around center of capsule is broken or missing.

### Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

Distributed By: AmerisourceBergen  
1 West First Avenue  
Conshohocken PA 19428  
Made in India REV 01/23

150091186

LOT/EXP

PEEL HERE →

### Directions (continued)

#### 14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

#### Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

CONTINUED →

### Directions (continued)

- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

### Other information

- read the directions and warnings before use
- keep the carton. It contains important information
- store at 20° - 25°C (68° - 77° F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use
- **If pregnant or breast-feeding,** ask a health care professional before use.
- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Questions or comments?

call 1-888-375-3784

14ct Carton Label:





**Tips for Managing Heartburn**

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

**Lansoprazole**  
delayed-release capsules USP 15 mg

**Acid Reducer**

1 BOTTLE INSIDE

---



**Compare to Prevacid® 24 HR active ingredient\***

NDC 46122-744-74

**Lansoprazole** °24 HR  
delayed-release capsules USP, 15 mg

**Acid Reducer**

**Treats Frequent Heartburn**

• May take 1 to 4 days for full effect

1 BOTTLE INSIDE  
14 CAPSULES TOTAL

**TAMPER EVIDENT:** Do not use if foil seal under cap printed with "swallow" for your protection" or if foil seal around center of capsule is broken or missing.

**IMPORTANT:** Read the directions and warnings before use. Keep the cap on. It contains important information.

**14 Capsules** One 14-day Course of Treatment



actual size

\*This product is not manufactured or distributed by Perrigo, distributor of Prevacid® 24HR Capsules. Prevacid® is a registered trademark of Helsiris Pharmaceuticals U.S.A., Inc.

Distributed By:  
AmerisourceBergen  
1 West First Avenue  
Conshohocken PA 19428  
Questions or Concerns?  
[www.mygnp.com](http://www.mygnp.com)




REV 01/23

LOT

EXP

Made in India

**Drug Facts**

**Active ingredient (in each capsule)**  
Lansoprazole USP, 15 mg

**Purpose**  
Acid Reducer

**Use**

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**

**Allergy alert:**

- lansoprazole may cause severe skin reactions.

Symptoms may include:

- skin redness
- blisters
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have:**

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss

**Drug Facts (continued)**

• nausea or vomiting

• stomach pain

Ask a doctor or pharmacist before use if you are:

- taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

• adults: 18 years of age and older

- this product is to be used once a day (every 24 hours), every day for 14 days
- if may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

**Drug Facts (continued)**

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

**Other information**

- read the directions and warnings before use
- keep the cap on. It contains important information.
- store at 20° - 25°C (68° - 77°F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

**Inactive ingredients**  
D&C Red 28, FD&C Blue No. 1, FD&C Green 3, FD&C Red 40, gelatin, hydroxypropyl cellulose, iron oxide black, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer, polyethylene glycol 6000, polyorbate 80, sodium lauryl sulfate, starch (corn), sucrose, sugar spheres, talc, titanium dioxide

**Questions or comments?**  
call 1-888-325-3264



<b>LANSOPRAZOLE</b>			
lansoprazole capsule, delayed release			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:46122-744
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>	
Lansoprazole (UNII: 0K5C5T2QPG) (Lansoprazole - UNII:0K5C5T2QPG)	Lansoprazole	15 mg	
<b>Inactive Ingredients</b>			
<b>Ingredient Name</b>	<b>Strength</b>		
D&C RED NO. 28 (UNII: 7671P0Y5NH)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
GELATIN (UNII: 2G86QN327L)			
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)			
FERROSO FERRIC OXIDE (UNII: XMOM87F357)			
LOW-SUBSTITUTED HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 2165RE0K14)			
MAGNESIUM CARBONATE (UNII: 0E53J927NA)			
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)			
POLYETHYLENE GLYCOL 6000 (UNII: 30IQX730WE)			
POLYSORBATE 80 (UNII: 6OZP39ZG8H)			
SODIUM LAURYL SULFATE (UNII: 368GB5141J)			

<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	BLUE (opaque cyan colored cap) , PINK (opaque pink colored body)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	3mm
<b>Flavor</b>		<b>Imprint Code</b>	RDY;398
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-744-74	1 in 1 CARTON	08/30/2023	
1		14 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:46122-744-04	3 in 1 CARTON	08/30/2023	
2		14 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202194	04/12/2022	

### Labeler - AMERISOURCEBERGEN DRUG CORPORATION (007914906)

Revised: 7/2025

AMERISOURCEBERGEN DRUG CORPORATION