# WINCO FOODS ULTRA STRENGTH STOMACH RELIEF- bismuth subsalicylate suspension WINCO FOODS

-----

## WinCo FOODS Ultra Strength Stomach Relief Drug Facts

### Active ingredient (in each 15 mL dose)

Bismuth subsalicylate 525 mg

## **Purposes**

Upset stomach reliever and antidiarrheal

#### Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching
- fullness

## Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea or vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

## Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

## Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

## Ask a doctor before use if you have

- fever
- mucus in the stool

## Ask a doctor or pharmacist before use if you are

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

## When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

## Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

## If pregnant or breast-feeding,

ask a health professional before use.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away at (1-800-222-1222).

#### **Directions**

- shake well before use
- only use dose cup provided
- adults and children 12 years and over: 15 mL (1 dose) every 1/2 or 30 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea.
- 15 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (120 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

#### Other information

- each 15 mL contains: sodium 10 mg
- salicylate 230 mg
- low sodium
- sugar free
- store at room temperature

- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

## **Inactive ingredients**

carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

#### **Questions?**

Call 800-824-1706 Monday - Friday 9am - 4pm MST

DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN.

## **Principal Display Panel**

WinCo® FOODS

## Compare to active ingredient PEPTO-BISMOL® Ultra Strength

**ULTRA STRENGTH** 

Stomach Relief

BISMUTH SUBSALICYLATE


**UPSET STOMACH RELIEVER & ANTI-DIARRHEAL** 

Soothing Relief for: Upset Stomach, Nausea, Heartburn & Diarrhea

#### **DISTRIBUTED BY:**

WINCO FOODS, LLC, BOISE, ID 83704

\*This product is not manufactured or distributed by PROCTER & GAMBLE INC., distributor of PEPTO-BISMOL $^{\circledR}$  Ultra Strength



If pregnant or breast-feeding, ask a health professional before use

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

Directions ■ shake well before use ■ only use dose cup provided ■ adults and children 12 years and over ■ 15 mL (1 dose) every ½ hour or 30mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea. ■ 15 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ do not exceed 8 doses (120 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information = each 15 mL contains: sodium 10 mg = salicylate 230 mg = low sodium = sugar free = store at room temperature = protect from freezing = avoid excessive heat (over than 104°F or 40°C)

Inactive ingredients carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

Questions? Call 800-824-1706 Monday - Friday 9am - 4pm MST

UNVARNISHED AREA DISTRIBUTED BY:
WINCO FOODS, LLC, BOISE, ID 83704

Winco

ULTRA STRENGTH

Stomach

Relief

BISMUTH SUBSALICYLATE/
UPSET STOMACH RELIEVER & ANTI-DIARRHEAL

Soothing Relief for: Upset Stomach,
Indigestion, Nausea, Heartburn & Diarrhea

Do not use if imprinted shrink band is missing or broken

#### **Drug Facts**

Active ingredient (in each 15 mL dose) Purpose

sismuth subsalicylate 525 mg......Upset stomach relieve and antidiarrheal

Uses relieves = travelers' diarrhea = diarrhea = upset stomach due to overindulgence in food and drink, including: = heartburn = indigestion = nausea = gas = belching = fullness

#### Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are allergic to salicylates (including aspirin)

taking other salicylate products

Do not use if you have man ulcer ma bleeding problem mbloody or black stool

Ask a doctor before use if you have m fever m mucus in the stool
Ask a doctor or pharmacist before use if you are taking any
drug for m anticoagulation (thinning the blood) m diabetes
gout arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if m symptoms get worse or last more than 2 days m ringing in the ears or loss of hearing occum diarrhea lasts more than 2 days

\*This product is not manufactured or distributed by PROCTER & GAMBLE INC., distributor of PEPTO-BISMOL® Ultra Strength.

## WINCO FOODS ULTRA STRENGTH STOMACH RELIEF

bismuth subsalicylate suspension

#### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:67091-363

Route of Administration ORAL

# Active Ingredient/Active Moiety

Ingredient Name

Basis of Strength

Strength

BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD118YE, SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD118YE, SUBSALICYLATE in 15 mL

## **Inactive Ingredients**

Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K6790BS311)	
DS.C PED NO. 22 (LINII) 1679DKY9DT)	

**D&C RED NO. 22** (UNII: 1678RKX8RT) **D&C RED NO. 28** (UNII: 767IP0Y5NH)

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)

POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)

WATER (UNII: 059QF0KO0R)

SALICYLIC ACID (UNII: O414PZ4LPZ)

DIMETHICONE (UNII: 92RU3N3Y1O)

SODIUM BENZOATE (UNII: OJ245FE5EU)

SUCRALOSE (UNII: 96K6UQ3ZD4)

XANTHAN GUM (UNII: TTV12P4NEE)

l	P	ackaging			
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
l	1	NDC:67091-363- 24	236 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/25/2020	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	02/25/2020	

# Labeler - WINCO FOODS (056098817)

Revised: 12/2023 WNCO FOODS