

**GENCARE HIVES RELIEF- fexofenadine hydrochloride tablet**  
**Pioneer Life Sciences, LLC**

---

**Gencare Hives Relief**

**Active Ingredient (in each caplet)**

Fexofenadine HCl 180 mg

**Purpose**

Antihistamine

**Uses**

reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

**Warnings**

**Severe Allergy Warning:**

Get emergency help immediately if you have hives along with any of the following symptoms:

- trouble swallowing
- dizziness or loss of consciousness
- swelling of tongue
- swelling in or around mouth
- trouble speaking
- drooling
- wheezing or problems breathing

These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.

**Not a Substitute for Epinephrine.** If your doctor has prescribed an epinephrine injector for "anaphylaxis or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.

**Do not use**

- to prevent hives from any known cause such as:
- foods ■ insect stings ■ medicines ■ latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious.

If you do not know the cause of your hives, see your doctor for a medical exam. Your

doctor may be able to help you find a cause.

■ If you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

■ kidney disease. Your doctor should determine if you need a different dose. ■ hives that are an unusual color, look bruised or blistered ■hives that do not itch

### **When using this product**

■do not take more than directed  
■do not take at the same time as aluminum or magnesium antacids  
■do not take with fruit juices (see Directions)

### **Stop use and ask a doctor if**

■ an allergic reaction to this product occurs. Seek medical help right away.  
■ symptoms do not improve after 3 days of treatment  
■ the hives have lasted more than 6 weeks

**If pregnant or breast feeding,** ask a health professional before use.

**Keep out of reach of children.** in case of overdose, get medical help or contact a poison control center right away.

### **Directions**

adults and children 12 years of age and over: take one 180mg tablet with water once a day; do not take more than 1 tablet in 24 hours

■children under 12 years of age: do not use  
■adults 65 years of age and older: ask a doctor  
■consumers with kidney disease: ask a doctor

### **Other information**


■ Store between 20°C and 25 °C (68 °and 77°F)  
■ Protect from excessive moisture

### **Inactive ingredient**

Colloidal Silicon Dioxide, Croscarmellose Sodium Hypromellose, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol/macrogol, Povidone, Pregelatinized Starch, Red iron oxide, Silica, Titanium dioxide, Yellow iron oxide.

## Questions or comments

Call +1 (732) 994-2808 or email support@gencare.health



### Drug Facts (cont.)

**Not a Substitute for Epinephrine.** If your doctor has prescribed an epinephrine injector for anaphylaxis or severe allergic symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.

#### Do not use

- to prevent hives from any known cause such as:
  - foods
  - insect stings
  - medicines
  - latex or rubber gloves
- because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious.
- If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.
- If you have ever had an allergic reaction to this product or any of its ingredients.

#### Ask a doctor before use if you have

kidney disease. Your doctor should determine if you need a different dose. ■ hives that are an unusual color, look bruised or blistered. ■ hives that do not itch

**When using this product** ■ do not take more than directed

- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

**Stop use and ask a doctor if** ■ an allergic reaction to this product occurs. Seek medical help right away.

- symptoms do not improve after 3 days of treatment
- the hives have lasted more than 6 weeks

**If pregnant or breast feeding**, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a poison control center right away.

**Directions:** adults and children 12 years of age and over: take one 180mg tablet with water once a day, do not take more than 1 tablet in 24 hours.

- children under 12 years of age: do not use
- adults 65 years of age and older: ask a doctor
- consumers with kidney disease: ask a doctor

#### Other information:

- Store between 20°C and 25°C (68 ° and 77°F)
- Protect from excessive moisture

#### Inactive ingredient:

Colloidal Silicon Dioxide, Croscarmellose Sodium, Hydroxyose, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol/Macrogol, Povidone, Pre-gelatinized Starch, Red Iron Oxide, Silica, Titanium Dioxide, Yellow Iron Oxide.

#### Questions or comments:

Call +1 (732) 994-2808 or email support@gencare.health

## GENCARE HIVES RELIEF

fexofenadine hydrochloride tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72090-058
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>FEXOFENADINE HYDROCHLORIDE</b> (UNII: 2S068B75ZU) (FEXOFENADINE - UNII: E6582LOH6V)	FEXOFENADINE HYDROCHLORIDE	180 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	

## Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	capsule	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	180
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72090-058-01	48 in 1 BOTTLE; Type 0: Not a Combination Product	01/21/2026	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210137	01/21/2026	

**Labeler** - Pioneer Life Sciences, LLC (014092742)

Revised: 1/2026

Pioneer Life Sciences, LLC