

**MUCINEX SINUS-MAX SEVERE CONGESTION AND PAIN MAXIMUM STRENGTH-  
acetaminophen, dextromethorphan hydrobromide, and phenylephrine  
hydrochloride capsule, liquid filled  
Reckitt Benckiser LLC**

-----  
**Mucinex® Sinus-Max® Severe Congestion & Pain  
Maximum Strength**

***Drug Facts***

<b><i>Active ingredients (in each liquid gel)</i></b>	<b><i>Purposes</i></b>
<b>Acetaminophen 325 mg</b>	<b>Pain reliever</b>
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

- temporarily relieves:
  - nasal congestion
  - headache
  - minor aches and pains
  - sinus congestion and pressure
  - cough
- temporarily promotes nasal and/or sinus drainage

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's Disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

### **When using this product do not use more than directed**

#### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

### **Overdose warning**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

### **Other information**

- store at 20-25°C (68-77°F)
- avoid excessive heat

### **Inactive ingredients**

FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water

### **Questions?**

#### **1-866-MUCINEX (1-866-682-4639)**

You may also report side effects to this phone number.

Dist. by: RB Health (US)  
Parsippany, NJ 07054-0224

### **PRINCIPAL DISPLAY PANEL - 16 Capsule Blister Pack Carton**

Fast Dissolving Liquid Gels!

NDC 72854-203-16

MAXIMUM STRENGTH

Mucinex®  
SINUS-MAX®

SEVERE CONGESTION  
& PAIN

Acetaminophen – Pain Reliever  
Dextromethorphan HBr – Cough Suppressant  
Phenylephrine HCl – Nasal Decongestant

- Clears Sinus Congestion
- Relieves Headache
- Controls Cough

Actual Size

16  
LIQUID GELS  
(Liquid Filled Capsules)

DAY TIME  
FOR AGES 12+

MAXIMUM STRENGTH

**Mucinex**  
**SINUS-MAX**

SEVERE CONGESTION  
& PAIN

16 LIQUID GELS

Tamper evident: Do not use if carton is damaged  
or if printed seal on blister is broken or missing.

*Fast Dissolving Liquid Gels!*

NDC 72854-203-16

MAXIMUM STRENGTH

**Mucinex**  
**SINUS-MAX**

SEVERE CONGESTION  
& PAIN

SEVERE CONGESTION  
& PAIN

**Mucinex**  
**SINUS-MAX**

MAXIMUM STRENGTH

Dist. by: RB Health (US)  
Passippany, NJ 07054-0224  
©2021 RB Health

030221 3176792

Maximum Strength per 4-hour dose  
Do not take more than a total of  
12 liquid gels in a 24-hour period.  
Take only as directed.  
Keep carton for full information.

**Acetaminophen – Pain Reliever**  
**Dextromethorphan HBr – Cough Suppressant**  
**Phenylephrine HCl – Nasal Decongestant**

- ✓ Clears Sinus Congestion
- ✓ Relieves Headache
- ✓ Controls Cough

Actual Size



16 LIQUID GELS  
(Liquid Filled Capsules)

DAY TIME  
FOR AGES 12+



HEALTH • HYGIENE • HOME  
[www.mucinex.com](http://www.mucinex.com)  
Patents: [www.rb.com/patents](http://www.rb.com/patents)


**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

LOT:

EXP.:

3176792





**Drug Facts**

**Active ingredients (in each liquid gel)**

**Purposes**

Acetaminophen 325 mg .....Pain reliever  
 Dextromethorphan HBr 10 mg.....Cough suppressant  
 Phenylephrine HCl 5 mg.....Nasal decongestant

**Uses**

- temporarily relieves:
  - nasal congestion
  - headache
  - minor aches and pains
  - sinus congestion and pressure
  - cough
- temporarily promotes nasal and/or sinus drainage

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:  
 ■ more than 12 liquid gels in 24 hours, which is the maximum daily amount  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks daily while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ skin reddening ■ blisters ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's Disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

**Drug Facts (continued)**

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

**Other information**

- store at 20-25°C (68-77°F)
- avoid excessive heat

**Inactive ingredients**

FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water

**Questions? 1-866-MUCINEX (1-866-682-4639)**

You may also report side effects to this phone number.

**MUCINEX SINUS-MAX SEVERE CONGESTION AND PAIN MAXIMUM STRENGTH**

acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride capsule, liquid filled

**Product Information**

**Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:72854-203

<b>Route of Administration</b>	ORAL
--------------------------------	------

<b>Active Ingredient/Active Moiety</b>		
<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

<b>Inactive Ingredients</b>	
<b>Ingredient Name</b>	<b>Strength</b>
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>WATER</b> (UNII: 059QF0KOOR)	

<b>Product Characteristics</b>			
<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	24mm
<b>Flavor</b>		<b>Imprint Code</b>	AR04
<b>Contains</b>			

<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-203-16	2 in 1 CARTON	07/26/2021	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/26/2021	

