

SENNA-S- docusate sodium and sennosides tablet, coated RITE AID

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

192R Rite aid 11822 3065 DOC SOD 50MG/SENNOSIDES 8.6MG TABLETS

DRUG FACTS

Active ingredients (in each tablet)

Docustate sodium 50 mg

Sennosides 8.6 mg

Purposes

Stool softener

Stimulant laxative

Uses

- relieves occasional constipation (irregularity)
- generally causes bowel movement in 6-12 hours

Warnings

Do not use

- this product if you are presently taking mineral oil, unless directed by a doctor
- laxative products for longer than 1 week unless directed by a doctor

Stop use and ask a doctor if you have rectal bleeding or fail to have to have a bowel movement after use of a laxative. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take preferably at bedtime or as directed by a doctor
- if you do not have a comfortable bowel movement by the second day, increase dose by one tablet (do not exceed maximum dosage) or decrease dose until you are comfortable

age	starting dosage	maximum dosage
adults and children 12 years and over	2 tablets once a day	4 tablets twice a day
children 6 to under 12 years	1 tablet once a day	2 tablets twice a day

Lot No.:
Exp Date:

LAXATIVE

DOCUSATE SODIUM 50 mg
SENNOSIDES 8.6 mg

Print/Varnish
Omit Area

Drug Facts (continued)
Other information
■ each tablet contains: calcium 20 mg, sodium 6 mg (LOW SODIUM)
■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
Inactive ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, dibasic calcium phosphate dihydrate, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, sodium benzoate, stearic acid, titanium dioxide
Questions or comments?
Call 1-877-290-4008

Compare to the active ingredients in **Senokot-S®**
NDC 11822-3065-9

LAXATIVE

DOCUSATE SODIUM 50 mg
SENNOSIDES 8.6 mg

NATURAL VEGETABLE
STIMULANT LAXATIVE
STOOL SOFTENER
Gentle, overnight relief

RETAIN CARTON FOR COMPLETE
PRODUCT INFORMATION

Drug Facts
Active ingredients (in each tablet)
Docusate sodium 50 mg.....Stool softener
Sennosides 8.6 mg.....Stimulant laxative
Purposes
Uses
■ relieves occasional constipation (irregularity)
■ generally causes bowel movement in 6-12 hours
Warnings
Do not use
■ if you are now taking mineral oil, unless directed by a doctor
■ laxative products for longer than 1 week unless directed by a doctor
Ask a doctor before use if you have
■ stomach pain ■ nausea ■ vomiting
■ noticed a sudden change in bowel habits that continues over a period of 2 weeks
Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative.
These may indicate a serious condition. ▶

Drug Facts (continued)
If pregnant or breast-feeding, ask a health professional before use.
Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ take preferably at bedtime or as directed by a doctor
■ if you do not have a comfortable bowel movement by the second day, increase dose by one tablet (do not exceed maximum dosage) or decrease dose until you are comfortable

Age	Starting dose	Maximum dosage
Adults and children 12 years of age and over	2 tablets once a day	4 tablets twice a day
Children 6 to under 12 years	1 tablet once a day	2 tablets twice a day
Children 2 to under 6 years	1/2 tablet once a day	1 tablet twice a day
Children under 2 years	ask a doctor	ask a doctor

*This product is not manufactured or distributed by Avrio Health L.P., owner of the registered trademark Senokot-S®. 192R 0123

PACKAGE NOT CHILD-RESISTANT

ACTUAL SIZE
30 TABLETS

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING



DISTRIBUTED BY: RITE AID,
200 NEWBERRY COMMONS
ETTERS, PA 17319
www.riteaid.com MADE IN U.S.A.
SATISFACTION GUARANTEE:
If you're not satisfied,
we'll happily refund your money.

Print/Varnish
Omit Area

0783 Varnish/Print Omit Area

NDC 11822-3065-3

LAXATIVE

DOCUSATE SODIUM 50 mg
SENNOSIDES 8.6 mg

NATURAL VEGETABLE
STIMULANT LAXATIVE
STOOL SOFTENER
Gentle, overnight relief



60 TABLETS

RETAIN CARTON FOR COMPLETE
PRODUCT INFORMATION

Uses ■ relieves occasional constipation (irregularity)
■ generally causes bowel movement in 6-12 hours
Warnings
Do not use ■ this product if you are presently taking mineral oil, unless directed by a doctor ■ laxative products for longer than 1 week unless directed by a doctor
Ask a doctor before use if you have ■ stomach pain ■ nausea ■ vomiting ■ a sudden change in bowel movements that persists over 2 weeks
Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions ■ take preferably at bedtime or as directed by a doctor ■ if you do not have a comfortable bowel movement by the second day, increase dose by one tablet (do not exceed maximum dosage) or decrease dose until you are comfortable
■ adults and children 12 years and over: ■ starting dosage 2 tablets once a day ■ maximum dosage 4 tablets twice a day
■ children 6 to under 12 years: ■ starting dosage 1 tablet once a day ■ maximum dosage 2 tablets twice a day
■ children 2 to under 6 years: ■ starting dosage 1/2 tablet once a day ■ maximum dosage 1 tablet twice a day
■ children under 2 years: ■ ask a doctor
Other information ■ TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING ■ each tablet contains: calcium 20 mg, sodium 6 mg (LOW SODIUM) ■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
Questions or comments? Call 1-877-290-4008
DISTRIBUTED BY: RITE AID
30 HUNTERLANE CAMP HILL, PA 17011 192R 0321
www.riteaid.com

Varnish
Omit Area

Lot No.:
Exp. Date:



Drug Facts (continued)

Other information

- each tablet contains: calcium 20 mg, sodium 6 mg (LOW SODIUM)
- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, dibasic calcium phosphate dihydrate, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, sodium benzoate, stearic acid, titanium dioxide

Questions or comments?
Call 1-877-290-4008

*This product is not manufactured or distributed by Avrio Health L.P., owner of the registered trademark Senokot-SP. 192R 0321

DISTRIBUTED BY: RITE AID
30 HUNTER LANE CAMP HILL, PA 17011
www.riteaid.com

SATISFACTION GUARANTEE
If you're not satisfied, we'll happily refund your money.

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Coating/Print Omit Area



Compare to the active ingredients in Senokot-SP

LAXATIVE

DOCUSATE SODIUM 50 mg • SENNOSIDES 8.6 mg
NATURAL VEGETABLE STIMULANT LAXATIVE
STOOL SOFTENER
Gentle, overnight relief

60 TABLETS

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredients (in each tablet)

Docusate sodium 50 mg, Stool softener
Sennosides 8.6 mg, Stimulant laxative

Uses
■ relieves occasional constipation (irregularity)
■ generally causes bowel movement in 6-12 hours

Warnings
Do not use
■ this product if you are presently taking mineral oil, unless directed by a doctor
■ laxative products for longer than 1 week unless directed by a doctor

Ask a doctor before use if you have
■ stomach pain ■ nausea ■ vomiting
■ a sudden change in bowel movements that persists over 2 weeks

Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.

Drug Facts (continued)

if pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ take preferably at bedtime or as directed by a doctor
■ if you do not have a comfortable bowel movement by the second day, increase dose by one tablet (do not exceed maximum dosage) or decrease dose until you are comfortable

age	starting dosage	maximum dosage
adults and children 12 years and over	2 tablets once a day	4 tablets twice a day
children 6 to under 12 years	1 tablet once a day	2 tablets twice a day
children 2 to under 6 years	1/2 tablet once a day	1 tablet twice a day
children under 2 years	ask a doctor	ask a doctor

E826R1 Coating/Print Omit Area

Coating Omit

NDC 11822-3065-6

LAXATIVE

DOCUSATE SODIUM 50 mg
SENNOSIDES 8.6 mg

NATURAL VEGETABLE
STIMULANT LAXATIVE
STOOL SOFTENER

Gentle, overnight relief



120 TABLETS

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Uses ■ relieves occasional constipation (irregularity)
■ generally causes bowel movement in 6-12 hours

Warnings
Do not use ■ this product if you are presently taking mineral oil, unless directed by a doctor ■ laxative products for longer than 1 week unless directed by a doctor

Ask a doctor before use if you have ■ stomach pain ■ nausea ■ vomiting
■ a sudden change in bowel movements that persists over 2 weeks
Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ take preferably at bedtime or as directed by a doctor
■ if you do not have a comfortable bowel movement by the second day, increase dose by one tablet (do not exceed maximum dosage) or decrease dose until you are comfortable

age	starting dosage	maximum dosage
adults and children 12 years of age and older	2 tablets once a day	4 tablets twice a day
children 6 to under 12 years	1 tablet once a day	2 tablets twice a day
children 2 to under 6 years	1/2 tablet once a day	1 tablet twice a day
children under 2 years	ask a doctor	ask a doctor

Other information
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING
■ each tablet contains: calcium 20 mg, sodium 6 mg (LOW SODIUM)
■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
Questions or comments? Call 1-877-290-4008

SATISFACTION GUARANTEE
If you're not satisfied, we'll happily refund your money.

DISTRIBUTED BY: RITE AID
30 HUNTER LANE CAMP HILL, PA 17011
www.riteaid.com

192F 0321

Varnish Omit Area

Lot No.:
Exp. Date:



SENNAS-S

docusate sodium and sennosides tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-3065
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SENNOSIDES (UNII: 3FYP5M0IJX) (SENNOSIDES - UNII:3FYP5M0IJX)	SENNOSIDES	8.6 mg
DOCUSATE SODIUM (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG)	DOCUSATE SODIUM	50 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	TCL081
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-3065-3	60 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2020	
2	NDC:11822-3065-6	120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/01/2020	
3	NDC:11822-3065-9	1 in 1 CARTON	04/15/2023	
3		30 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	06/01/2020	

Labeler - RITE AID (014578892)

Registrant - TIME CAP LABORATORIES, INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

Revised: 4/2023

RITE AID