

TOBRAMYCIN- tobramycin solution/ drops
Preferred Pharmaceuticals Inc.

Rx Only
For Topical Ophthalmic Use Only

DESCRIPTION

Tobramycin ophthalmic solution 0.3% is a sterile topical ophthalmic antibiotic formulation prepared specifically for topical therapy of external ophthalmic infections.

Each mL of Tobramycin ophthalmic solution USP, 0.3% contains: Active: tobramycin 0.3% (3 mg). **Preservative:** benzalkonium chloride 0.01% (0.1 mg). **Inactives:** boric acid, sodium sulfate, sodium chloride, tyloxapol, sodium hydroxide and/or sulfuric acid (to adjust pH) and water for injection. Tobramycin ophthalmic solution 0.3% has a pH range between 7.0 and 8.0 and an osmolality of 260-320 mOsm/kg.

Tobramycin is a water-soluble aminoglycoside antibiotic active against a wide variety of gram- negative and gram-positive ophthalmic pathogens.

The chemical structure of tobramycin is:

Molecular Weight = 467.52 Molecular Formula:

$C_{18}H_{37}N_5O_9$

Chemical Name:

0-{3-amino-3-deoxy- α -D-gluco-pyranosyl (1 \rightarrow 4) }-0-{2,6-diamino-2,3,6-trideoxy α -D-ribohexo-pyranosyl-(1 \rightarrow 6) }-2 deoxystreptamine.

CLINICAL PHARMACOLOGY

In Vitro Data: *In vitro* studies have demonstrated tobramycin is active against susceptible strains of the following microorganisms: *Staphylococci*, including *S. aureus* and *S. epidermidis* (coagulase-positive and coagulase-negative), including penicillin-resistant strains.

Streptococci, including some of the Group A-beta-hemolytic species, some nonhemolytic species, and some *Streptococcus pneumoniae*.

Pseudomonas aeruginosa, *Escherichia coli*, *Klebsiella pneumoniae*, *Enterobacter aerogenes*, *Proteus mirabilis*, *Morganella morganii*, most *Proteus vulgaris* strains, *Haemophilus influenzae* and *H. aegyptius*, *Moraxella lacunata*, *Acinetobacter calcoaceticus* and some *Neisseria* species.

Bacterial susceptibility studies demonstrate that in some cases, microorganisms resistant to gentamicin retain susceptibility to tobramycin.

INDICATIONS AND USAGE

Tobramycin ophthalmic solution 0.3% is a topical antibiotic indicated in the treatment of external infections of the eye and its adnexa caused by susceptible bacteria. Appropriate monitoring of bacterial response to topical antibiotic therapy should accompany the use of tobramycin ophthalmic solution 0.3%. Clinical studies have shown tobramycin to be safe and effective for use in children.

CONTRAINDICATIONS

Tobramycin ophthalmic solution 0.3% is contraindicated in patients with known hypersensitivity to any of its components.

WARNINGS

FOR TOPICAL OPHTHALMIC USE. NOT FOR INJECTION INTO THE EYE. Sensitivity to topically applied aminoglycosides may occur in some patients. Severity of hypersensitivity reactions may vary from local effects to generalized reactions such as erythema, itching, urticaria, skin rash, anaphylaxis, anaphylactoid reactions, or bullous reactions. If a sensitivity reaction to tobramycin ophthalmic solution 0.3% occurs, discontinue use.

PRECAUTIONS

General: As with other antibiotic preparations, prolonged use may result in overgrowth of nonsusceptible organisms, including fungi. If superinfection occurs, appropriate therapy should be initiated.

Cross-sensitivity to other aminoglycoside antibiotics may occur; if hypersensitivity develops with this product, discontinue use and institute appropriate therapy. Patients should be advised not to wear contact lenses if they have signs and symptoms of bacterial ocular infection.

Information for Patients: Do not touch dropper tip to any surface, as this may contaminate the solution.

Pregnancy: Reproduction studies in 3 types of animals at doses up to 33 times the normal human systemic dose have revealed no evidence of impaired fertility or harm to the fetus due to tobramycin. There are, however, no adequate and well-controlled studies in pregnant women. Because animal studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers: Because of the potential for adverse reactions in nursing infants from tobramycin ophthalmic solution 0.3%, a decision should be made whether to discontinue nursing the infant or discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use: Safety and effectiveness in pediatric patients below the age of 2 months has not been established.

Geriatric Use: No overall differences in safety or effectiveness have been observed between elderly and younger patients.

ADVERSE REACTIONS

The most frequent adverse reactions to tobramycin ophthalmic solution 0.3% are hypersensitivity and localized ocular toxicity, including lid itching and swelling, and conjunctival erythema. These reactions occur in less than three of 100 patients treated with tobramycin ophthalmic solution 0.3%.

Postmarketing Experience: Additional adverse reactions identified from post-marketing use include anaphylactic reaction, Stevens-Johnson syndrome, and erythema multiforme.

The following additional adverse reactions have been reported with systemic aminoglycosides:

Neurotoxicity, ototoxicity and nephrotoxicity have occurred in patients receiving systemic aminoglycoside therapy. Aminoglycosides may aggravate muscle weakness in patients with known or suspected neuromuscular disorders, such as myasthenia gravis or Parkinson's disease, because of their potential effect on neuromuscular function.

DOSAGE AND ADMINISTRATION

In mild to moderate disease, instill 1 or 2 drops into the affected eye(s) every 4 hours. In severe infections, instill 2 drops into the eye(s) hourly until improvement, following which treatment should be reduced prior to discontinuation.

HOW SUPPLIED

Tobramycin ophthalmic solution USP, 0.3% is supplied as a 5 mL sterile solution,

packaged in a 10 mL white LDPE bottle and natural LDPE nozzle and White HDPE cap as follows:

5 mL containing tobramycin 0.3% (3 mg/mL) NDC 68788-7252-5.

Storage: Store at 2° to 25°C (36° to 77°F).

After opening, tobramycin ophthalmic solution 0.3% can be used until the expiration date on the bottle.

For Product Inquiry call +1-800-417-9175

Revised: June, 2021

Manufactured for:

Somerset Therapeutics, LLC

Hollywood, FL 33024

Made in India

Code No.: KR/DRUGS/KTK/28/289/97

PSSO0377

ST-TBR11/P/05


Relabeled By: Preferred Pharmaceuticals Inc.,

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

Tobramycin Ophthalmic Solution 0.3%

5 mL

Container Label

<p>Tobramycin Ophthalmic Sol. USP 0.3% Generic for Tobrex</p> <p>Each mL contains: Tobramycin 0.3% (3mg)</p> <p>Pkg Size: Exp Date: ####/#### Lot#: Batch#:</p> <p>Ins: Mfg: Somerset Therapeutics, LLC Prod#:</p> <p>Warning Store between 2° - 25°C (36° - 77°F). Do not touch dropper tip to any surface as this may contaminate the solution. For use in the eye. RX Only. Avoid exposure to excessive heat. Keep out of reach of children. For ophthalmic use only. Do not use if imprinted neckband is not intact.</p>	 <p>Directions English</p> <p>Use as directed by your doctor Instill _____ drops every _____ hours.</p>	<p>CAUTION: Federal law PROHIBITS transfer of this drug to any person other than the patient for whom it was prescribed.</p>  <p>GTIN ##### SN ##### EXP ####/####</p>	<p>Instrucciones Espanol: Usó según lo dirigido por su doctor Póngase _____ gota(s) cada _____ horas.</p>	<p>Tobramycin Ophthalmic Sol. USP 0.3% Qty: Ins: Lot: Bat: Prod# (NDC):</p> <p>Tobramycin Ophthalmic Sol. USP 0.3% Qty: Ins: Lot: Bat: Prod# (NDC):</p> <p>Tobramycin Ophthalmic Sol. USP 0.3% Qty: Insurance NDC: Lot: Bat:</p> <p>Tobramycin Ophthalmic Sol. USP 0.3% Qty: Ins: Lot: Bat: Prod# (NDC):</p> <p>Log Chart Billing Patient</p>
---	---	---	--	---

TOBRAMYCIN
tobramycin solution/ drops

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:68788-7252(NDC:70069-131)
Route of Administration	OPHTHALMIC		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TOBRAMYCIN (UNII: VZ8RRZ51VK) (TOBRAMYCIN - UNII:VZ8RRZ51VK)	TOBRAMYCIN	3 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	
BORIC ACID (UNII: R57ZHV85D4)	
SODIUM SULFATE ANHYDROUS (UNII: 36KCS0R750)	
TYLOXAPOL (UNII: Y27PUL9H56)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SULFURIC ACID (UNII: O40UQP6WCF)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	WHITE (Clear, colorless solution)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68788-7252-5	1 in 1 CARTON	08/29/2018	
1		5 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA207444	08/29/2018	

Labeler - Preferred Pharmaceuticals Inc. (791119022)

Registrant - Preferred Pharmaceuticals Inc. (791119022)

Establishment

Name	Address	ID/FEI	Business Operations
Preferred Pharmaceuticals Inc.		791119022	RELABEL(68788-7252)

Revised: 9/2025

Preferred Pharmaceuticals Inc.