## DG HEALTH CHILDRENS IBUPROFEN- ibuprofen suspension Dolgencorp, LLC

-----

## Dolgencorp, LLC Children's Ibupropfen Drug Facts

#### Active ingredient (in each 5 mL)

Ibuprofen 100 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

#### Purposes

Pain reliever/fever reducer

## Uses

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

#### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more

than directed or for longer than directed.

**Sore throat warning**: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

## Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

## Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

# Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

# When using this product

give with food or milk if stomach upset occurs

## Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

## **Dosing Chart**

Weight (lb)	Age (yr)	Dose (mL)**	
under 24 lbs	under 2 years	ask a doctor	
24-35 lbs	2-3 years	5 mL	
36-47 lbs	4-5 years	7.5 mL	
48-59 lbs	6-8 years	10 mL	
60-71 lbs	9-10 years	12.5 mL	
72-95 lbs	11 years	15 mL	

\*\*or as directed by a doctor

## Other information

- each 5 mL contains: sodium 2 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not freeze

## Inactive ingredients

anhydrous citric acid, D&C red #33, FD&C blue #1, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, natural and artificial grape flavor, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

## **Questions or comments?**

### 1-888-309-9030

## Package/Label Principal Display Panel

SEE NEW WARNINGS For Ages 2 to 11 Years children's Ibuprofen Oral Suspension 100mg per 5mL Pain Reliever/Fever Reducer (NSAID) LASTS UP TO 8 HOURS Alcohol Free Gluten Free Gluten Free 4 FL OZ (118 mL)



DG HEALTH CHILDRENS IBUPROFEN ibuprofen suspension						
Product Information						
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-102			
Route of Administration	ORAL					

		ive Moiety			
	Ing	redient Name	Basis of Streng	th Strength	
BUPROFEN (UNII:	WK2XYI10C	9M) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg in 5 ml	
Inactive Ingre	dients				
	Strength				
ANHYDROUS CITR					
D&C RED NO. 33 (					
FD&C BLUE NO. 1					
FD&C RED NO. 40					
GLYCERIN (UNII: PE					
HIGH FRUCTOSE (					
HYPROMELLOSE,					
POLYSORBATE 80					
WATER (UNII: 059Q					
SODIUM BENZOAT					
SORBITOL (UNII: 50					
XANTHAN GUM (UI	NII: TTV12P	4NEE)			
Color	PURPLE (dark) Se		Score		
Shape			ize		
Flavor		GRAPE	Imprint Code		
Contains					
Packaging		Package Description	Marketing Start Date	Marketing End Date	
# Item Code	1 in 1 CAF	RTON	04/20/2016		
<ul> <li># Item Code</li> <li>1 NDC:55910-102- 26</li> </ul>		RTON 1 BOTTLE; Type 0: Not a Combinatio			
<ul> <li># Item Code</li> <li>1 NDC:55910-102- 26</li> <li>1</li> </ul>	118 mL in Product	1 BOTTLE; Type 0: Not a Combinatio			
<ul> <li># Item Code</li> <li>DDC:55910-102- 26</li> <li>I</li> </ul>	118 mL in Product	1 BOTTLE; Type 0: Not a Combinatio	n		
NDC:55910-102-	118 mL in Product	1 BOTTLE; Type 0: Not a Combinatio	n	Marketing Enc Date	

# Labeler - Dolgencorp, LLC (068331990)