

ACID REDUCER- famotidine tablet
Allegiant Health

443 - Acid Reducer

Active ingredient(s)

Famotidine USP 20mg

Purpose

Acid reducer

Use(s)

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if

you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breastfeeding,

ask a health professional before use.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions**adults and children 12 years and over:**

- to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn n do not use more than 2 tablets in 24 hours
- **children under 12 years:** ask a doctor

Other information

- read the directions and warnings before use
- do not use if imprinted safety seal under cap is broken or missing
- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients

Hydroxypropyl cellulose, hypromellose, macrogol, magnesium stearate, microcrystalline cellulose, pre-gelatinized starch, sodium starch glycolate, talc, titanium dioxide, triacetin

Questions/Comments

Call 1-888-952-0050 Monday through Friday 9AM – 5PM EST

Principal Display Panel



Drug Facts

Active ingredient (in each tablet)
Famotidine USP 20mg.....Acid reducer

Purpose

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Drug Facts (continued on inside)

†This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Maximum Strength Pepcid AC®. Pepcid AC® is a registered trademark of Johnson & Johnson.

Manufactured for:
Allegiant Health
 Deer Park, NY 11729

LB2135
R0925

X0040DLHD3
HealthA2Z® Acid Red... to Acid Indigestion New

Lot: _____
 PEELED HERE
 Exp: _____

UNVARNISHED

Drug Facts (continued)

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath, sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Drug Facts (continued)

Directions

- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
 - children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- do not use if imprinted safety seal under cap is broken or missing
- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients

Hydroxypropyl cellulose, hypromellose, macrogol, magnesium stearate, microcrystalline cellulose, pre-gelatinized starch, sodium starch glycolate, talc, titanium dioxide, triacetin

Questions or Comments?
 Call 1-888-952-0050 Monday through Friday 9AM – 5PM EST

Acid Reducer

ACID REDUCER

famotidine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69168-443
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

Inactive Ingredients

Ingredient Name	Strength
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	V;15
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69168-443-32	100 in 1 BOTTLE; Type 0: Not a Combination Product	01/15/2024	
2	NDC:69168-443-52	225 in 1 BOTTLE; Type 0: Not a Combination Product	01/15/2024	
3	NDC:69168-443-09	10 in 1 BLISTER PACK; Type 0: Not a Combination Product	02/29/2024	
4	NDC:69168-443-50	1 in 1 CARTON	03/05/2024	
4		50 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:69168-443-99	365 in 1 BOTTLE; Type 0: Not a Combination Product	10/28/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215822	01/15/2024	

