

**ORAJEL 2X GUM PAIN ALCOHOL-FREE- menthol and hydrogen peroxide rinse**  
**Church & Dwight Co., Inc.**

-----  
**Orajel™ 2x Gum Pain Rinse**  
**Alcohol-Free**

***Drug Facts***

<b><i>Active ingredients</i></b>	<b><i>Purpose</i></b>
Hydrogen peroxide 1.5%	Oral debriding agent/Oral antiseptic
Menthol 0.1%	Oral pain reliever

**Uses**

- first aid to help reduce bacteria in minor oral wounds
- for temporary pain relief and
- use in cleansing minor wounds or minor gum inflammation resulting from:
  - minor dental procedures
  - accidental injury
  - orthodontic appliances
  - canker sores
  - dentures
  - other irritations of the mouth and gums

Aids in the removal of:

- phlegm
- mucus
- other secretions associated with occasional sore mouth

**Warnings**

**Do not use** this product for more than 7 days unless directed by a dentist or healthcare provider

**When using this product** do not swallow

- do not exceed recommended dosage

**Stop use and see your physician promptly if**

- swelling, rash or fever develops
- irritation, pain or redness persists or worsens
- sore mouth symptoms do not improve in 7 days

**Keep out of reach of children.** If more than used for rinsing is accidentally swallowed, get medical help or contact a Poison Control Center right away

**Directions**

- remove imprinted safety seal from bottle cap
- to remove child-resistant cap, squeeze smooth sides of cap while turning. Reclose tightly. Ready to use, no mixing needed.

Adults and children 2 years of age and older	Swish one-half capful (2 teaspoons = 10mL) around the mouth over the affected area for at least 1 minute and then spit out. Use up to 4 times daily after meals and at bedtime or as directed by a dentist or healthcare provider
Children under 12 years of age	Should be supervised in the use of this product
Children under 2 years of age	Consult a dentist or healthcare provider

### Other information

- cap tightly
- keep away from heat or direct sunlight
- do not use if safety seal is broken or missing

### Inactive ingredients

disodium EDTA, FD&C blue no.1, methyl salicylate, phosphoric acid, poloxamer 338, polysorbate 20, propylene glycol, sodium saccharin, sorbitol, water

### Questions or comments?

call us at **1-800-952-5080** M-F 9am-5pm ET or visit our website at [www.oraljel.com](http://www.oraljel.com)

### PRINCIPAL DISPLAY PANEL - 473.2 mL Bottle Label

1#  
ORAL PAIN  
RELIEF BRAND  
FOR ADULTS

Orajel <sup>TM</sup>  
2X  
GUM PAIN RINSE  
MEDICATED  
ALCOHOL FREE

EFFECTIVE ORAL PAIN RELIEF

- Soothes & Cools Gums
- Kills Harmful Bacteria
- Promotes Healing

- Relieves Gum Irritation  
Caused by Dentures or Braces

ORAL DEBRIDING AGENT/  
ANTISEPTIC RINSE/PAIN RELIEVER

Soothing  
MINT

16 FL OZ (473.2 mL)

OJLBF-01904-01 72025286

**#1 ORAL PAIN  
RELIEF BRAND  
FOR ADULTS**

**Orajel™**  
**2X GUM PAIN RINSE**  
**MEDICATED**

**ALCOHOL FREE**

**EFFECTIVE ORAL PAIN RELIEF**

- ✓ Soothes & Cools Gums
- ✓ Kills Harmful Bacteria
- ✓ Promotes Healing
- ✓ Relieves Gum Irritation  
Caused by Dentures or Braces



16 FL. OZ (473.2 mL)

ORAL DEBRIDING AGENT/  
ANTISEPTIC RINSE/PAIN RELIEVER

OJLBF-01904-01 72025286

**Soothing  
MINT**

**Drug Facts**

**Active ingredients Purpose**

Hydrogen peroxide 1.5% ..... Oral debriding agent/  
Oral antiseptic

Menthol 0.1% ..... Oral pain reliever

**Uses** ■ first aid to help reduce bacteria in minor oral  
wounds ■ for temporary pain relief and ■ use in cleansing

relieves ■ for temporary pain relief and ■ use in cleaning minor wounds or minor gum inflammation resulting from:

- minor dental procedures
- orthodontic appliances
- dentures
- accidental injury
- canker sores
- other irritations of the mouth and gums

Aids in the removal of: ■ phlegm ■ mucus ■ other secretions associated with occasional sore mouth

**Warnings**

Do not use this product for more than 7 days unless directed by a dentist or healthcare provider

When using this product do not swallow ■ do not exceed recommended dosage

Stop use and see your physician promptly if ■ swelling, rash or fever develops ■ irritation, pain or redness persists or worsens ■ sore mouth symptoms do not improve in 7 days

Keep out of reach of children. If more than used for rinsing is accidentally swallowed, get medical help or contact a Poison Control Center right away

**Directions** ■ remove imprinted safety seal from bottle cap ■ to remove child-resistant cap, squeeze smooth sides of cap while turning. Reclose tightly. Ready to use, no mixing needed.


Adults and children 2 years of age and older	Swish one-half capful (2 teaspoons = 10mL) around the mouth over the affected area for at least 1 minute and then spit out. Use up to 4 times daily after meals and at bedtime or as directed by a dentist or healthcare provider
Children under 12 years of age	Should be supervised in the use of this product
Children under 2 years of age	Consult a dentist or healthcare provider

**Other information** ■ cap tightly ■ keep away from heat or direct sunlight ■ do not use if safety seal is broken or missing

**Inactive ingredients** disodium EDTA, FD&C blue no.1, methyl salicylate, phosphoric acid, poloxamer 338, polysorbate 20, propylene glycol, sodium saccharin, sorbitol, water

**Questions or comments?** call us at 1-800-952-5080 M-F 9am-5pm ET or visit our website at [www.oraljel.com](http://www.oraljel.com)


**SCAN ME**



LEARN MORE

**SAFETY SEALED**

Church & Dwight Co., Inc.  
Ewing, N.J. 08628  
TM 2021 Church & Dwight Co., Inc.  
0JLBB-01904-01 72025285  
Made in Canada



3 10310 01904 4

## ORAJEL 2X GUM PAIN ALCOHOL-FREE

menthol and hydrogen peroxide rinse

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:10237-799
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>MENTHOL, UNSPECIFIED FORM</b> (UNII: L7T10EIP3A) (MENTHOL, UNSPECIFIED FORM - UNII:L7T10EIP3A)	MENTHOL, UNSPECIFIED FORM	1 mg in 1 mL
<b>HYDROGEN PEROXIDE</b> (UNII: BBX060AN9V) (HYDROGEN PEROXIDE - UNII:BBX060AN9V)	HYDROGEN PEROXIDE	15 mg in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
<b>EDETATE DISODIUM ANHYDROUS</b> (UNII: 8NLQ36F6MM)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>METHYL SALICYLATE</b> (UNII: LAV5U5022Y)	
<b>PHOSPHORIC ACID</b> (UNII: E4GA8884NN)	
<b>POLOXAMER 338</b> (UNII: F75JV2T505)	
<b>POLYSORBATE 20</b> (UNII: 7T1F30V5YH)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>WATER</b> (UNII: 059QF0KO0R)	

Product Characteristics			
Color	blue	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10237-799-16	473.2 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/15/2023	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	01/15/2023	

**Labeler -** Church & Dwight Co., Inc. (001211952)

Establishment			
Name	Address	ID/FEI	Business Operations
Church & Dwight Canada Corp.		253933600	manufacture(10237-799)