# MUCUS RELIEF DM MAXIMUM STRENGTH- dextromethorphan hbr, guaifenesin liquid Hy-Vee

-----

## **Drug Facts**

## Active ingredients (in each 20 mL)

Dextromethorphan HBr 20 mg Guaifenesin 400 mg

## **Purposes**

Cough suppressant

Expectorant

### Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of botherosme mucus and make coughs more productive
- temporarily relieves:
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - · the impulse to cough to help you get to sleep

# Warnings

### Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

# Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm(mucus)

When using this product, do not use more than directed

# Stop use and ask a doctor if

cough lasts more than 7 days, comes back, or occurs with fever, rash or headache that

lasts. These could be signs of a serious condition.

## If pregnant or breast-feeding,

ask a health professional before use.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### **Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours
- children under 12 years of age: do not use

### Other information

- each 20 mL contains: sodium 20 mg
- store between 20-25°C (68-77°F). Do not refrigerate

# **Inactive ingredients**

anhydrous citric acid, disodium EDTA, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

# **Principal Display Panel**

\*COMPARE TO the Active Ingredients in Maximum Strength Mucinex® Fast-Max® DM Max

Maximum Strength

Mucus Relief

Dextromethorphan HBr-Cough Suppressant

Guaifenesin-Expectorant

### DM Max

- Controls Cough
- Relieves Chest Congestion
- Thins & Loosens Mucus
- 4-Hour Dosing

• Ages 12 +

FL OZ (mL)

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® DM Max.

# TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: HY-VEE, INC. WEST DES MOINES, IA 50266 CALL 1-800-289-8343 www.hy-vee.com

# Package Label



\*COMPARE TO

Hyvee. Thealth

NDC 42507-505-06

# Maximum Strength Mucus Relief

Dextromethorphan HBr - Cough Suppressant Guaifenesin - Expectorant

### **DM Max**

- Controls Cough
- Relieves Chest Congestion
- Thins & Loosens Mucus
- 4-Hour Dosing
- Ages 12+

DISTRIBUTED BY HY-VEE, INC. WEST DES MOINES, IA 50266 SATISFACTION GUARANTEED CALL 1-800-289-8343 www.hy-vee.com

6 FL OZ (177 mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® DM Max.



PLD-D409A LB004108

## **Drug Facts**

## Active ingredients (in each 20 mL)

Dextromethorphan HBr 20 mg.....

......Cough suppressant Guaifenesin 400 mg......Expectorant

PEEL CORNER FOR MORE DRUG FACTS

Purposes

### Drug Facts (continued)

#### Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves
- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the intensity of coughing
- the impulse to cough to help you get to sleep

### Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Drug Facts (continued)

### Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product, do not use more than directed.

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

## Drug Facts (continued)

### Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours
- children under 12 years of age: do not use

### Other information

- each 20 mL contains: sodium 20 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

## Drug Facts (continued)

### Inactive ingredients

citric acid, disodium EDTA, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

# **MUCUS RELIEF DM MAXIMUM STRENGTH**

dextromethorphan hbr, guaifenesin liquid

<b>Product Information</b>	Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:42507-505	
Route of Administration	ORAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL	

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
EDETATE CALCIUM DISODIUM (UNII: 25IH6R4SGF)		
PROPYL GALLATE (UNII: 8D4SNN7V92)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SODIUM CITRATE (UNII: 1Q73Q2JULR)		
SORBITOL (UNII: 506T60A25R)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
GLYCERIN (UNII: PDC6A3C0OX)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SUCRALOSE (UNII: 96K6UQ3ZD4)		
XANTHAN GUM (UNII: TTV12P4NEE)		

l	Pa	Packaging			
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
		NDC:42507- 505-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/31/2016	06/30/2024

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/31/2016	06/30/2024

Revised: 4/2024 Hy-Vee