

**BENZETHONIUM CHLORIDE AND DYCLONINE HYDROCHLORIDE- liquid
bandage liquid
Rite Aid Corporation**

Liquid Bandage 1 fl oz (30 mL)

Active ingredient

Benzethonium chloride 0.2%

Dyclonine hydrochloride 0.75%

Purpose

First aid antiseptic

Topical analgesic

Uses

- first aid to prevent bacterial contamination in minor cuts, scrapes and burns
- for temporary relief of pain

Warnings

For external use only.

Flammable:

Keep away from fire or flame.

Do not use

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- deep or puncture wounds
- deep cuts
- animal bites
- serious burns

When using this product

- avoid contact with eyes

Stop use and ask a doctor if

- condition persists or gets worse
- symptoms persists for more than 7 days or clear up and occur again within a few

days

- infection occurs

Keep out of reach of children.

If swallowed, get medical help or call a Poison Control Center right away.

Directions

- for adults and children 2 years of age or older
- clean affected area
- apply a small amount on the area 1 to 3 times daily and let dry
- a second coating may be applied for extra protection
- children under 2 years of age: consult a doctor

Other information

- cap bottle tightly
- store between 20° to 25°C (68° to 77°F)
- may temporarily sting upon application
- to remove, apply more Liquid Bandage and quickly wipe off

Inactive ingredients

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

Questions?

Call 1-866-964-0939

Principal Display Panel

Rite Aid

LIQUID BANDAGE

TOPICAL ANALGESIC & ANTISEPTIC

For small cuts & wounds

To help prevent infection

Antiseptic formula dries to form an invisible protective cover that is flexible

NET 1.0 FL OZ (29.6 mL)

Use Liquid Bandage for blisters, chapped & cracked fingertips, hangnails, paper cuts, shaving nicks, & help in preventing the formulation of calluses

Great for: athletes, active lifestyles, outdoor sportsmen & musicians



LIQUID BANDAGE

TOPICAL ANALGESIC & ANTISEPTIC

For small cuts & wounds

To help prevent infection

Antiseptic formula dries to form an invisible protective cover that is flexible

NET 1.0 FL OZ (29.6 mL)

LIQUID BANDAGE

TOPICAL ANALGESIC & ANTISEPTIC

Use Liquid Bandage for: blisters, chapped & cracked fingertips, hangnails, paper cuts, shaving nicks & help in preventing the formation of calluses

Great for: athletes, active lifestyles, outdoor sportsmen & musicians

53-033RA-B01

LIQUID BANDAGE

TOPICAL ANALGESIC & ANTISEPTIC

For small cuts & wounds
To help prevent infection

LIQUID BANDAGE

TOPICAL ANALGESIC & ANTISEPTIC

For small cuts & wounds
To help prevent infection

NET 1.0 FL OZ (29.6 mL)



DISTRIBUTED BY: RITE AID
30 HUNTER LANE
CAMP HILL, PA 17011
www.riteaid.com
MADE IN USA WITH U.S.
AND IMPORTED PARTS

SATISFACTION
GUARANTEE

If you're not satisfied, we'll
happily refund your money.



Care ■ Wipe top of bottle clean after each use.

Questions? Call 1-866-964-0939

Inactive ingredients: acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40


Other information ■ cap bottle tightly ■ store between 20° to 25°C (68° to 77°F) ■ may temporarily sting upon application ■ to remove, apply more Liquid Bandage and quickly wipe off

Directions ■ for adults and children 2 years of age and older ■ clean affected area ■ apply a small amount on the area 1 to 3 times daily and let dry ■ a second coating may be applied for extra protection ■ children under 2 years of age: consult a doctor

Keep out of reach of children. If swallowed, get medical help or call a Poison Control Center right away.

■ symptoms persist for more than 7 days or clear up and occur again within a few days ■ infection occurs

■ common persists or gets worse



DO NOT USE IF NECKBAND IS BROKEN

Active ingredients: Benzethonium Chloride 0.2% and Dyclonine Hydrochloride 0.75%. **Purpose:** First aid antiseptic and topical analgesic. **Uses:** First aid to prevent bacterial contamination in minor cuts, scrapes, and burns. For the temporary relief of pain. **Warnings:** For external use only. **Flammable:** Keep away from fire or flame. **Do not use** in eyes, over large areas of the body, and longer than 1 week unless directed by a doctor.

LIQUID BANDAGE

TOPICAL ANALGESIC & ANTISEPTIC

For small cuts & wounds
To help prevent infection

NET 1.0 FL OZ (29.6 mL)

Ask a doctor before use if you have deep or puncture wounds, deep cuts, animal bites and serious burns.

When using this product avoid contact with eyes.

Stop use and ask a doctor if condition persists or gets worse, symptoms persist for more than 7 days or clear up and occur again within a few days, and infection occurs.

Keep out of reach of children. If swallowed, get medical help or call a Poison Control Center right away.

Directions: For adults and children 2 years of age and older. Clean affected area. Apply a small amount on the area 1 to 3 times daily and let dry. A second coating may be applied for extra protection. Children under 2 years of age: consult a doctor.

Other information: Cap bottle tightly. Store between 20° to 25°C (68° to 77°F). May temporarily sting upon application. To remove, apply more Liquid Bandage and quickly wipe off.

Inactive ingredients: acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

Questions? Call 1-866-964-0939

DISTRIBUTED BY: RITE AID, 30 HUNTER LANE
CAMP HILL, PA 17011 www.riteaid.com

MADE IN USA WITH U.S. AND IMPORTED PARTS

SATISFACTION GUARANTEE: if you're not satisfied, we'll happily refund your money. 50-033RA-B01

BENZETHONIUM CHLORIDE AND DYCLONINE HYDROCHLORIDE

liquid bandage liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-0779
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	0.2 mg in 30 mL
DYCLONINE HYDROCHLORIDE (UNII: ZEC193879Q) (DYCLONINE -	DYCLONINE	0.75 mg

UNII:078A24Q300)

HYDROCHLORIDE

in 30 mL

Inactive Ingredients

Ingredient Name	Strength
PYROXYLIN (UNII: KYR8BR2X6O)	
ETHYL ACETATE (UNII: 76845O8NMZ)	
ACETONE (UNII: 1364PS73AF)	
AMYL ACETATE (UNII: 92Q24NH7AS)	
CASTOR OIL (UNII: D5340Y2I9G)	
ALCOHOL (UNII: 3K9958V90M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0779-3	30 mL in 1 BOTTLE, WTH APPLICATOR; Type 0: Not a Combination Product	02/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	02/01/2020	

Labeler - Rite Aid Corporation (014578892)

Revised: 12/2024

Rite Aid Corporation