

ASPIRIN PAIN RELIEF- aspirin tablet, delayed release
Rite Aid Corporation

Rite Aid 44-227

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - headache
 - menstrual pain
 - a cold
 - toothache
 - muscle pain
- or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- hives
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- new symptoms occur
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

black iron oxide, cellulose, colloidal anhydrous silica, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, propylene glycol, shellac wax, simethicone, sodium alginate, sodium bicarbonate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal Display Panel

NDC 11822-2270-4

**Compare to the active ingredient
in **Ecotrin®** Regular Strength

**Talk to your doctor or other healthcare provider
before using this product for your heart.**

REGULAR STRENGTH
PAIN RELIEF
ASPIRIN

**ASPIRIN 325 mg
PAIN RELIEVER (NSAID)**

aspirin regimen
safety coated

500 ENTERIC
COATED TABLETS

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**This product is not manufactured or distributed by Medtech Products Inc.,
owner of the registered trademark Ecotrin® Regular Strength. 50844
REV0122C22714

**DISTRIBUTED BY: RITE AID
30 HUNTER LANE
CAMP HILL, PA 17011
www.riteaid.com**

SATISFACTION GUARANTEE

If you're not satisfied, we'll happily refund your money.

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Ecotrin® Regular Strength. 50844 REV0122C22714

DISTRIBUTED BY: RITE AID
30 HUNTER LANE,
CAMP HILL, PA 17011
www.riteaid.com

SATISFACTION GUARANTEE
If you're not satisfied, we'll happily refund your money.

0 11822 42300 7

PEEL HERE FOR MORE DRUG FACTS

NDC 11822-2270-4
Talk to your doctor or other healthcare provider before using this product for your heart.

**Compare to the active ingredient in Ecotrin® Regular Strength

REGULAR STRENGTH PAIN RELIEF

ASPIRIN

ASPIRIN 325 mg

PAIN RELIEVER (NSAID)

aspirin regimen
safety coated

500 ENTERIC COATED TABLETS

No Print/No Varnish Area
Lot no. & Exp. Date

Drug Facts

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Active ingredient (in each tablet)	Purpose
Aspirin 325 mg (NSAID)	Pain reliever
*nonsteroidal anti-inflammatory drug	

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis ■ headache ■ menstrual pain ■ a cold
- toothache ■ muscle pain
- or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling ■ hives ■ shock
- asthma (wheezing)

STOP PEELING

Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout ■ diabetes ■ arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood ■ have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child

Drug Facts (continued)

or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients black iron oxide, cellulose, colloidal anhydrous silica, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, propylene glycol, shellac wax, simethicone, sodium alginate, sodium bicarbonate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?
1-800-426-9391

Rite Aid 44-227

ASPIRIN PAIN RELIEF

aspirin tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-2270
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	

POWDERED CELLULOSE (UNII: SMD1X3XO9M)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
STARCH, CORN (UNII: O8232NY3SJ)
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)
HYMPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)
POLYDEXTROSE (UNII: VH2XOU12IE)
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)
Polyvinyl Acetate Phthalate (UNII: 58QVG85GW3)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
SHELLAC (UNII: 46N107B71O)
DIMETHICONE (UNII: 92RU3N3Y1O)
WATER (UNII: 059QF0KO0R)
SODIUM ALGINATE (UNII: C269C4G2ZQ)
SODIUM BICARBONATE (UNII: 8MDF5V39QO)
STEARIC ACID (UNII: 4ELV7Z65AP)
TALC (UNII: 7SEV7J4R1U)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
TRIACETIN (UNII: XHX3C3X673)
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	44;227
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-2270-4	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/14/1992	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/14/1992	

Labeler - Rite Aid Corporation (014578892)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	manufacture(11822-2270) , pack(11822-2270)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11822-2270)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(11822-2270)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(11822-2270)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(11822-2270)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(11822-2270)

Revised: 12/2025

Rite Aid Corporation