

**ACTICON- dexbrompheniramine maleate, pseudoephedrine hydrochloride solution**  
**ACTIPHARMA, LLC**

-----  
**NEW ACTICON® Cold & Allergy Oral Solution**

**Drug Facts**

**Active Ingredients**  
**(in each 5 mL tsp)**

Dexbrompheniramine Maleate, USP 2 mg  
Pseudoephedrine HCl, USP 60 mg

**Purpose**

Antihistamine  
Nasal Decongestant

**Uses**

- Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:
- relieves sinus congestion and pressure, helps decongest sinus openings and passages
- restores freer breathing through the nose
- runny nose • sneezing • itching of the nose or throat • itchy, watery eyes • nasal congestion

**Warnings**

**Do not exceed recommended dosage**

**Do not use this product**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis • glaucoma • heart disease • high blood pressure • thyroid disease • diabetes • difficulty in urination due to enlargement of the prostate gland

**Do not take this product if you are taking sedatives or tranquilizers, without first consulting your doctor.**

**When using this product**

- excitability may occur, especially in children
- may cause drowsiness
- alcohol, sedatives and tranquilizers may increase drowsiness effect
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- if symptoms do not improve within 7 days or are accompanied by fever
- new symptoms occur

**If pregnant or breast feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

**Directions**

**Do not exceed more than 4 doses in 24 hours, or as directed by a doctor.**

<b>AGE</b>	<b>DOSE</b>
Adults and children 12 years of age and over	1 teaspoonful (5 ml) every 4-6 hours

**Other information**

- Tamper Evident Feature: Do not use if inner seal is torn, cut, or opened.
- Store at controlled room temperature 15°- 30°C (59°- 86°F).

**Inactive ingredients**

Citric acid, flavor, methylparaben, potassium citrate, propylene glycol, propylparaben, purified water, sorbitol, sucralose.

**Questions or Comments?**

Call weekdays from 8 AM to 4 PM AST at **1.787.608.0882.**

**Contains the same active ingredients as Conex®\***

**Cherry Flavor**

**ActiPharma**

COMMITTED TO HEALTH AND WELL-BEING

WWW.ACTIPHARMA.NET

Manufactured in the USA with imported ingredients for ActiPharma, LLC San Juan, PR 00917. \* Conex® is a registered trademark of Llorens Pharmaceutical Corp. This product is not manufactured, distributed or marketed by Llorens Pharmaceutical Corp.

**Packaging**

NDC 63102-108-16

**NEW**  
**ACTICON®**

**Cold & Allergy**  
**Oral Solution**

Contains the same active ingredients as Conex®\*

**ANTIHISTAMINE**  
**NASAL DECONGESTANT**

Each 5 mL contains:  
Dexbrompheniramine Maleate...2 mg  
Pseudoephedrine HCl.....60 mg

**Cherry Flavor**



16 Fl.oz. (474 mL)

Lot #  
Exp. Date

**Drug Facts**

**Active Ingredients**  
(in each 5 mL tsp)  
Dexbrompheniramine Maleate, USP 2 mg ..... Antihistamine  
Pseudoephedrine HCl, USP 60 mg ..... Nasal Decongestant

**Purpose**

**Uses**

- Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:
- relieves sinus congestion and pressure, helps decongest sinus openings and passages
- restores freer breathing through the nose
- runny nose • sneezing • itching of the nose or throat • itchy, watery eyes • nasal congestion

**Warnings**

**Do not exceed recommended dosage**

**Do not use this product**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma • heart disease • high blood pressure • thyroid disease
- diabetes • difficulty in urination due to enlargement of the prostate gland

**Do not take this product if you are taking sedatives or tranquilizers, without first consulting your doctor.**

**When using this product**

- excitability may occur, especially in children • may cause drowsiness • alcohol, sedatives and tranquilizers may increase drowsiness effect • avoid alcoholic beverages • use caution when driving a motor vehicle or operating machinery

**Drug Facts (continued)**

**Warnings (continued)**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur • if symptoms do not improve within 7 days or are accompanied by fever • new symptoms occur

**If pregnant or breast feeding, ask a health professional before use. Keep out of reach of children.** In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

**Directions**

**Do not exceed more than 4 doses in 24 hours, or as directed by a doctor.**

AGE	DOSE
Adults and children 12 years of age and over	1 teaspoonful (5 ml) every 4-6 hours

**Other information**

- Tamper Evident Feature: Do not use if inner seal is torn, cut, or opened.
- Store at controlled room temperature 15°- 30°C (59°- 86°F).

**Inactive ingredients** Citric acid, flavor, methylparaben, potassium citrate, propylene glycol, propylparaben, purified water, sorbitol, sucralose.

**Questions or Comments?** Call weekdays from 8 AM to 4 PM AST at 1.787.608.0882.

Manufactured in the USA with imported ingredients for ActiPharma, LLC San Juan, PR 00917. \* Conex® is a registered trademark of Llorens Pharmaceutical Corp. This product is not manufactured, distributed or marketed by Llorens Pharmaceutical Corp.

Rev. 11/24



**ACTICON**

dexbrompheniramine maleate, pseudoephedrine hydrochloride solution

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63102-108
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>DEXBROMPHENIRAMINE MALEATE</b> (UNII: BPA9UT29BS) (DEXBROMPHENIRAMINE - UNII: 75T64B71RP)	DEXBROMPHENIRAMINE MALEATE	2 mg in 5 mL
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b> (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII: 7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	60 mg in 5 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>POTASSIUM CITRATE</b> (UNII: EE90ONI6FF)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

## Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63102-108-16	474 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/15/2022	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/15/2022	

**Labeler** - ACTIPHARMA, LLC (079340948)

Revised: 7/2025

ACTIPHARMA, LLC