

**TARGET MULTI SYMPTOM FLU RELIEF- acetaminophen, dextromethorphan hbr, and acetaminophen, chlorpheniramine maleate, dextromethorphan hbr
TARGET CORPORATION**

Up & up Multi-Symptom Flu Relief Combo Pack *Drug Facts*

Flu Relief Max Strength Daytime Powder**

Active ingredients (in each packet)

Acetaminophen 1000 mg

Dextromethorphan HBr 30 mg

Purposes

Pain reliever/Fever reducer

Cough suppressant

Uses

- temporarily relieves these symptoms due to a common cold or flu:
 - headache
 - minor aches and pains
 - cough due to minor throat and bronchial irritation
 - minor sore throat pain
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

When using this product

- **do not exceed recommended dosage.**

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- take every 6 hours, while symptoms persist. Do not take more than 3 packets in 24 hours unless directed by a doctor.

1. Age	1. Dose
1. adults and children 12 years of age and over	1. one packet

1. children under 12 years of age	1. do not use
-----------------------------------	---------------

- dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume the entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other information

- **each packet contains:** potassium 6 mg
- store at room temperature. Protect from excessive heat and moisture.

Inactive ingredients

anhydrous citric acid, caramel, flavor, maltodextrin, potassium chloride, silica, sucralose, sucrose.

Questions or comments?

1-866-467-2748

Flu Relief Max Strength Nighttime Powder**

Drug Facts

Active ingredients (in each packet)

Acetaminophen 1000 mg

Chlorpheniramine maleate 4 mg

Dextromethorphan HBr 30 mg

Purposes

Pain reliever/Fever reducer

Antihistamine

Cough suppressant

Uses

- temporarily relieves these symptoms due to a common cold or flu:
 - headache
 - minor aches and pains

- cough due to minor throat and bronchial irritation
- minor sore throat pain
- runny nose
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAO, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- avoid alcoholic drinks

- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- take every 6 hours, while symptoms persist. Do not take more than 3 packets in 24 hours unless directed by a doctor.

1. Age	1. Dose
1. adults and children 12 years of age and over	1. one packet
1. children under 12 years of age	1. do not use

- dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat

Other information

- **each packet contains:** potassium 6 mg
- store at room temperature. Protect from excessive heat and moisture.

Inactive ingredients

anhydrous citric acid, caramel, flavor, maltodextrin, potassium chloride, silica, sucralose, sucrose.

Questions or comments?

1-866-467-2748

Other Safety Information

READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE. KEEP CARTON FOR REFERENCE. DO NOT DISCARD.

DO NOT TAKE BOTH PRODUCTS AT THE SAME TIME.

DO NOT TAKE MORE THAN 3 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.

DO NOT TAKE A DOSE OF THE NIGHTTIME PRODUCT SOONER THAN 4 HOURS AFTER THE LAST DOSE OF THE DAYTIME PRODUCT UNLESS DIRECTED BY YOUR DOCTOR.

TAMPER-EVIDENT INNER UNIT. DO NOT USE IF NECKBAND PRINTED WITH "SEALED FOR SAFETY" IS TORN OR MISSING.

Distributed by:

Principal Display Panel

Compare to the active ingredients in Theraflu Multi-Symptom Flu Relief Max Strength** Daytime* & Nighttime*.

NDC 82442-559-12

MULTI SYMPTOM FLU RELIEF

Daytime Maximum Strength** Flu Relief

AcetaminophenPain Reliever/Fever Reducer

Dextromethorphan HBrCough Suppressant

6 PACKETS

Nighttime Maximum Strength** Flu Relief

AcetaminophenPain Reliever/Fever Reducer

Chlorpheniramine MaleateAntihistamine

Dextromethorphan HBrCough Suppressant

6 PACKETS

Hot liquid therapy that relieves:

- Fever
- Body ache
- Headache
- Sore throat pain

Daytime
Maximum Strength**
Flu Relief



Nighttime
Maximum Strength**
Flu Relief



MULTI-SYMPTOM FLU RELIEF

Daytime
Maximum Strength**
Flu Relief



Nighttime
Maximum Strength**
Flu Relief



MULTI-SYMPTOM FLU RELIEF

Daytime
Maximum Strength**
Flu Relief



Nighttime
Maximum Strength**
Flu Relief



MULTI-SYMPTOM FLU RELIEF

Daytime
Maximum Strength**
Flu Relief



Nighttime
Maximum Strength**
Flu Relief



MULTI-SYMPTOM FLU RELIEF

Daytime
Maximum Strength**
Flu Relief



Nighttime
Maximum Strength**
Flu Relief



MULTI-SYMPTOM FLU RELIEF

Daytime
Maximum Strength**
Flu Relief



Nighttime
Maximum Strength**
Flu Relief



MULTI-SYMPTOM FLU RELIEF

TARGET MULTI SYMPTOM FLU RELIEF

acetaminophen, dextromethorphan hbr, and acetaminophen, chlorpheniramine maleate, dextromethorphan hbr kit

Product Information				
Product Type		HUMAN OTC DRUG	Item Code (Source)	NDC:82442-559
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82442-559-12	1 in 1 CARTON; Type 1: Convenience Kit of Co-Package	06/06/2025	

Quantity of Parts		
Part #	Package Quantity	Total Product Quantity
Part 1	1 CARTON	6
Part 2	1 CARTON	6

Part 1 of 2

TARGET DAYTIME MS FLU

acetaminophen, dextromethorphan hbr powder

Product Information	
Item Code (Source)	NDC:82442-558
Route of Administration	ORAL

Active Ingredient/Active Moiety		
Ingredient Name		Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		1000 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)		30 mg

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARAMEL (UNII: T9D99G2B1R)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
POTASSIUM CHLORIDE (UNII: 660YQ98I10)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

SUCROSE (UNII: C151H8M554)				
Product Characteristics				
Color	white (to off white, yellow, beige, and brown color)	Score		
Shape		Size		
Flavor	HONEY (Lemon)	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82442-558-06	6 in 1 CARTON; Type 1: Convenience Kit of Co-Package		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M012	06/06/2025	
Part 2 of 2				
TARGET NIGHTTIME MS FLU RELIEF				
acetaminophen, chlorpheniramine maleate, dextromethorphan hbr powder				

Product Information	
Item Code (Source)	NDC:82442-560
Route of Administration	ORAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	1000 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	4 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARAMEL (UNII: T9D99G2B1R)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	

POTASSIUM CHLORIDE (UNII: 660YQ98I10)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
SUCROSE (UNII: C151H8M554)				
Product Characteristics				
Color	white (white to off-white, yellow, beige, and brown color)	Score		
Shape		Size		
Flavor	HONEY (lemon)	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82442-560-06	6 in 1 CARTON; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012		06/06/2025	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012		06/06/2025	

Labeler - TARGET CORPORATION (006961700)