

TOPCO MICONAZOLE 3- miconazole nitrate cream
Topco Associates LLC

Topco Associates LLC. Miconazole 3 Drug Facts

Active ingredient

Miconazole nitrate 4% (200 mg in each applicator)

Purpose

Vaginal antifungal

Uses

- treats vaginal yeast infections

Warnings

For vaginal use only

Do not use

if you have never had a vaginal yeast infection diagnosed by a doctor

Ask a doctor before use if you have

- **vaginal itching and discomfort for the first time**
- **lower abdominal, back or shoulder pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge. You may have a more serious condition.**
- vaginal yeast infections often (such as once a month or 3 in 6 months). You could be pregnant or have a serious underlying medical cause for your symptoms, including diabetes or a weakened immune system.
- been exposed to the human immunodeficiency virus (HIV) that causes AIDS

Ask a doctor or pharmacist before use if you are

taking the prescription blood thinning medicine warfarin, because bleeding or bruising may occur

When using this product

- do not use tampons, douches, spermicides or other vaginal products. Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs).
- do not have vaginal intercourse
- mild increase in vaginal burning, itching or irritation may occur
- if you do not get complete relief ask a doctor before using another product

Stop use and ask a doctor if

- symptoms do not get better in 3 days
- symptoms last more than 7 days
- you get a rash or hives, abdominal pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- before using this product read the enclosed consumer information leaflet for complete directions and information
- adults and children 12 years of age and over:
- insert 1 applicatorful into the vagina at bedtime for 3 nights in a row. Throw applicator away after use.
- **children under 12 years of age: ask a doctor**

Other information

- do not use if **printed** applicator wrapper is torn, open, or incompletely sealed
- do not purchase if carton is open
- store at 20°-25°C (68°-77°F)

Inactive ingredients

benzoic acid, cetyl alcohol, isopropyl myristate, polysorbate 60, potassium hydroxide, propylene glycol, purified water, stearyl alcohol

Questions or comments?

1-888-423-0139

Consumer Information

3-Day Treatment

CONSUMER INFORMATION LEAFLET

Miconazole Nitrate Vaginal Cream (4%)

VAGINAL ANTIFUNGAL

Cures Most Vaginal Yeast Infections

Why should I use this product?

This product contains 4% miconazole nitrate cream in prefilled applicators that cures most vaginal yeast infections. ***Do not use this product if this is the first time you have vaginal discharge, itching, burning and***

discomfort. See your doctor or health professional first to find out the cause of your symptoms. If a doctor has told you in the past that you had a vaginal yeast infection and you have the same symptoms now (such as vaginal discharge, itching or burning), then this product may work for you.

What is a vaginal yeast infection?

A vaginal yeast infection is a common condition caused by an overgrowth of yeast (Candida) that may normally live in the vagina. Your doctor may call this infection "monilia" or "candidiasis." Some women may have a yeast infection on the skin outside of the vagina (vulva) at the same time that they have a vaginal infection.

Who can get a vaginal yeast infection?

You can get a vaginal yeast infection at any age. It is most common during the childbearing years. Women who are pregnant or diabetic, taking antibiotics, birth control pills or steroids, or who have a weakened immune system are more likely to get repeated yeast infections that may not clear up easily with proper treatment.

Some medical conditions can weaken the body's normal ability to fight infection. One of the most serious of these conditions is infection with the human immunodeficiency virus (HIV - the virus that causes AIDS). The HIV virus causes the body to be more likely to get infections, including vaginal yeast infections that may not clear up easily with proper treatment. If you may have been exposed to HIV and get repeated vaginal yeast infections, you should see your doctor right away. For more information on HIV infection, please contact your doctor or the CDC National AIDS HOTLINE. The CDC phone numbers are: 1-800-232-4636 (English/Spanish), or 1-888-232-6348 (hearing impaired, TTY).

How can I tell if I have a vaginal yeast infection?

When you have a vaginal yeast infection, you may have one or more of the following symptoms:

- vaginal itching
- vaginal discharge that may be thick, white, and lumpy like cottage cheese
- vaginal soreness, irritation, or burning
- rash or redness on the skin outside the vagina (vulva)
- burning on urination
- painful vaginal intercourse (sex)

Note: Vaginal yeast infections do NOT cause fever, chills, lower abdominal, back or shoulder pain, foul-smelling vaginal discharge, or a missed period. These may be signs of a sexually transmitted disease (STD) or a tubal pregnancy. If you have these symptoms, call your doctor right away.

What are other causes of a vaginal discharge?

It is normal to have a small amount of vaginal discharge at certain times of the month. This normal discharge may be clear or slightly white and does not cause itching, pain or a foul odor.

The most common cause of an abnormal vaginal discharge is an infection. These infections include bacterial vaginosis (BV), trichomoniasis (Trich), gonorrhea (GC) and/or chlamydia. All of these may be transmitted sexually and are called sexually transmitted diseases (STDs). If you have more questions about sexually transmitted diseases (STDs) call the CDC Hotline at 1-800-232-4636.

Although many of the infections mentioned above can cause symptoms similar to a vaginal yeast infection (vaginal discharge, irritation and itching), their diagnosis must be made by a doctor so that proper treatment can be given.

If these infections are not properly treated or if proper treatment is delayed, serious problems, such as pelvic inflammatory disease (PID) may result, which may prevent you from having children in the future. If you are pregnant and do not get the proper treatment, the infection may be passed to your baby before

or during delivery and may cause your baby to have permanent damage. If you have multiple sex partners or a new sex partner, you should also ask a doctor before use to make sure you do not have an STD.

Why do women get repeated vaginal yeast infections?

Women may get repeated vaginal yeast infections that may not clear up easily with proper treatment. Listed below are some of the causes of repeated yeast infections:

- **hormonal changes** occurring a few days before the monthly period
- **use of antibiotics**
- **use of some birth control pills**
- **pregnancy**
- **diabetes ("sugar" or "high blood sugar")**
- **clothing** - wearing tight layers or moist clothing in the genital area
- **weakened immune system** - some drugs (such as chemotherapy or steroids) and medical conditions can weaken the body's normal ability to fight infection. One of the most serious of these conditions is infection with the human immunodeficiency virus (HIV - the virus that causes AIDS). Infection with HIV causes the person to be more likely to get infections, including vaginal yeast infections.

If you get vaginal yeast infections often (such as once a month or 3 in 6 months), you should talk to a doctor.

Are vaginal yeast infections sexually transmitted?

Vaginal yeast infections are usually not spread by having intercourse (sex). However, if your partner has a rash, itching or discomfort in his genital area, he should contact a doctor to find out the cause of his symptoms and tell the doctor that you are treating your vaginal yeast infection with this product.

How can I prevent repeated vaginal yeast infections?

To lower your chances of getting another yeast infection:

- **Try to keep the genital area cool and dry.** Yeast grow well in warm, moist areas. The following suggestions may be helpful:
 - (1) Wear cotton underwear and loose-fitting clothes.
 - (2) Change out of damp clothes or a wet bathing suit as soon as possible.
 - (3) If you use minipads when you are not having a menstrual period, change the minipads often.
- **Talk with your doctor about any drugs you are now taking.** You are more likely to get a vaginal yeast infection if you are taking certain drugs such as antibiotics, steroids, or birth control pills. Do not stop taking these drugs without first asking your doctor. A doctor may need to see you to make sure that you do not have other medical conditions such as diabetes or a weakened immune system.

Can I use this product during my menstrual period?

Yes, this product can be used during your menstrual period. In fact, many women get vaginal yeast infections just before their period because of hormonal changes. Using this product during your period will not affect how well this product works. If you have started treatment and your period occurs, you should complete the full course of treatment.

Do not use tampons while using this product, because tampons may remove some of the drug from the vagina. Use deodorant-free sanitary napkins or pads instead, and change them often.

Can I use other vaginal products with this product?

This drug should **not** be used with other vaginal products.

- Douches and tampons may remove some of the cream from the vagina.
- Spermicides may interfere with this product.
- Condoms and diaphragms may be damaged by this product and fail to prevent pregnancy or sexually transmitted diseases (STDs).

How can I get the best results when treating my infection?

- Use 1 prefilled applicator of vaginal cream at bedtime, even during your menstrual period for 3 nights in a row.
- Dry the genital area thoroughly after a shower, bath or swim. Change out of a wet bathing suit or damp clothes as soon as possible. A dry area is less likely to lead to the overgrowth of yeast.
- Wear cotton underwear and loose-fitting clothes.
- Wipe from front to back after a bowel movement or after urination.
- Do not douche, because douching may wash the drug out of the vagina.
- Do not use tampons, because they remove some of the drug from the vagina. Use deodorant-free sanitary napkins or pads as needed.
- Do not use spermicides, as they may interfere with this product.
- Do not have vaginal intercourse while using this product.
- Do not scratch the skin outside the vagina. Scratching can cause more irritation and can spread the infection.
- **Tell your doctor about any drugs you are now taking.** Certain drugs such as antibiotics, steroids, and birth control pills, may make it more likely for you to get a vaginal yeast infection. If you are taking any of these drugs do not stop taking them without first asking a doctor.
- If you have any other medical questions or concerns about vaginal yeast infections, call your doctor.

What warnings should I know about when using this product?

For vaginal use only.

Do not use if you have never had a vaginal yeast infection diagnosed by a doctor.

Ask a doctor before use if you have:

- **vaginal itching and discomfort for the first time. You may need a different treatment.**
- **lower abdominal, back or shoulder pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge. You could have a more serious condition.**
- vaginal yeast infections often (such as once a month or 3 in 6 months). You could be pregnant or have a serious underlying medical cause for your symptoms, including diabetes or a weakened immune system.
- been exposed to the human immunodeficiency virus (HIV) that causes AIDS.

Ask a doctor or pharmacist before use if you are taking the prescription blood thinning medicine, warfarin (coumadin), because bleeding or bruising may occur.

When using this product:

- do not use tampons, douches, spermicides, or other vaginal products. Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs).

- do not have vaginal intercourse
- mild increase in vaginal burning, itching or irritation may occur
- if you do not get complete relief ask a doctor before using another product


How Should I Use This Product?

This product is for adults and children 12 years of age and over. For children under 12 years, ask a doctor.


Directions for using the Applicator containing the Vaginal Cream:

Begin treatment before going to bed:

- 1 Remove the prefilled applicator from the wrapper.
- 2 Unscrew and remove violet cap from prefilled applicator.
- 3 Insert plunger into prefilled applicator by placing small end of plunger into the ribbed end of the applicator.



- 4 Gently insert the applicator into the vagina as far as it will go comfortably. This can be done while lying on your back with your knees bent (as shown in the picture) or while standing with your feet apart and your knees bent.



- 5 With one hand holding the barrel, use the other hand to push the plunger all the way in to place the cream as far back in the vagina as possible. Then remove both parts of the applicator from the vagina.
- 6 Throw the applicator away after use. Do not flush in toilet.
- 7 Lie down as soon as possible after inserting the cream. This will reduce leakage.
- 8 Repeat steps 1 through 7 before going to bed for the next 2 days.

You may want to use deodorant-free pads or pantyshields to protect your clothing during the time that you are using this product. This is because the cream can leak or you may get some discharge. **Do not use tampons, douches, spermicides, condoms or diaphragms** until after you have completed treatment and your symptoms are gone.

Stop use and ask your doctor if:

- **symptoms do not get better in 3 days**
- **symptoms last more than 7 days**
- **you get a rash or hives, abdominal pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge.**

These may be signs that this product is not working, or you may have a more serious condition or an allergic reaction.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

What side effects may occur with this product?

A mild increase in vaginal burning, itching, or irritation may occur when the applicator containing the cream is inserted. Abdominal cramping has also been reported.

Stop using this product and consult your doctor if you have abdominal pain, hives, skin rash, or if you have severe vaginal burning, itching, or irritation or swelling.

What should I do if I have questions about this product?

If you have questions of a medical nature, please contact your pharmacist, doctor or health care professional. If you have any other questions or need more information on this product, call our toll-free number **1-800-719-9260**.

Other Information:

- **TAMPER-EVIDENT UNIT** – do not use if **printed** applicator wrapper is torn, open or incompletely sealed.
- store at 20°-25°C (68°-77°F)

Active Ingredient: miconazole nitrate 4% (200 mg per applicator)

Inactive Ingredients: benzoic acid, cetyl alcohol, isopropyl myristate, polysorbate 60, potassium hydroxide, propylene glycol, purified water, stearyl alcohol.

Distributed By

Perrigo®

Allegan, MI 49010

10/16

: 07000 00 J1

Package/Label Principal Display Panel

3-DAY TREATMENT

3 Ready-to-Use Prefilled Disposable Cream Applicators

COMPARE TO MONISTAT® 3 ACTIVE INGREDIENT

3-DAY TREATMENT

Miconazole 3

MICONAZOLE NITRATE VAGINAL CREAM (4%)

VAGINAL ANITFUNGAL

(Miconazole Nitrate 200 mg per Applicator)
CURES Most Vaginal YEAST INFECTIONS
NET WT. 0.18 OZ (5 g) EACH APPLICATOR



3-DAY TREATMENT

- 3 Ready-to-Use Prefilled Disposable Cream Applicators



Miconazole 3

MICONAZOLE NITRATE VAGINAL CREAM (4%)
VAGINAL ANTIFUNGAL
(Miconazole Nitrate 200 mg per Applicator)

TopCare
health™

3-DAY TREATMENT

Miconazole 3

MICONAZOLE NITRATE VAGINAL CREAM (4%)
VAGINAL ANTIFUNGAL

- CURES Most Vaginal YEAST INFECTIONS

NET WT. 0.18 OZ (5 g) EACH APPLICATOR

COMPARE TO MONISTAT® 3
ACTIVE INGREDIENT*

NDC 36800-782-14

3

07014 88 C1

CODE AREA

3-DAY TREATMENT

TopCare
health™

Miconazole 3

MICONAZOLE NITRATE VAGINAL CREAM (4%)
VAGINAL ANTIFUNGAL



3-DAY TREATMENT

Miconazole 3

MICONAZOLE NITRATE VAGINAL CREAM (4%)
VAGINAL ANTIFUNGAL

Drug Facts

Active ingredient
Miconazole nitrate 4% (200 mg in each applicator).....
Purpose
Vaginal antifungal

Uses
■ treats vaginal yeast infections

Warnings
For vaginal use only
Do not use if you have never had a vaginal yeast infection diagnosed by a doctor

Ask a doctor before use if you have
■ vaginal itching and discomfort for the first time
■ lower abdominal, back or shoulder pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge. You may have a more serious condition.
■ vaginal yeast infections often (such as once a month or 3 in 6 months). You could be pregnant or have a serious underlying medical cause for your symptoms, including diabetes or a weakened immune system.
■ been exposed to the human immunodeficiency virus (HIV) that causes AIDS

Ask a doctor or pharmacist before use if you are taking the prescription blood thinning medicine warfarin, because bleeding or bruising may occur

When using this product
■ do not use tampons, douches, spermicides or other vaginal products. Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs).
■ do not have vaginal intercourse
■ mild increase in vaginal burning, itching or irritation may occur
■ if you do not get complete relief ask a doctor before using another product

Drug Facts (continued)

Stop use and ask a doctor if
■ symptoms do not get better in 3 days
■ symptoms last more than 7 days
■ you get a rash or hives, abdominal pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ before using this product read the enclosed consumer information leaflet for complete directions and information
■ adults and children 12 years of age and over:
■ insert 1 applicatorful into the vagina at bedtime for 3 nights in a row. Throw applicator away after use.
■ children under 12 years of age: ask a doctor

Other information
■ do not use if printed applicator wrapper is torn, open, or incompletely sealed
■ do not purchase if carton is open
■ store at 20°-25°C (68°-77°F)

Inactive ingredients
benzoic acid, cetyl alcohol, isopropyl myristate, polysorbate 60, potassium hydroxide, propylene glycol, purified water, stearyl alcohol

Questions or comments? 1-888-423-0139



Scan here or call
1-888-423-0139 for
more information



This TopCare® product is laboratory tested to guarantee its highest quality. Your total satisfaction is guaranteed.
This product is not manufactured or distributed by INSIGHT Pharmaceuticals, LLC, distributor of Miconazole 3.

TAMPER-EVIDENT UNIT
DO NOT USE IF PRINTED APPLICATOR WRAPPER IS TORN, OPEN OR INCOMPLETELY SEALED.

DISTRIBUTED BY TOPCO ASSOCIATES LLC
ELK GROVE VILLAGE, IL 60007
© TOPCO PERA1119 QUESTIONS? 1-888-423-0139
topcare@topco.com www.topcarebrand.com

TOPCO MICONAZOLE 3

miconazole nitrate cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-782
Route of Administration	VAGINAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MICONAZOLE NITRATE (UNII: VW4HCYW1K) (MICONAZOLE - UNII:7NNO0D7S5M)	MICONAZOLE NITRATE	40 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
POLYSORBATE 60 (UNII: CAL22UVI4M)	

POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
STEARYL ALCOHOL (UNII: 2KR89I4HIY)	

Product Characteristics

Color	WHITE (to off white)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-782-14	3 in 1 CARTON	04/27/2020	
1		5 g in 1 APPLICATOR; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091366	04/27/2020	

Labeler - Topco Associates LLC (006935977)

Revised: 4/2020

Topco Associates LLC