

**LORATADINE- loratadine tablet**  
**Publix Super Markets Inc**

-----  
**Loratadine**

***Drug Facts***

**Active ingredient (in each tablet)**

Loratadine, USP 10 mg

**Purpose**

Antihistamine

**Uses**

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

**Warnings**

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.

**When using this product** do not take more than directed. Taking more than directed may cause drowsiness.

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

adults and children 6 years and over	1 tablet daily; not more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Other Information**

- store between 68° to 77° F (20° to 25° C)
- protect from excessive moisture
- **TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS ARE TORN, BROKEN OR**

**SHOW ANY SIGNS OF TAMPERING.**

**Inactive ingredients**

corn starch, lactose monohydrate, magnesium stearate, pregelatinized starch

**DISTRIBUTED BY PUBLIX SUPER MARKETS, INC.,  
3300 PUBLIX CORPORATE PARKWAY, LAKELAND, FL 33811**

**PRINCIPAL DISPLAY PANEL - 10 mg Tablet Blister Pack Carton**

NDC 56062-526-69

INDOOR & OUTDOOR ALLERGIES  
24-HOUR

allergyrelief

LORATADINE TABLETS, USP 10 mg

ANTIHISTAMINE

Non-drowsy\*

Relief of:

- Sneezing • Runny nose
- Itchy, watery eyes
- Itchy throat or nose


10 TABLETS

\*When taken as directed. See Drug Facts Panel.

ACTUAL SIZE

†Compare to the active  
ingredient of Claritin®

5199416



Keep the carton. It contains important information. See end panel for expiration date.

**Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.**

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

**Warnings**

■ itching of the nose or throat  
■ sneezing  
■ itchy, watery eyes  
■ runny nose  
or other upper respiratory allergies;  
temporarily relieves these symptoms due to hay fever

**Uses**

Loratadine, USP 10 mg, .....Antihistamine

**Active ingredient (in each tablet)**

**Purpose**

**Drug Facts (continued)**  
before use.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center. right away (1-800-222-1222).

INDOOR & OUTDOOR ALLERGIES  
24-HOUR



**allergyrelief**

LORATADINE TABLETS, USP 10 mg  
ANTIHISTAMINE



INDOOR & OUTDOOR ALLERGIES  
24-HOUR

**allergyrelief**

LORATADINE TABLETS, USP 10 mg  
ANTIHISTAMINE

NDC 56062-526-69



ACTUAL SIZE

<sup>†</sup>Compare to the active ingredient of Claritin®

**Non-drowsy\***

**Relief of:**

- Sneezing • Runny nose
- Itchy, watery eyes
- Itchy throat or nose

**10** TABLETS

\*When taken as directed. See Drug Facts Panel.

Expiration Date:

Batch No.

Non Varnish Area

<sup>†</sup>All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Claritin®.

**Inactive ingredients**  
corn starch, lactose monohydrate, magnesium stearate, pregelatinized starch

**Other information**  
■ store between 68° to 77° F (20° to 25° C)  
■ protect from excessive moisture  
■ TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.

**Directions**  
adults and children 6 years and over  
1 tablet daily; not more than 1 tablet in 24 hours  
ask a doctor  
children under 6 years of age  
ask a doctor  
consumers with liver or kidney disease  
ask a doctor

**Drug Facts (continued)**  
before use.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center. right away (1-800-222-1222).



5199416

**publix.**

R0619  
DISTRIBUTED BY PUBLIX SUPER MARKETS, INC.,  
3300 PUBLIX CORPORATE PARKWAY  
LAKELAND, FL 33811  
1-888-267-3037 publix.com  
PUBLIX GUARANTEES COMPLETE  
SATISFACTION OR YOUR MONEY BACK



5

# LORATADINE

loratadine tablet

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:56062-526
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LORATADINE (UNII: 7AJ03BO7QN) (LORATADINE - UNII:7AJ03BO7QN)	LORATADINE	10 mg

## Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

## Product Characteristics

Color	white (White to Off-White)	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	RX526
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:56062-526-69	10 in 1 BLISTER PACK; Type 0: Not a Combination Product	08/19/2003	
2	NDC:56062-526-31	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	08/19/2003	
3	NDC:56062-526-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	08/19/2003	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076134	08/19/2003	

**Labeler** - Publix Super Markets Inc (006922009)

**Registrant** - Sun Pharmaceutical Industries Inc. (146974886)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Ohm Laboratories Inc.		051565745	manufacture(56062-526)

Revised: 9/2019

Publix Super Markets Inc