TUSNEL COUGH SUPPRESSANT EXPECTORANT- dextromethorphan hbr, guaifenesin liquid

Llorens Pharmaceutical International Division, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

In each 5 mL

Dextromethorphan - 10 mg

Guaifenesin - 100 mg

Purpose

Cough Suppressant

Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation
- helps loosen phlegm (mucus)
- helps thin bronchial secretions to make coughs more productive

Warnings

Do not use if you are now taking a prescritpion monamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's diesease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescirption drug contains an MAOI, aska doctor or pharmacist before taking this product.

Do not use if you have ever had an allergic reaction to any of the ingredients in this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or head that lasts. These could be signs of a serious condition.

If pregnant or breast-feeing, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions:

- take every 4 hours as needed, or as directed by a doctor
- do not take more than 6 doses in 24 hours
- do not exceed recommended dose

Age	Dose
Adults and children over 12 years and over	2 teaspoonfuls (10 mL)
Children under 12 years	do not use

Inactive ingredients cherry flavor, citric acid, FD&C #40, menthol, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate, sucralose, sucrose.

Questions or comments? 1-866-595-5598

Llorens Pharmaceutical

Drug Facts Active Ingredients (per 5 mL teaspoonful) Dextromethorphan HBr, USP 10mg Cough Suppressant Guaifenesin, USP 100mg..... Expectorant temporarily relieves cough due to minor throat and bronchial irritation · helps loosen phlegm (mucus) · helps thin bronchial secretions to make coughs more productive Do not use if you are now taking a prescription monoamine

DO NOT USE IT YOU GIT I TOWN TOWING A PRESCRIPTION THAT THE WORK OF THE WORK O pharmacist before taking this product.

Do not use if you have ever had an allergic reation to any of the ingredients in this product.

Ask a doctor before use if you have

a cough with too much phlegm (mucus) - a cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema.

Stop use and ask a doctor if cough lasts more than 7 days, comes back or is accompanied by fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional

Keep out of reach of children. In case of overdose, get cal help or contact a Poison Control Center (1-800-222-1222) right away.

COUGH **SUPPRESSANT EXPECTORANT**

Guaifenesin Dextromethorphan

- CHERRY **FLAVOR**
- √ Controls Cough
- √ Loosens and Relieves Chest Congestion
- √ Alcohol Free

Compare to the active ingredients in Robitussin® DM*

16 FL OZ (473 mL)

Drug Facts (continued)

DIRECTIONS

onections

toke every 4 hours as needed, or as directed by a doctor
do not take more than 6 doses in 24 hours
do not exceed recommended dose

Adults and children 12 years and over 2 teaspoonfuls (10 mL)

OTHER INFORMATION

· each teaspoonful (5 mL) contains: sodiu store at room temperature 15°-30°C (59°-86°F)

protect from freezing - do not refrigerate - protect from light Pharmacists-Preserve and dispense in a tight, light resistant container with a child resistant cap as defined in the USP TAMPER-EVIDENT: Do not use if foil seal over bottle opening is torn broken or missing.

Inactive ingredients: cherry flavor, citric acid, FD&C Red #40, menthol, methylparaben, propylene glycol, propylpa purified water, sodium citrate, sucralose, sucrose.

Questions or comments? 1-866-595-5598

This product is not manufactured or distributed by the owner of the registered trademark Robitussin®DM.

THIS IS A BULK CONTAINER NOT INTENDED FOR DISPENSING Manufactured by: Code #: L-159 Rev.: 05/22

Llorens Pharmaceutical International Division, Inc. 7080 NW 37th Court Miami, FL 33147 | www.LlorensPharm.com

University Llorens Pharmaceutical

Exp Date :



TUSNEL COUGH SUPPRESSANT EXPECTORANT

dextromethorphan hbr, quaifenesin liquid

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:54859-509

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name Strength

ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
MENTHOL (UNII: L7T10EIP3A)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SUCROSE (UNII: C151H8M554)	
f .	

l	P	Packaging			
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:54859-509- 16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2022	

Marketing In	Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part341	05/01/2019		

Labeler - Llorens Pharmaceutical International Division, Inc. (037342305)

Revised: 7/2022 Llorens Pharmaceutical International Division, Inc.