

BMD SOLUBLE- bacitracin methylene disalicylate powder, for solution
Zoetis Inc.

BMD® Soluble

BMD® Soluble

(bacitracin methylenedisalicylate soluble powder)

Proportioner Pack

For poultry and livestock drinking water

Broiler and replacement chickens

An aid in prevention and control of necrotic enteritis caused by *Clostridium perfringens* susceptible to bacitracin methylenedisalicylate.

Growing turkeys

An aid in control of transmissible enteritis (blue comb, mud fever) in growing turkeys complicated by organisms susceptible to bacitracin methylenedisalicylate.

Growing quail

For the prevention of ulcerative enteritis due to *Clostridium colinum* susceptible to bacitracin methylenedisalicylate.

Swine

For use in the treatment of swine dysentery associated with *Brachyspira hyodysenteriae*.

Active ingredient

Contains 51.2 g bacitracin activity from bacitracin methylenedisalicylate equivalent to 200 g bacitracin activity per pound or to 18,520 units bacitracin (master standard) per gram.

Ingredients

Granulated sugar, sodium bicarbonate, anti-caking agent.

Caution

Not for use in humans. Keep out of reach of children.
For oral use in animals only.

Concentrated

For water medicators.

Directions for use

Animal	Condition	Prevention Control	
Broiler and replacement chickens	Necrotic enteritis caused by <i>Clostridium perfringens</i> susceptible to bacitracin methylenedisalicylate.	100 mg / gal	200 - 400 mg / gal
Growing turkeys	Transmissible enteritis (blue comb, mud fever) complicated by organisms susceptible to bacitracin methylenedisalicylate.	-	400 mg / gal
Growing quail	Ulcerative enteritis due to <i>Clostridium colinum</i> susceptible to bacitracin methylenedisalicylate..	400 mg / gal	-
Swine	Swine dysentery (bloody scours) associated with <i>Brachyspira hyodysenteriae</i> .	-	1000 mg / gal

Net Weight

4.1 oz (116.2 g)

Store between 20°C -25°C (68°F - 77°F);

excursions permitted between 15°C - 37°C (59°F - 99°F).

Restricted Drug (California) - Use only as directed.

Prepare a fresh solution daily and use as sole source of drinking water.

Mixing instructions

For Proportioners

Select the treatment dosage. Set the proportioner at the desired delivery rate. To prepare the proportioner's stock solution, place the indicated quantity of BMD Soluble in a 2-gallon container, fill with water and stir until dissolved.

For Tanks

One package of BMD Soluble will medicate approximately 50 gallons of water at a dosage of 1000 mg/gallon.

Treatment dosage	PROPORTIONER SET AT:	
	1 ounce/gallon	2 ounces/gallon
100 mg/gal	1/2 pack	1/4 pack
200 mg/gal*	1 pack	1/2 pack
400 mg/gal	2 packs	1 pack
1000 mg/gal	5 packs	2-1/2 packs

* 200 mg/gallon is equivalent to about 100 g of feed grade BMD per ton of feed

*200 mg/gallon is equivalent to about 100 g of feed grade BMD per ton of feed

To control an outbreak, start medication at first clinical signs of disease. Consult a diagnostic laboratory or veterinarian to determine the diagnosis and advice regarding the optimal level of drug. For necrotic enteritis in broiler and replacement chickens, administer continuously 5-7 days or as long as clinical signs persist, then reduce medication to prevention levels (100 mg/gal).

For transmissible enteritis in turkeys, administer continuously as long as clinical signs persist. For swine dysentery start medication at first signs of the disease or at time of exposure and administer continuously for 7 days or until signs of dysentery disappear. Treatment not to exceed 14 days.

CAUTION: Not for use in swine weighing more than 250 lb.

zoetis

Distributed by:
Zoetis Inc.
Kalamazoo, MI 49007



Net Weight 4.1 oz (116.2 g)

Restricted Drug (California) - Use only as directed.

40038555

Approved by FDA under NADA # 065-470

PRINCIPAL DISPLAY PANEL - bag label

BMD[®] Soluble

(bacitracin methylenedisalicylate soluble powder)

Proportioner Pack

For poultry and livestock drinking water

Follow directions for use and mixing instructions on packet label.

Store between 20°C - 25°C (68°F - 77°F); excursions permitted between 15°C - 37°C (59°F - 99°F).

CAUTION: Not for use in humans. Keep out of reach of children. For oral use in animals only.

Contents: 50 packs, 4.1 oz (116.2 g) each

zoetis

Distributed by:
Zoetis Inc., Kalamazoo, MI 49007

Take Time



Observe Label
Directions

10006716

40038554

Restricted Drug (California) - Use only as directed.

Approved by FDA under NADA # 065-470

Lot:	IMPRINT AREA
Exp.:	



BMD SOLUBLE

bacitracin methylene disalicylate powder, for solution

Product Information

Product Type	OTC TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:54771-1320
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BACITRACIN METHYLENEDISALICYLATE (UNII: JGV6OJ52FT) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN	200 mg in 0.45 kg

Product Characteristics

Color	brown (Tan to light brown)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54771-1320-1	50 in 1 CONTAINER		
1		0.1162 kg in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA065470	03/22/2007	

Labeler - Zoetis Inc. (828851555)

Revised: 7/2022

Zoetis Inc.