

COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated
VALU MERCHANDISERS COMPANY

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

1163-BST-2021-1118

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- for the temporary relief of the following cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

-
- take 2 caplets every 4 hours

adults and children 12 years and over	<ul style="list-style-type: none">▪ swallow whole – do not crush, chew, or dissolve▪ do not take more than 10 caplets in 24 hours
children under 12 years	<ul style="list-style-type: none">▪ ask a doctor

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, D&C yellow #10, FD&C blue #2, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

PRINCIPAL DISPLAY PANEL

COMPARE TO THE ACTIVE INGREDIENTS IN TYLENOL® COLD & FLU SEVERE†

Best Choice®

SEVERE

Cold & Flu

Headache, Fever, Sore Throat, Nasal Congestion,

Cough & Chest Congestion, Clears Out Mucus

Pain Reliever / Fever Reducer - Acetaminophen

Nasal Decongestant - Phenylephrine HCl

Cough Suppressant - Dextromethorphan HBr

Expectorant - Guaifenesin

Cool Taste

24 CAPLETS

NC

NC

NC

Drug Facts (continued)

Warnings
 Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
 ■ liver disease
 ■ heart disease ■ high blood pressure ■ thyroid disease
 ■ diabetes ■ trouble urinating due to an enlarged prostate gland
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Directions
 ■ do not take more than directed (see overdose warning)
 adults and children 12 years and over
 ■ take 2 caplets every 4 hours
 ■ swallow whole – do not crush, chew, or dissolve
 ■ do not take more than 10 caplets in 24 hours
 children under 12 years
 ■ ask a doctor

Drug Facts (continued)

When using this product do not exceed recommended dosage
 ■ when using this product do not exceed recommended dosage
 ■ when using this product do not exceed recommended dosage
 ■ when using this product do not exceed recommended dosage

Stop use and ask a doctor if
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present ■ new symptoms occur
 ■ cough comes back or occurs with rash or headache that lasts
 ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
 ■ nervousness, dizziness, or sleeplessness occur
 ■ throat gets worse or lasts more than 7 days

Keep out of reach of children.
 Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

If pregnant or breast-feeding, ask a health professional before use. These could be signs of a serious condition.
 ■ cough gets worse or lasts more than 7 days
 ■ nervousness, dizziness, or sleeplessness occur
 ■ throat gets worse or lasts more than 7 days

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin
 ■ when using this product do not exceed recommended dosage
 ■ when using this product do not exceed recommended dosage
 ■ when using this product do not exceed recommended dosage

Stop use and ask a doctor if
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present ■ new symptoms occur
 ■ cough comes back or occurs with rash or headache that lasts
 ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
 ■ nervousness, dizziness, or sleeplessness occur
 ■ throat gets worse or lasts more than 7 days

Keep out of reach of children.
 Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

If pregnant or breast-feeding, ask a health professional before use. These could be signs of a serious condition.
 ■ cough gets worse or lasts more than 7 days
 ■ nervousness, dizziness, or sleeplessness occur
 ■ throat gets worse or lasts more than 7 days

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin
 ■ when using this product do not exceed recommended dosage
 ■ when using this product do not exceed recommended dosage
 ■ when using this product do not exceed recommended dosage

Stop use and ask a doctor if
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present ■ new symptoms occur
 ■ cough comes back or occurs with rash or headache that lasts
 ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
 ■ nervousness, dizziness, or sleeplessness occur
 ■ throat gets worse or lasts more than 7 days

Keep out of reach of children.
 Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

If pregnant or breast-feeding, ask a health professional before use. These could be signs of a serious condition.
 ■ cough gets worse or lasts more than 7 days
 ■ nervousness, dizziness, or sleeplessness occur
 ■ throat gets worse or lasts more than 7 days

Drug Facts

Active ingredients (in each caplet) Purpose
 Acetaminophen 325 mg Pain reliever/fever reducer
 Dextromethorphan HBr 10 mg Cough suppressant
 Guaifenesin 200 mg Expectorant
 Phenylephrine HCl 5 mg Nasal decongestant

Drug Facts (continued)

Uses ■ for the temporary relief of the following cold/flu symptoms:
 ■ minor aches and pains ■ headache ■ sore throat
 ■ nasal congestion ■ cough
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
 ■ temporarily reduces fever

Best Choice®

SEVERE Cold & Flu

Headache, Fever, Sore Throat, Nasal Congestion, Cough & Chest Congestion, Clears Out Mucus

Pain Reliever / Fever Reducer - **Acetaminophen**
 Nasal Decongestant - **Phenylephrine HCl**
 Cough Suppressant - **Dextromethorphan HBr**
 Expectorant - **Guaifenesin**

Cool Taste
24 CAPLETS

PROVIDED BY:
 VALU MERCHANDISERS, CO.
 5000 KANSAS AVE
 KANSAS CITY, KS 66106

You may report serious side effects to:
 881 Main Street
 Lumberton, NJ 08048

Best Choice
 100% Guaranteed
 www.bestchoicebrand.com

†This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Cold & Flu Severe.

DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN

F116305BST_R0

Drug Facts (continued)

Other information
 ■ store between 20-25°C (68-77°F) in a dry place
 ■ retain carton for complete product information

Inactive ingredients
 acetylsalicylic acid, croscarmellose sodium, colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone,

Drug Facts (continued)

D&C yellow #10, FD&C blue #2, flavor, magnesium stearate, malto-dextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

BEST CHOICE

0 70038 58079 9



COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-163
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (UNII: 2S7830E561)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	yellow	Score	no score
Shape	OVAL	Size	19mm
Flavor	MINT	Imprint Code	AAA;1136
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-163-05	2 in 1 CARTON	03/09/2012	12/31/2025
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	03/09/2012	12/31/2025

Labeler - VALU MERCHANDISERS COMPANY (868703513)

Revised: 9/2023

VALU MERCHANDISERS COMPANY