

**FEXOFENADINE HYDROCHLORIDE AND PSEUDOEPHEDRINE HYDROCHLORIDE-
fexofenadine hydrochloride and pseudoephedrine hydrochloride tablet,
extended release
CVS Health Corp**

**Fexofenadine Hydrochloride and Pseudoephedrine Hydrochloride Extended-
Release Tablets USP, 60 mg/120 mg**

Active ingredient(s)

Fexofenadine HCl USP, 60 mg

Pseudoephedrine HCl USP, 120 mg

Purpose

Antihistamine

Nasal decongestant

Use(s)

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have difficulty swallowing

Ask a doctor before use if you have

- heart disease
- thyroid disease

- glaucoma
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.

When using this product

- **do not take more than directed**
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)
- the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).

Stop use and ask doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- you get nervous, dizzy, or sleepless

If pregnant or breast-feeding

ask a health professional before use.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not divide, crush, chew or dissolve the tablet; swallow tablet whole

adults and children 12 years of age and over	take 1 tablet with a glass of water every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other information

- safety sealed: do not use if carton is opened or if individual blister units are torn or opened
- store between 20° to 25°C (68° to 77°F) store between 20° to 25°C (68° to 77°F)
- FDA approved dissolution test specifications differ from USP.
- FDA approved organic impurities test procedure differs from USP Procedure 1.

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, ferric oxide (iron oxide yellow), HPMC 2910 / hypromellose (6 Cps), hydroxypropyl cellulose, hypromellose (methocel K100M DC2), macrogol (polyethylene glycol MW 400), macrogol (polyethylene glycol MW 8000), magnesium stearate, microcrystalline cellulose (avicel PH 101), microcrystalline cellulose (avicel PH102), pregelatinized starch.

Questions?

Call **1-888-375-3784**

Principal Display Panel

Blister carton



No Coating Area

Drug Facts (continued)

- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)
- the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).
- do not take with fruit juices (see Directions)
- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- you get nervous, dizzy, or sleepless
- if program or breast-feeding, ask a health professional before use
- Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
do not chew, crush, chew or dissolve the tablet; swallow tablet whole

adults and children 12 years of age and over
take 1 tablet with a glass of water every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours

children under 12 years of age
do not use

adults 65 years of age and older
ask a doctor

consumers with kidney disease
ask a doctor

Questions or comments? call toll-free weekdays 9 AM to 8 PM EST. Call 1-888-375-3784

Inactive ingredients coloidal silicon dioxide, croscarmellose sodium, ferric oxide (iron oxide yellow), hydroxypropyl methylcellulose, hypromellose (K100M DC2), macrogol (polyethylene glycol) MW 8000, magnesium stearate, microcrystalline cellulose (rancel PH11), microcrystalline cellulose (wavel PH102), pregelatinized starch

Other information
safety sealed do not use if carton is opened or if individual blister units are torn or opened

store between 20° and 25°C (68° and 77°)

store in original container

USP Procedure 1

do not take more than directed



No Coating Area

Drug Facts

Active ingredients (in each tablet)
Pseudoephedrine HCl USP, 120 mg; Fexofenadine HCl USP, 60 mg; Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
runny nose
itchy, watery eyes
sneezing

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies due to the common cold, hay fever or other upper respiratory allergies

temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies

temporarily relieves sinus congestion and pressure

temporarily relieves itchy throat through the nose

temporarily relieves itchy, watery eyes

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
runny nose
itchy, watery eyes
sneezing

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
runny nose
itchy, watery eyes
sneezing

When using this product
do not take more than directed

ask a doctor before use if you have:
heart disease
high blood pressure
glaucoma
diabetes
kidney disease. Your doctor should determine if you need a different dose.

if you have difficulty swallowing

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug

if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

if you have difficulty swallowing

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug

LOT
BP

No Coating Area



#197233

CVS Health. Compare to the active ingredients in Allegra-D® 12 Hour Allergy & Congestion Tablets* NDC 51316-015-30

ALLERGY & CONGESTION RELIEF

Allergy Relief D

FEXOFENADINE HCl 60 mg/Antihistamine
PSEUDOEPHEDRINE HCl 120 mg/Nasal decongestant
EXTENDED RELEASE TABLETS, USP

INDOOR/OUTDOOR RELIEF OF:

- Nasal & sinus congestion due to colds or allergies
- Sneezing; Runny nose; Itchy, watery eyes & Itchy nose or throat due to allergies

Non-drowsy

30 EXTENDED-RELEASE TABLETS



Actual Size



*This product is not manufactured or distributed by Chattem, Inc. (part of the Sanofi Group), distributor of Allegra-D 12 Hour Allergy & Congestion Tablets. Allegra-D is a registered trademark of Aventisub LLC.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2023 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
Made in India
V-13254



Rev: 01/23

CVS Health.

ALLERGY & CONGESTION RELIEF

Allergy Relief D

FEXOFENADINE HCl 60 mg/Antihistamine
PSEUDOEPHEDRINE HCl 120 mg/Nasal decongestant
EXTENDED RELEASE TABLETS, USP

INDOOR/OUTDOOR RELIEF OF:

- Nasal & sinus congestion due to colds or allergies
- Sneezing; Runny nose; Itchy, watery eyes & Itchy nose or throat due to allergies

Non-drowsy



FEXOFENADINE HYDROCHLORIDE AND PSEUDOEPHEDRINE HYDROCHLORIDE

fexofenadine hydrochloride and pseudoephedrine hydrochloride tablet, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51316-015
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FEXOFENADINE HYDROCHLORIDE (UNII: 2S068B75ZU) (FEXOFENADINE - UNII:E6582LOH6V)	FEXOFENADINE HYDROCHLORIDE	60 mg
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	120 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE 101 (UNII: 7T9FYH5QMK)	
Croscarmellose Sodium (UNII: M28OL1HH48)	
Ferric Oxide Yellow (UNII: EX438O2MRT)	
Silicon Dioxide (UNII: ETJ7Z6XBU4)	
Magnesium Stearate (UNII: 70097M6I30)	
Polyethylene Glycol 400 (UNII: B697894SGQ)	
Hypromellose 2208 (100000 Mpa.S) (UNII: VM7F0B23ZI)	
HYDROXYPROPYL CELLULOSE (45000 WAMW) (UNII: 8VAB711C5E)	
Hypromellose 2910 (6 Mpa.S) (UNII: 0WZ8WG20P6)	
Polyethylene Glycol 8000 (UNII: Q662QK8M3B)	
Starch, Corn (UNII: O8232NY3SJ)	
MICROCRYSTALLINE CELLULOSE 102 (UNII: PNR0YF693Y)	

Product Characteristics

Color	WHITE (one white to off-white color layer and other light yellow to yellow color)	Score	no score
Shape	CAPSULE	Size	16mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51316-015-20	4 in 1 CARTON	03/17/2023	
1		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:51316-015-30	6 in 1 CARTON	03/17/2023	

2	5 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215434	06/03/2022	

Labeler - CVS Health Corp (062312574)

Revised: 12/2025

CVS Health Corp