

STILA SHEER COLOR TINTED MOISTURIZER BS SPF20 (ALL SHADES)- titanium dioxide and zinc oxide lotion

Stila Styles, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Stila Sheer color tinted moisturizer broad spectrum SPF 20 (All Shades)

Drug Facts

| Active Ingredients | Purpose |
|---------------------------|----------------|
| Titanium Dioxide 5.00% | sunscreen |
| Zinc Oxide 5.00% | sunscreen |

Uses

"Helps prevent sunburn"

"If used as directed with other protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun."

Warnings

For external use only

Do not use on damaged or broken skin

Stop use and ask a doctor if rash occurs

When using this product keep out of eyes. Rinse with water to remove.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Apply liberally 15 minutes before sun exposure
- Reapply:
 - After 80 minutes of swimming or sweating
 - Immediately after towel drying
 - At least every 2 hours
- Children under 6 months: Ask a doctor
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - Limit time in the sun, especially from 10 am – 2 pm
 - Wear long-sleeved shirts, pants, hats and sunglasses

Inactive Ingredients

aluminum hydroxide, anthemis nobilis (chamomile) flower extract, boron nitride, butylene glycol, camellia oleifera (camilia oleifera) leaf extract, caprylyl glycol, cyclohexasiloxane, cyclopentasiloxane, dextrin palmitate, dimethicone, ginkgo biloba leaf extract, glycerin, hexylene

glycol, lysine, magnesium ascorbyl phosphate, magnesium chloride, octyldodecyl neopentanoate, palmitic acid, panax ginseng (ginseng) root extract, peg/ppg-18/18 dimethicone, phenoxyethanol, phytantriol, potassium chloride, propylene glycol, retinyl palmitate, sodium chloride, sodium dehydroacetate, sodium hyaluronate, stearic acid, tetrahexyldecyl ascorbate, tocopherol, tocopheryl acetate, triethoxycaprylylsilane, water (aqua), zinc chloride, **may contain (+/-)**: mica, titanium dioxide (ci 77891), iron oxides (ci 77491, 77492, 77499) <iln c121004>.

Other Information

Protect this product from excessive heat and direct sun

Questions?

stilacosmetics.com

dist. by stila styles, llc
los angeles, ca 90027

PRINCIPAL DISPLAY PANEL - 50 ml Tube Carton

stila

sheer color
tinted moisturizer
broad spectrum SPF 20


oil-free
water-resistant (80 minutes)

1.7 fl. oz./50 ml

SHADE LABEL
FPO
1 X .5, CR= .0625

stila

sheer color
tinted moisturizer
broad spectrum SPF 20



oil-free
water-resistant (80 minutes)

1.7 fl. oz./50 ml

Drug Facts

Active Ingredients

| | |
|------------------|-------|
| Titanium Dioxide | 5.00% |
| Zinc Oxide | 5.00% |

| |
|-----------|
| Purpose |
| sunscreen |
| sunscreen |

Uses:

"Helps prevent sunburn"
"If used as directed with other protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun."

Warnings:

For external use only

usage: blend over entire face with fi
benefits:

- light-diffusing pigments help dimin
- ginseng root extract and ginkgo b
- help prevent premature aging
- chamomile helps eliminate rednes
- vitamins A and E condition and pi
- advanced emollient system helps t
- sheer to medium coverage for all:

NO COVERTING

ngertips. layer for additional coverage.

ish the look of fine lines and wrinkles
iloba leaf extract provide potent antioxidant benefits and

s, soothes the skin and improves skin elasticity
rovide antioxidant benefits
hydrate and moisturize the skin
skin types

Do not use on damaged or broken skin
Stop use and ask a doctor if rash occurs
When using this product keep out of eyes. Rinse with water to remove.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions:

- Apply liberally 15 minutes before sun exposure
- Reapply:
 - After 80 minutes of swimming or sweating
 - Immediately after towel drying
 - At least every 2 hours
- Children under 6 months: Ask a doctor
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - Limit time in the sun, especially from 10 am – 2 pm
 - Wear long-sleeved shirts, pants, hats and sunglasses

Inactive Ingredients

aluminum hydroxide, anthemis nobilis (chamomile) flower extract, boron nitride, butylene glycol, camellia oleifera (camilia oleifera) leaf extract, caprylyl glycol, cyclohexasiloxane, cyclopentasiloxane, dextrin palmitate, dimethicone, ginkgo biloba leaf extract, glycerin, hexylene glycol, lysine, magnesium ascorbyl phosphate, magnesium chloride, octyldodecyl neopentanoate, palmitic acid, panax ginseng (ginseng) root extract, peg/ppg-18/18 dimethicone, phenoxyethanol, phytantriol, potassium chloride, propylene glycol, retinyl palmitate, sodium chloride, sodium dehydroacetate, sodium hyaluronate, stearic acid, tetrahexyldecyl ascorbate, tocopherol, tocopheryl acetate, triethoxycaprylylsilane, water (aqua), zinc chloride, **may contain (+/-):** mica, titanium dioxide (ci 77891), iron oxides (ci 77491, 77492, 77499) <iln c121004>.

Other Information

Protect this product from excessive heat and direct sun


Questions? stilacosmetics.com

UPC LABEL
FPO
1.25 X .75, CR=.0625

stilacosmetics.com

dist. by © stila styles, llc
los angeles, ca 90027

biorius sprl • 2 rue rené descartes
b-7000 mons, belgium
made in u.s.a. S556



STILA SHEER COLOR TINTED MOISTURIZER BS SPF20 (ALL SHADES)

titanium dioxide and zinc oxide lotion

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:76049-556 |
| Route of Administration | TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|---------------|
| Titanium Dioxide (UNII: 15FIX9V2JP) (Titanium Dioxide - UNII:15FIX9V2JP) | Titanium Dioxide | 50 mg in 1 mL |
| Zinc Oxide (UNII: SOI2LOH54Z) (Zinc Oxide - UNII:SOI2LOH54Z) | Zinc Oxide | 50 mg in 1 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| aluminum hydroxide (UNII: 5QB0T2IUN0) | |
| chamaemelum nobile flower (UNII: O2T154T6OG) | |
| boron nitride (UNII: 2U4T60A6YD) | |
| butylene glycol (UNII: 3XUS85K0RA) | |
| camellia oleifera leaf (UNII: 5077EL0C60) | |
| caprylyl glycol (UNII: 00YIU5438U) | |
| cyclomethicone 6 (UNII: XHK3U310BA) | |
| cyclomethicone 5 (UNII: 0THT5PCI0R) | |
| dimethicone (UNII: 92RU3N3Y1O) | |
| ginkgo (UNII: 19FUJ2C58T) | |
| glycerin (UNII: PDC6A3C0OX) | |
| hexylene glycol (UNII: KEH0A3F75J) | |
| lysine (UNII: K3Z4F929H6) | |
| magnesium ascorbyl phosphate (UNII: 0R822556M5) | |
| magnesium chloride (UNII: 02F3473H9O) | |
| octyldodecyl neopentanoate (UNII: X8725R883T) | |
| palmitic acid (UNII: 2V16EO95H1) | |
| asian ginseng (UNII: CUQ3A77YXI) | |
| peg/ppg-18/18 dimethicone (UNII: 9H0AO7T794) | |

| | |
|--|--|
| phenoxyethanol (UNII: HIE492ZZ3T) | |
| phytantriol (UNII: 8LVI07A72W) | |
| potassium chloride (UNII: 660YQ98I10) | |
| propylene glycol (UNII: 6DC9Q167V3) | |
| vitamin a palmitate (UNII: 1D1K0N0VVC) | |
| sodium chloride (UNII: 451W47IQ8X) | |
| sodium dehydroacetate (UNII: 8W46YN971G) | |
| hyaluronate sodium (UNII: YSE9PPT4TH) | |
| stearic acid (UNII: 4ELV7Z65AP) | |
| tetrahexyldecyl ascorbate (UNII: 9LBV3F07AZ) | |
| tocopherol (UNII: R0ZB2556P8) | |
| .alpha.-tocopherol acetate (UNII: 9E8X80D2L0) | |
| triethoxycaprylsilane (UNII: LDC331P08E) | |
| water (UNII: 059QF0K00R) | |
| zinc chloride (UNII: 86Q357L16B) | |
| mica (UNII: V8A1AW0880) | |
| ferric oxide red (UNII: 1K09F3G675) | |
| ferric oxide yellow (UNII: EX438O2MRT) | |
| ferrosoferric oxide (UNII: XM0M87F357) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:76049-556-01 | 1 in 1 CARTON | | |
| 1 | | 50 mL in 1 TUBE | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part347 | 08/31/2013 | |

Labeler - Stila Styles, LLC (809192896)

Establishment

| Name | Address | ID/FEI | Business Operations |
|---------------|---------|-----------|------------------------|
| MANA PRODUCTS | | 078870292 | MANUFACTURE(76049-556) |

Revised: 11/2013

Stila Styles, LLC