

**GOOD SENSE COLD- acetaminophen, chlorpheniramine maleate,  
dextromethorphan hbr, phenylephrine hcl suspension  
L. Perrigo Company**

-----  
**Perrigo Multi-Symptom Cold Plus Drug Facts**

**Active ingredients (in each 5 mL)**

Acetaminophen 160 mg

Chlorpheniramine maleate 1 mg

Dextromethorphan HBr 5 mg

Phenylephrine HCl 2.5 mg

**Purposes**

Pain reliever/fever reducer

Antihistamine

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves
- minor aches and pains
- headache
- minor sore throat pain
- runny nose
- itchy nose or throat
- sneezing
- itchy, watery eyes due to hay fever
- cough due to minor throat and bronchial irritation as may occur with a cold
- nasal and sinus congestion
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- in a child under 4 years of age
- if your child has ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- to make a child sleepy

### **Ask a doctor before use if the child has**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

### **Ask a doctor or pharmacist before use if the child is**

taking

- the blood thinning drug warfarin
- sedatives or tranquilizers

### **When using this product**

- **do not exceed recommended dosage**
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness
- excitability may occur, especially in children

## Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

## Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical even if you do not notice any signs or symptoms.

## Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see Overdose warning)**
- **shake well before using**
- remove the child protective cap and squeeze your child's dose into the dosing cup
- may be given every 4 hours, while symptoms persist. Do not give more than 5 doses in 24 hours unless directed by a doctor.
- use enclosed dosing cup only. Keep for use with this product. Do not use any other dosing device.

Age	Dose
<b>children under 4 years of age</b>	<b>do not use</b>
children 4 to under 6 years of age	do not use unless directed by a doctor
children 6 to under 12 years of age	10 mL

## Other information

- **each 5 mL contains:** sodium 3 mg
- contains no aspirin
- store between 20-25°C (68-77°F). Protect from light.
- close cap tightly

## Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, tribasic sodium phosphate, xanthan gum

## Questions or comments?

**1-800-719-9260**

**Principal Display Panel**

GOODSENSE®

Ages 6 to 11 Years

Children's Triacting Suspension Multi-Symptom Cold Plus

Fever Reducer/Pain Reliever (Acetaminophen)

Cough Suppressant (Dextromethorphan HBr)

Nasal Decongestant (Phenylephrine HCl)

Antihistamine (Chlorpheniramine Maleate)

- Fever • Cough
- Runny & Stuffy Nose
- Aches & Pains
- Sore Throat

Grape Flavor

Compare to active ingredients of Children's Triaminic® Multi-Symptom Fever & Cold

4 FL OZ (118 mL)



**Drug Facts (continued)**

- diabetes ■ glaucoma
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

**Ask a doctor or pharmacist before use if the child is taking**

- sedatives or tranquilizers
- the blood thinning drug warfarin

**When using this product**

- do not exceed recommended dosage
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness
- excitability may occur, especially in children

**Stop use and ask a doctor if**

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical even if you do not notice any signs or symptoms.



**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- shake well before using
- remove the child protective cap and squeeze your child's dose into the dosing cup
- may be given every 4 hours, while symptoms persist. Do not give more than 5 doses in 24 hours unless directed by a doctor.
- use enclosed dosing cup only. Keep for use with this product. Do not use any other dosing device.

Age	Dose
children under 4 years of age	do not use
children 4 to under 6 years of age	do not use unless directed by a doctor
children 6 to under 12 years of age	10 mL

**Other Information**

- each 5 mL contains: sodium 3 mg
- contains no aspirin



**Drug Facts (continued)**

- store at 20-25°C (68-77°F). Protect from light.
- close cap tightly

**Inactive Ingredients** anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, tribasic sodium phosphate, xanthan gum

**Questions or comments? 1-800-719-9260**

**DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING**

Gluten Free  
Distributed By  
**Perrigo®**  
Allagan, MI 49010



**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

GOODSENSE®  
Children's  
Triacting Suspension  
**Multi-Symptom Cold Plus**  
Grape Flavor

NDC 0113-0903-26  
GOODSENSE®  
Ages 6 to 11 Years

Children's  
Triacting Suspension  
**Multi-Symptom Cold Plus**

Fever Reducer/Pain Reliever  
(Acetaminophen)  
Cough Suppressant  
(Dextromethorphan HBr)  
Nasal Decongestant (Phenylephrine HCl)  
Antihistamine  
(Chlorpheniramine Maleate)




- Fever • Cough
- Runny & Stuffy Nose
- Aches & Pains
- Sore Throat

Compare to active ingredients of  
Children's Triaminic®  
Multi-Symptom Fever & Cold

4 FL OZ (118 mL)



90326 C2 C5



**Drug Facts**

**Active Ingredients (In each 5 mL)**

Acetaminophen 160 mg.....Pain reliever/fever reducer  
Chlorpheniramine maleate 1 mg.....Antihistamine  
Dextromethorphan HBr 5 mg.....Cough suppressant  
Phenylephrine HCl 2.5 mg.....Nasal decongestant

**Purposes**

**Uses**

- temporarily relieves
- minor aches and pains ■ headache
- minor sore throat pain ■ runny nose
- itchy nose or throat ■ sneezing
- itchy, watery eyes due to hay fever
- nasal and sinus congestion
- cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount.

■ with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- in a child under 4 years of age
- if your child has ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- to make a child sleepy

**Ask a doctor before use if the child has**

- liver disease ■ heart disease
- high blood pressure ■ thyroid disease

**GOOD SENSE COLD**  
acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, phenylephrine hcl suspension

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0113-0903
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	1 mg in 5 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

Inactive Ingredients	
Ingredient Name	Strength
<b>CARRAGEENAN</b> (UNII: 5C69YCD2YJ)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>CALCIUM SULFATE, UNSPECIFIED FORM</b> (UNII: WATODDB505)	
<b>SODIUM PHOSPHATE, TRIBASIC</b> (UNII: A752Q30A6X)	

Product Characteristics			
<b>Color</b>	PURPLE (opaque)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE	<b>Imprint Code</b>	
<b>Contains</b>			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-0903-26	1 in 1 CARTON	12/22/2010	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/22/2010	

**Labeler** - L. Perrigo Company (006013346)

Revised: 10/2023

L. Perrigo Company