

**LOPERAMIDE HYDROCHLORIDE- loperamide hydrochloride tablet**  
**Chain Drug Consortium, LLC**

-----  
**Loperamide Hydrochloride**

***Drug Facts***

**Active ingredient (in each caplet)**

Loperamide Hydrochloride USP, 2 mg

**Purpose**

Anti-diarrheal

**Use**

controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert**

Do not use if you have ever had a rash or other allergic reaction to loperamide hydrochloride

**Heart alert**

Taking more than directed can cause serious heart problems or death

**Do not use** if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product** tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding**,ask a health professional before use.

**Keep out of reach of children.**In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

### Other information

- store between 20° - 25°C (68° - 77°F)
- see side panel for lot number and expiration date
- **TAMPER EVIDENT: THIS PRODUCT IS PROTECTED WITH SEALED BLISTER UNITS. DO NOT USE IF ANY ARE TORN OR BROKEN.**

### Inactive ingredients

anhydrous lactose, croscarmellose sodium, crospovidone, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

### Questions?

Call toll-free Monday-Friday 8:30 am to 5 pm EST at **1800-406-7984**.

Distributed By:  
Pharmacy Value Alliance, LLC

407 East Lancaster Avenue,  
Wayne, PA 19087

**PRINCIPAL DISPLAY PANEL - 2 mg Caplet Blister Pack Carton**

COMPARE TO THE ACTIVE  
INGREDIENT OF IMODIUM<sup>®</sup>A-D †

See New Warnings  
and Directions

Premier  
Value<sup>®</sup>

LOPERAMIDE HYDROCHLORIDE  
TABLETS USP, 2 mg

Anti-Diarrheal

Controls The Symptoms of Diarrhea

6 Caplets\*

Each caplet (\*capsule-shaped tablet)  
contains Loperamide Hydrochloride USP, 2 mg

INDEPENDENTLY TESTED

PV

SATISFACTION GUARANTEED

5197452



5197452



**Questions?** Call toll-free Monday-Friday 8:30 am to 5 pm EST at 1 800-406-7984.

**Directions**  
 ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea  
 ■ in night do as on chart. If possible, use weight to dose; otherwise, use age.  
 adults and children 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours

**Drug Facts (continued)**  
 Do not use if you have bloody or black stool  
 ■ Ask a doctor before use if you have  
 ■ fever  
 ■ a history of liver disease  
 ■ a history of abnormal heart rhythm  
 Ask a doctor before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.  
 When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.  
 Stop use and ask a doctor if  
 ■ symptoms get worse  
 ■ diarrhea lasts for more than 2 days  
 ■ you get abdominal swelling or bloating. These may be signs of a serious condition.  
 If pregnant or breast-feeding, ask a health professional before use.  
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Other Information**  
 ■ store between 20° - 25°C (68° - 77°F)  
 ■ see side panel for lot number and expiration date  
 ■ TAMPER EVIDENT: THIS PRODUCT IS PROTECTED WITH SEALED BLISTER UNITS. DO NOT USE IF ANY ARE TO RN OR BROKEN.

**Inactive Ingredients**  
 amyloride, croscarmellose sodium, croscopolone, D&C Yellow No. 10 aluminum lake, vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

**Drug Facts (continued)**  
 children 9-11 years 1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours  
 children 6-8 years 1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours  
 children 2-5 years ask a doctor (34 to 47 lb)  
 children under 2 years do not use (up to 33 lbs)

**Drug Facts**  
 Loperamide Hydrochloride USP, 2 mg (Anti-diarrhea)

**Use**  
 Use to control symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**  
 Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide hydrochloride

**Heart Alert:** Taking more than directed can cause serious heart problems or death

**Premier Value**

**LOPERAMIDE HYDROCHLORIDE TABLETS USP, 2 mg**

**Anti-Diarrheal**

**Controls The Symptoms of Diarrhea**

**6 Caplets\***  
 Each caplet (capsule-shaped tablet) contains Loperamide Hydrochloride USP, 2 mg

**COMPARE TO THE ACTIVE INGREDIENT OF IMODIUM® A-D<sup>+</sup>**

**See New Warnings and Directions**

**NON VARNISH**

**Keep the carton. It contains important information.**  
 \*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Imodium® A-D.

Lot No.: \_\_\_\_\_ Expiration Date: \_\_\_\_\_

**Pharmacy Value Alliance, LLC**  
 407 East Lancaster Avenue,  
 Wayne, PA 19087

**Pharmacy Value Alliance**

**Ro419**

**8 40986 02762 8**

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.



## LOPERAMIDE HYDROCHLORIDE

loperamide hydrochloride tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68016-123
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LOPERAMIDE HYDROCHLORIDE</b> (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>CROSPVIDONE (15 MPA.S AT 5%)</b> (UNII: 68401960MK)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>HYDROGENATED COTTONSEED OIL</b> (UNII: Z82Y2C65EA)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>POWDERED CELLULOSE</b> (UNII: SMD1X3XO9M)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	

### Product Characteristics

<b>Color</b>	green	<b>Score</b>	2 pieces
<b>Shape</b>	CAPSULE	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	123
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-123-06	6 in 1 BLISTER PACK; Type 0: Not a Combination Product	02/01/1993	

2	NDC:68016-123-18	18 in 1 BLISTER PACK; Type 0: Not a Combination Product	02/01/1993	
3	NDC:68016-123-24	24 in 1 BLISTER PACK; Type 0: Not a Combination Product	02/01/1993	
4	NDC:68016-123-97	96 in 1 BOTTLE; Type 0: Not a Combination Product	02/01/1993	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074091	02/01/1993	

**Labeler** - Chain Drug Consortium, LLC (101668460)

**Registrant** - Ranbaxy Pharmaceuticals Inc. (937890044)

## Establishment

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		051565745	manufacture(68016-123)

Revised: 1/2024

Chain Drug Consortium, LLC