ORAL SALINE LAXATIVE CHERRY LAXATIVE- sodium phosphate, dibasic and sodium phosphate, monobasic, unspecified form liquid Natureplex, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

ORAL SALINE LAXATIVE CHERRY LAXATIVE

Drug Facts

Active ingredients (in each tablespoon)	Purpose
Dibasic Sodium Phosphate 2.7g	Saline laxative
Monobasic Sodium Phosphate 7.2g	Saline laxative

Use

- relieves occasional constipation
- this product generally produces bowel movement in 30 minutes to 6 hours

Warnings

Dosage Warning

Taking more than the recommended dosage in 24 hours can be harmful.

Do not use

- if you have a previous kidney problem, congestive heart failure, impaired renal functions, decreased intervascular volume, dehydration, or uncorrected electrolyte abnormalities
- in children under 6 years old
- for bowel cleansing

Ask a doctor before use if you

- have abdominal pain, nausea, or vomiting
- have kidney disease
- have heart problems
- have a sudden change in bowel habits lasting more than 2 weeks
- have already used a laxative for more than 3 days
- are dehydrated
- are 55 years of age or older
- are on a sodium-restricted diet
- under a doctor's care for any medical condition
- are taking any other prescription or non-prescription medications

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

When using this product

- do not use more than directed. Serious side effects may occur from excess dosage.
- do not use for more than 3 days, without asking a doctor

Stop use and ask a doctor if

- you have rectal bleeding
- you do not have a bowel movement within 6 hours
- you have any symptoms that your body is losing more fluid than you are drinking
- you have early symptoms of dehydration including
 - thirstiness
 - dizziness
 - urinating less often than normal
 - vomiting

These symptoms may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away: 800-222-1222.

Directions (or as directed by a doctor)

- Use as a laxative for relief of occasional constipation
- Dilute dose in the table below with one glass (8 fl oz) of cool water. Drink, then follow with at least one additional glass (8 fl oz) of cool water. There is significant loss of fluid when using this product. To prevent dehydration, drink plenty of clear liquids after use.
- Do not use for bowel cleansing
- There are 3 teaspoons in 1 TABLEspoon
- Do not take more unless directed by a doctor. See Warnings

Ages (years)	Dosage	Maximum dose
12 and older	1 TABLEspoon	2 to 3 TABLEspoons [*]
10 to 11	1 TABLEspoon [*]	1 TABLEspoon [*]
6 to 9	1 teaspoons*	2 teaspoons*
Under 6	ASK A DOCTOR	ASK A DOCTOR

⁶ SINGLE DAILY DOSAGE. DO NOT TAKE MORE OF THIS PRODUCT IN A 24-HOUR PERIOD

Other information

- each tables poon contains: 1.6 g sodium
- saline laxatives increase retention of water in the intestines, promoting bowel movements
- this product generally produces a bowel movement in 30 minutes to 6 hours
- this product is sugar-free
- do not freeze
- store at 15 to 30°C (59 to 86°F)
- **Tamper Evident:** DO NOT USE IF SEAL ON BOTTLE IS PUNCTURED OR MISSING.

Inactive ingredients

flavor, glycerin, purified water, saccharin sodium, sodium benzoate

Questions or comments?

866-323-0107 or www.natureplex.com

PRINCIPAL DISPLAY PANEL - 45 mL Bottle Carton

FOR DIRECTIONS LIFT SIDE PANEL

*Natureplex*TM

ORAL SALINE NDC 67234-041-01

Laxative Phosphate

For Relief Of Occasional Constipation

Cherry Splash

1.5 Fl Oz (45 mL)



NOTTAKE MORE OF THIS PRODUCT IN A 24-HOUR PERIOD ▼ from excess dosage. ■ do not use for more than 3 days, without asking a doctor from excess dosage. ■ do not use for more than 3 days, without asking a doctor <i>Drug Facts</i> (<i>continued</i>) Stop use and ask a doctor If ■ you have rectal bleeding ■ you do not have a bowel movement within 6 hours = you have rectal bleeding = you do not have a bowel wou are drinking = you have early symptoms of dehydration including • thirstiness	octor. See War	consupation tass (8 fl oz) of cool water. Drink, then follow with at least There is significant loss of fluid when using this ity of clear liquids after use.	octor)		Drug Facts	Purpose Saline laxative Saline laxative Saline laxative Saline laxative sage in 24 hours can be harmful. sage in 24 hours can be harmful. stive heart failure, impaired renal corrected electrolyte abnormalities icorrected electrolyte abnormalities is lasting more than 2 weeks = have d = are 55 years of age or older = are edical condition = are taking any other y other drug. Take this product two or now other drugs work.	Drug Facts (continued) Use Traileves occasional constipation Ihis product gene minutes to 6 hours Varnings Warnings Taking more than the recommended do Dosage Warning: Taking more than the recommended do Dosage Warning: Taking more than the recommended do Warnings Do not use If you have a previous kidney problem, conge tunctons, decreased intervascular volume, dehydration, or urance in children under 6 years old Tor bowel cleansing Ask a doctor before use If you Inve a addem change in bowel hal aready used a laxative for more than 3 days aready used a laxative for more than 3 days aready do Ask a doctor or pharmacist before use If you are taking an prescription or non-prescription medications Ask a doctor or pharmacist before use If you are taking an more hours before or after other drugs. Laxatives may affect When using this product = do not use for more than 3 days, withom excess dosage. = do not use for more than 3 days, withom excess dosage. = do not use for more than 3 days, withom excess dosage. = do not use for more than 3 days, withom within 6 hours Brug Facts (continued) you have early symptoms of denydration Brow are addared before or after or beact of a port wave any symptoms of denydration	ask a health professional before use. In case of overdose, get medical help or contact a r 800-222-1222. <i>Dy a doctor</i>) <i>by a doctor</i>) <i>ceasional constitpation</i> with one glass (8 fi oz) of cool water. Drink, then follow with at least col water. There is significant loss of fluid when using this of drink plenty of clear liquids after use. nd, drink plenty of clear liquids after use. <i>Do sage</i> Maximum dose if 1 TABLEspoon* 1 1 TABLEspoon* 1 1 telsepoon* 2 to 3 TABLEspoon* 1 1 telsepoon* 2 to 3 TABLEspoon* 1 1 telsepoon* 2 to 3 TABLEspoon* 1 telsepoon* 1 telsepoon* 1 telsepoon* 1 telsepoon* 2 to 3 TABLEspoon* 1 telsepoon* 1 telsepoon* 2 to 3 TABLEspoon*
•	spoon 2 to 3 TABLEspoons* spoon* 1 TABLEspoon* ons* 2 teaspoons*	octor. See Warnings ge Maximum dose LEspoon* 2 to 3 TABLEspoons* LEspoon* 2 teaspoons*	east	east		y other drug. Take this product two or now other drugs work. 1. Serious side effects may occur	Ask a doctor or pharmacist before use if you are taking ar more hours before or after other drugs. Laxatives may affect When using this product ■ do not use more than directed	DCTOR ASK ADOCTOR RE OF THIS PRODUCT IN A 24-HOUR PERIOD
ASK A DOCTOR ASK A DOCTOR ASK A DOCTOR ASK A doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.		octor. See Warnings ge Maximum dose	least	east	ess d	d material and the second seco	have heart problems whave a sudden change in bowel hat already used a laxative for more than 3 days ware dehydrate on a sodium-restricted diet wunder a doctor's care for any m prescription or non-prescription medications	
east	least					Purpose Saline laxative Saline laxative		fessional before use. ose, get medical help or contact

Drug Facts (continued) If pregnant or breast-feeding, Keep out of reach of children. Poison Control Center right awa Poison Control Center right awa Drug Facts (continued) Drug Facts (continued) Drug facts (or as directed Bilute dose in the table below Dilute dose in the table below Do not use for bowel cleansi There are 3 teaspoons in 11 Do not take more unless dir Do not take more unless dir 10 to 11 6 to 9	*SINGLE DAILY DOSAGE. DO	
--	--------------------------	--

ORAL SALINE LAXATIVE CHERRY LAXATIVE

sodium phosphate, dibasic and sodium phosphate, monobasic, unspecified form liquid

		1 1		1				
Product Informa	tion							
Product Type		HUMAN OTC DRUG	N OTC DRUG Item Code (Source)			NDC:67234-041		
Route of Administra	ation	ORAL						
Active Ingredien	t/Active M	oiety						
	Ing	redient Name				Basis of Stre	ngth	Strength
		NSPECIFIED FORM (UN HR, SODIUM CATION - U				SODIUM PHOSPHAT DIBASIC, UNSPECIF FORM		1.4 g in 14.7868 mL
SODIUM PHO SPHATE, MONOBASIC, UNSPECIFIED FORM (UNII: 3980 JIH2SW) (PHOSPHATE ION - UNII:NK08 V8K8 HR, SODIUM CATION - UNII:LYR4M0 NH37)SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM6.2 g in 14.7							6.2 g in 14.7868 mL	
. .								
Inactive Ingredie	ents							
Ingredient Name GLYCERIN (UNII: PDC6A3C0OX)							Strength	
WATER (UNII: 059QF	· · · · ·							
SACCHARIN SODIUM		X40TY)						
SODIUM BENZOATE								
		,						
Product Charact	eristics							
Color			Sco	re				
Shape			Siz	e				
Flavor	lavor CHERRY Imprint			rint Co	Code			
Contains								
Packaging								
# Item Code		Package Descriptio	n		Mar	keting Start Date	Market	ing End Date
1 NDC:67234-041-01	1 in 1 CARTON	J			08/31	1/2012		
1	45 mL in 1 BO	TTLE; Type 0: Not a Com	nbination P	roduct				

Marketing Informat	ion		
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part334	08/31/2012	

Labeler - Natureplex, LLC (062808196)

Establishment

Name	Address	ID/FEI	Business Operations
Natureplex LLC		062808196	MANUFACTURE(67234-041)

Revised: 6/2018

Nature plex, LLC