

**AZELEX- azelaic acid cream**  
**Allergan, Inc.**

-----

**AZELEX<sup>®</sup>**  
**(azelaic acid cream) 20%**

**For Dermatologic Use Only**  
**Not for Ophthalmic Use**

**DESCRIPTION**

**AZELEX<sup>®</sup>** (azelaic acid cream) 20% contains azelaic acid, a naturally occurring saturated dicarboxylic acid.

Structural Formula:  $\text{HOOC}-(\text{CH}_2)_7-\text{COOH}$

Chemical Name: 1,7-heptanedicarboxylic acid

Empirical Formula:  $\text{C}_9\text{H}_{16}\text{O}_4$

Molecular Weight: 188.22

**Active Ingredient:** Each gram of **AZELEX<sup>®</sup>** contains azelaic acid..... 0.2 g (20% w/w).

**Inactive Ingredients:** cetearyl octanoate; glycerin; glyceryl stearate and cetearyl alcohol and cetyl palmitate and cocoglycerides; PEG-5 glyceryl stearate; propylene glycol; and purified water. Benzoic acid is present as a preservative.

**CLINICAL PHARMACOLOGY**

The exact mechanism of action of azelaic acid is not known. The following *in vitro* data are available, but their clinical significance is unknown. Azelaic acid has been shown to possess antimicrobial activity against *Propionibacterium acnes* and *Staphylococcus epidermidis*. The antimicrobial action may be attributable to inhibition of microbial cellular protein synthesis.

A normalization of keratinization leading to an anticomedonal effect of azelaic acid may also contribute to its clinical activity. Electron microscopic and immunohistochemical evaluation of skin biopsies from human subjects treated with **AZELEX<sup>®</sup>** cream demonstrated a reduction in the thickness of the stratum corneum, a reduction in number and size of keratohyalin granules, and a reduction in the amount and distribution of filaggrin (a protein component of keratohyalin) in epidermal layers. This is suggestive of the ability to decrease microcomedo formation.

**Pharmacokinetics**

Following a single application of **AZELEX<sup>®</sup>** cream to human skin *in vitro*, azelaic acid penetrates into the stratum corneum (approximately 3 to 5% of the applied dose) and other viable skin layers (up to 10% of the dose is found in the epidermis and dermis). Negligible cutaneous metabolism occurs after topical application. Approximately 4% of the topically applied azelaic acid is systemically absorbed. Azelaic acid is mainly excreted unchanged in the urine but undergoes some  $\beta$ -oxidation to shorter chain dicarboxylic acids. The observed half-lives in healthy subjects are approximately 45 minutes after oral dosing and 12 hours after topical dosing, indicating percutaneous absorption rate-limited kinetics.

Azelaic acid is a dietary constituent (whole grain cereals and animal products), and can be formed endogenously from longer-chain dicarboxylic acids, metabolism of oleic acid, and  $\psi$ -oxidation of monocarboxylic acids. Endogenous plasma concentration (20 to 80 ng/mL) and daily urinary excretion (4 to 28 mg) of azelaic acid are highly dependent on dietary intake. After topical treatment with

**AZELEX**<sup>®</sup> cream in humans, plasma concentration and urinary excretion of azelaic acid are not significantly different from baseline levels.

## **INDICATIONS AND USAGE**

**AZELEX**<sup>®</sup> cream is indicated for the topical treatment of mild-to-moderate inflammatory acne vulgaris.

## **CONTRAINDICATIONS**

**AZELEX**<sup>®</sup> cream is contraindicated in individuals who have shown hypersensitivity to any of its components.

## **WARNINGS**

**AZELEX**<sup>®</sup> cream is for dermatologic use only and not for ophthalmic use.

There have been isolated reports of hypopigmentation after use of azelaic acid. Since azelaic acid has not been well studied in patients with dark complexions, these patients should be monitored for early signs of hypopigmentation.

## **PRECAUTIONS**

### **General**

If sensitivity or severe irritation develop with the use of **AZELEX**<sup>®</sup> cream, treatment should be discontinued and appropriate therapy instituted.

### **Information for Patients**

Patients should be told:

1. To use **AZELEX**<sup>®</sup> cream for the full prescribed treatment period.
2. To avoid the use of occlusive dressings or wrappings.
3. To keep **AZELEX**<sup>®</sup> cream away from the mouth, eyes and other mucous membranes. If it does come in contact with the eyes, they should wash their eyes with large amounts of water and consult a physician if eye irritation persists.
4. If they have dark complexions, to report abnormal changes in skin color to their physician.
5. Due in part to the low pH of azelaic acid, temporary skin irritation (pruritus, burning, or stinging) may occur when **AZELEX**<sup>®</sup> cream is applied to broken or inflamed skin, usually at the start of treatment. However, this irritation commonly subsides if treatment is continued. If it continues, **AZELEX**<sup>®</sup> cream should be applied only once-a-day, or the treatment should be stopped until these effects have subsided. If troublesome irritation persists, use should be discontinued, and patients should consult their physician (see ADVERSE REACTIONS).

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Azelaic acid is a human dietary component of a simple molecular structure that does not suggest carcinogenic potential, and it does not belong to a class of drugs for which there is a concern about carcinogenicity. Therefore, animal studies to evaluate carcinogenic potential with **AZELEX**<sup>®</sup> cream were not deemed necessary. In a battery of tests ( assay, HGPRT test in Chinese hamster ovary cells, human lymphocyte test, dominant lethal assay in mice), azelaic acid was found to be nonmutagenic. Animal studies have shown no adverse effects on fertility.

### **Pregnancy Teratogenic Effects**

Embryotoxic effects were observed in Segment I and Segment II oral studies with rats receiving 2500 mg/kg/day of azelaic acid. Similar effects were observed in Segment II studies in rabbits given 150 to 500 mg/kg/day and in monkeys given 500 mg/kg/day. The doses at which these effects were noted were all within toxic dose ranges for the dams. No teratogenic effects were observed. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

### **Nursing Mothers**

Equilibrium dialysis was used to assess human milk partitioning *in vitro*. At an azelaic acid concentration of 25 mcg/mL, the milk/plasma distribution coefficient was 0.7 and the milk/buffer distribution was 1.0, indicating that passage of drug into maternal milk may occur. Since less than 4% of a topically applied dose is systemically absorbed, the uptake of azelaic acid into maternal milk is not expected to cause a significant change from baseline azelaic acid levels in the milk. However, caution should be exercised when **AZELEX**<sup>®</sup> cream is administered to a nursing mother.

### **Pediatric Use**

Safety and effectiveness in pediatric patients under 12 years of age have not been established.

### **Geriatric Use**

Clinical studies of **AZELEX**<sup>®</sup> cream did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects.

## **ADVERSE REACTIONS**

During clinical trials with **AZELEX**<sup>®</sup> cream, adverse reactions were generally mild and transient in nature. The most common adverse reactions occurring in approximately 1-5% of patients were pruritus, burning, stinging and tingling. Other adverse reactions such as erythema, dryness, rash, peeling, irritation, dermatitis, and contact dermatitis were reported in less than 1% of subjects. There is the potential for experiencing allergic reactions with use of **AZELEX**<sup>®</sup> cream.

In patients using azelaic acid formulations, the following additional adverse experiences have been reported rarely: worsening of asthma, vitiligo depigmentation, small depigmented spots, hypertrichosis, reddening (signs of keratosis pilaris), and exacerbation of recurrent herpes labialis.

## **DOSAGE AND ADMINISTRATION**

After the skin is thoroughly washed and patted dry, a thin film of **AZELEX**<sup>®</sup> cream should be gently but thoroughly massaged into the affected areas twice daily, in the morning and evening. The hands should be washed following application. The duration of use of **AZELEX**<sup>®</sup> cream can vary from person to person and depends on the severity of the acne. Improvement of the condition occurs in the majority of patients with inflammatory lesions within four weeks.

## **HOW SUPPLIED**

**AZELEX**<sup>®</sup> cream is supplied in a closed orifice tube with a white, spiked screwcap in the following sizes:

30 g - NDC 0023-8694-30

50 g - NDC 0023-8694-50

**Storage:** Protect from freezing. Store product on its side. Store at 15°-30° C (59°-86° F).

**Revised: 09/2015**

Distributed under license by Allergan, Inc., Irvine, CA 92612, U.S.A.

© 2015 Allergan. All right reserved.

® marks owned by Allergan, Inc.

Made in Italy by Bayer HealthCare Manufacturing S.r.l., Segrate (Milan), Italy


70896US17

## PRINCIPAL DISPLAY PANEL

NDC 0023-8694-50

AZELEX

(azelaic acid cream) 20%



## AZELEX

azelaic acid cream

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0023-8694
<b>Route of Administration</b>	CUTANEOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

azelaic acid (UNII: F2VW3D43YT) (azelaic acid - UNII:F2VW3D43YT)	azelaic acid	0.2 g in 1 g
--	--------------	--------------

### Inactive Ingredients

Ingredient Name	Strength
cetearyl ethylhexanoate (UNII: 9M64UO4C25)	
glycerin (UNII: PDC6A3C0OX)	
cetostearyl alcohol (UNII: 2DMT128M1S)	
cetyl palmitate (UNII: 5ZA2S6B08X)	
propylene glycol (UNII: 6DC9Q167V3)	
water (UNII: 059QF0K00R)	
glyceryl monostearate (UNII: 230OU9XXE4)	
coco-glycerides (UNII: ISE9I7DNUG)	
benzoic acid (UNII: 8SKN0B0MIM)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0023-8694-30	1 in 1 CARTON	03/21/1996	
1		30 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:0023-8694-50	1 in 1 CARTON	03/21/1996	
2		50 g in 1 TUBE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020428	03/21/1996	

**Labeler** - Allergan, Inc. (144796497)