

KETOROLAC TROMETHAMINE- ketorolac tromethamine solution

SOLA Pharmaceuticals, LLC

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use KETOROLAC TROMETHAMINE OPHTHALMIC SOLUTION safely and effectively. See full prescribing information for KETOROLAC TROMETHAMINE OPHTHALMIC SOLUTION.

KETOROLAC TROMETHAMINE ophthalmic solution, 0.5%

Initial U.S. Approval: 1991

INDICATIONS AND USAGE

Ketorolac tromethamine ophthalmic solution is a nonsteroidal, anti-inflammatory indicated for:

- The treatment of inflammation following cataract surgery. (1)
- The temporary relief of ocular itching due to seasonal allergic conjunctivitis. (1)

DOSAGE AND ADMINISTRATION

One drop of ketorolac tromethamine ophthalmic solution should be applied to the affected eye(s) four times a day for relief of ocular itching due to seasonal allergic conjunctivitis.

For the treatment of postoperative inflammation in patients who have undergone cataract extraction, one drop of ketorolac tromethamine ophthalmic solution should be applied to the affected eye four times daily beginning 24 hours after cataract surgery and continuing through the first 2 weeks of the postoperative period. (2.1)

DOSAGE FORMS AND STRENGTHS

Ophthalmic solution containing 5 mg/mL ketorolac tromethamine. (3)

- 5 mL size bottle filled with 3 mL of solution
- 10 mL size bottle filled with 5 mL of solution
- 10 mL size bottle filled with 10 mL of solution

CONTRAINDICATIONS

Hypersensitivity to any component of this product. (4)

WARNINGS AND PRECAUTIONS

- Delayed healing (5.1)
- Cross-sensitivity or hypersensitivity (5.2)
- Increased bleeding time due to interference with thrombocyte aggregation (5.3)
- Corneal effects including keratitis (5.4)

ADVERSE REACTIONS

The most frequent adverse reactions reported by up to 40% of patients participating in clinical trials have been transient stinging and burning on instillation. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact SOLA Pharmaceuticals at 1-866-747-7365 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 3/2024

FULL PRESCRIBING INFORMATION: CONTENTS*
1 INDICATIONS AND USAGE
2 DOSAGE AND ADMINISTRATION

- 2.1 Recommended Dosing
- 2.2 Use with Other Topical Ophthalmic Medications

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Delayed Healing
- 5.2 Cross-Sensitivity or Hypersensitivity
- 5.3 Increased Bleeding Time
- 5.4 Corneal Effects
- 5.5 Contact Lens Wear

6 ADVERSE REACTIONS

- 6.1 Clinical Studies Experience
- 6.2 Postmarketing Experience

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

- 17.1 Slow or Delayed Healing
- 17.2 Avoiding Contamination of the Product
- 17.3 Contact Lens Wear
- 17.4 Intercurrent Ocular Conditions
- 17.5 Concomitant Topical Ocular Therapy

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Ketorolac tromethamine ophthalmic solution is indicated for the temporary relief of ocular itching due to seasonal allergic conjunctivitis. Ketorolac tromethamine ophthalmic solution is also indicated for the treatment of postoperative inflammation in patients who have undergone cataract extraction.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosing

Patient Dosing

The recommended dose of ketorolac tromethamine ophthalmic solution is one drop four times a day to the affected eye(s) for relief of ocular itching due to seasonal allergic conjunctivitis.

For the treatment of postoperative inflammation in patients who have undergone cataract extraction, one drop of ketorolac tromethamine ophthalmic solution should be applied to the affected eye four times daily beginning 24 hours after cataract surgery and continuing through the first 2 weeks of the postoperative period.

2.2 Use with Other Topical Ophthalmic Medications

Ketorolac tromethamine ophthalmic solution has been safely administered in conjunction with other ophthalmic medications such as antibiotics, alpha-agonists, beta blockers, carbonic anhydrase inhibitors, cycloplegics, and mydriatics. Drops should be administered at least 5 minutes apart.

3 DOSAGE FORMS AND STRENGTHS

5 mL size bottle filled with 3 mL of ketorolac tromethamine ophthalmic solution, 0.5% (5 mg/mL)

10 mL size bottle filled with 5 mL of ketorolac tromethamine ophthalmic solution, 0.5% (5 mg/mL)

10 mL size bottle filled with 10 mL of ketorolac tromethamine ophthalmic solution, 0.5% (5 mg/mL)

4 CONTRAINDICATIONS

Ketorolac tromethamine ophthalmic solution is contraindicated in patients with previously demonstrated hypersensitivity to any of the ingredients in the formulation.

5 WARNINGS AND PRECAUTIONS

5.1 Delayed Healing

Topical nonsteroidal anti-inflammatory drugs (NSAIDs) may slow or delay healing. Topical corticosteroids are also known to slow or delay healing. Concomitant use of topical NSAIDs and topical steroids may increase the potential for healing problems.

5.2 Cross-Sensitivity or Hypersensitivity

There is the potential for cross-sensitivity to acetylsalicylic acid, phenylacetic acid derivatives, and other NSAIDs. There have been reports of bronchospasm or exacerbation of asthma associated with the use of ketorolac tromethamine ophthalmic solution in patients who have either a known hypersensitivity to aspirin/non-steroidal anti-inflammatory drugs or a past medical history of asthma. Therefore, caution should be used when treating individuals who have previously exhibited sensitivities to these drugs.

5.3 Increased Bleeding Time

With some NSAIDs, there exists the potential for increased bleeding time due to interference with thrombocyte aggregation. There have been reports that ocularly applied nonsteroidal anti-inflammatory drugs may cause increased bleeding of ocular tissues (including hyphemas) in conjunction with ocular surgery.

It is recommended that ketorolac tromethamine ophthalmic solution be used with caution in patients with known bleeding tendencies or who are receiving other medications, which may prolong bleeding time.

5.4 Corneal Effects

Use of topical NSAIDs may result in keratitis. In some susceptible patients, continued use of topical NSAIDs may result in epithelial breakdown, corneal thinning, corneal erosion, corneal ulceration, or corneal perforation. These events may be sight threatening. Patients with evidence of corneal epithelial breakdown should immediately discontinue use of topical NSAIDs and should be closely monitored for corneal health.

Postmarketing experience with topical NSAIDs suggests that patients with complicated ocular surgeries, corneal denervation, corneal epithelial defects, diabetes mellitus, ocular surface diseases (e.g., dry eye syndrome), rheumatoid arthritis, or repeat ocular surgeries within a short period of time may be at increased risk for corneal adverse events which may become sight threatening. Topical NSAIDs should be used with caution in these patients.

Postmarketing experience with topical NSAIDs also suggests that use more than 1 day prior to surgery or use beyond 14 days post-surgery may increase patient risk for the occurrence and severity of corneal adverse events.

5.5 Contact Lens Wear

Ketorolac tromethamine ophthalmic solution should not be administered while wearing contact lenses.

6 ADVERSE REACTIONS

6.1 Clinical Studies Experience

Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical studies of a drug cannot be directly compared to the rates in the clinical studies of another drug and may not reflect the rates observed in practice.

The most frequent adverse reactions reported with the use of ketorolac tromethamine ophthalmic solutions have been transient stinging and burning on instillation. These reactions were reported by up to 40% of patients participating in clinical trials.

Other adverse reactions occurring approximately 1 to 10% of the time during treatment with ketorolac tromethamine ophthalmic solutions included allergic reactions, corneal edema, iritis, ocular inflammation, ocular irritation, superficial keratitis, and superficial ocular infections.

Other adverse reactions reported rarely with the use of ketorolac tromethamine

ophthalmic solutions included: corneal infiltrates, corneal ulcer, eye dryness, headaches, and visual disturbance (blurry vision).

6.2 Postmarketing Experience

The following adverse reactions have been identified during post-marketing use of ketorolac tromethamine ophthalmic solution 0.5% in clinical practice. Because they are reported voluntarily from a population of unknown size, estimates of frequency cannot be made. The reactions, which have been chosen for inclusion due to either their seriousness, frequency of reporting, possible causal connection to topical ketorolac tromethamine ophthalmic solution 0.5% or a combination of these factors, include bronchospasm or exacerbation of asthma, corneal erosion, corneal perforation, corneal thinning, and epithelial breakdown [see Warnings and Precautions (5.2, 5.4)].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Teratogenic Effects. Pregnancy Category C

Pregnancy Category C: Ketorolac tromethamine, administered during organogenesis, was not teratogenic in rabbits and rats at oral doses of 3.6 mg/kg/day and 10 mg/kg/day, respectively. These doses are approximately 100 times and 250 times higher respectively than the maximum recommended human topical ophthalmic daily dose of 2 mg (5 mg/ mL x 0.05 mL/drop, x 4 drops x 2 eyes) to affected eyes on a mg/kg basis. Additionally, when administered to rats after Day 17 of gestation at oral doses up to 1.5 mg/kg/day (approximately 40 times the typical human topical ophthalmic daily dose), ketorolac tromethamine resulted in dystocia and increased pup mortality. There are no adequate and well-controlled studies in pregnant women. Ketorolac tromethamine ophthalmic solution should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nonteratogenic Effects:Because of the known effects of prostaglandin-inhibiting drugs on the fetal cardiovascular system (closure of the ductus arteriosus), the use of ketorolac tromethamine ophthalmic solution during late pregnancy should be avoided.

8.3 Nursing Mothers

Because many drugs are excreted in human milk, caution should be exercised when ketorolac tromethamine ophthalmic solution is administered to a nursing woman.

8.4 Pediatric Use


Safety and efficacy in pediatric patients below the age of 2 have not been established.

8.5 Geriatric Use

No overall clinical differences in safety or effectiveness have been observed between elderly and other adult patients.

11 DESCRIPTION

Ketorolac tromethamine ophthalmic solution 0.5% is a member of the pyrrolo-pyrrole group of nonsteroidal anti-inflammatory drugs (NSAIDs) for ophthalmic use. Its chemical name is (±)-5-Benzoyl-2, 3-dihydro-1H pyrrolizine-1-carboxylic acid compound with 2-amino-2-(hydroxymethyl)-1,3-propanediol (1:1) and it has the following structure:



Ketorolac tromethamine ophthalmic solution is supplied as a sterile isotonic aqueous 0.5% solution, with a pH of 7.4. Ketorolac tromethamine ophthalmic solution is a racemic mixture of R-(+) and S-(-) -ketorolac tromethamine. Ketorolac tromethamine may exist in three crystal forms. All forms are equally soluble in water. The pKa of ketorolac is 3.5. This white to off-white crystalline substance discolors on prolonged exposure to light. The molecular weight of ketorolac tromethamine is 376.41. The osmolality of ketorolac tromethamine ophthalmic solution is 290 mOsmol/kg.

Each mL of ketorolac tromethamine ophthalmic solution contains: **Active:**ketorolac tromethamine 0.5%. **Preservative:**benzalkonium chloride 0.01%. **Inactives:**edetate disodium 0.1%; octoxynol 40; water for injection; sodium chloride; hydrochloric acid and/or sodium hydroxide to adjust the pH.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Ketorolac tromethamine is a nonsteroidal anti-inflammatory drug which, when administered systemically, has demonstrated analgesic, anti-inflammatory, and anti-pyretic activity. The mechanism of its action is thought to be due to its ability to inhibit prostaglandin biosynthesis

12.3 Pharmacokinetics

Two drops of 0.5% ketorolac tromethamine ophthalmic solution instilled into the eyes of patients 12 hours and 1 hour prior to cataract extraction achieved a mean ketorolac concentration of 95 ng/mL in the aqueous humor of 8 of 9 eyes tested (range 40 to 170 ng/mL).

One drop of 0.5% ketorolac tromethamine ophthalmic solution was instilled into 1 eye and 1 drop of vehicle into the other eye TID in 26 healthy subjects. Five (5) of 26 subjects had detectable concentrations of ketorolac in their plasma (range 11 to 23 ng/mL) at Day 10 during topical ocular treatment. The range of concentrations following TID dosing of 0.5% ketorolac tromethamine ophthalmic solution are approximately 4 to 8% of the steady state mean minimum plasma concentration observed following four times daily oral administration of 10 mg ketorolac in humans (290 ± 70 ng/mL).

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Ketorolac tromethamine was not carcinogenic in either rats given up to 5 mg/kg/day orally for 24 months or in mice given 2 mg/kg/day orally for 18 months. These doses are approximately 125 times and 50 times higher respectively than the maximum recommended human topical ophthalmic daily dose given as QID for itching to affected eyes on a mg/kg basis.

Ketorolac tromethamine was not mutagenic *in vitro* in the Ames assay or in forward mutation assays. Similarly, it did not result in an *in vitro* increase in unscheduled DNA synthesis or an *in vivo* increase in chromosome breakage in mice. However, ketorolac tromethamine did result in an increased incidence in chromosomal aberrations in Chinese hamster ovary cells. Ketorolac tromethamine did not impair fertility when administered orally to male and female rats at doses up to 9 mg/kg/day and 16 mg/kg/day, respectively. These doses are respectively 225 and 400 times higher than the typical human topical ophthalmic daily dose.

14 CLINICAL STUDIES

Two controlled clinical studies showed that ketorolac tromethamine ophthalmic solution was significantly more effective than its vehicle in relieving ocular itching caused by seasonal allergic conjunctivitis.

Two controlled clinical studies showed that patients treated for two weeks with ketorolac tromethamine ophthalmic solution were less likely to have measurable signs of inflammation (cell and flare) than patients treated with its vehicle.

Results from clinical studies indicate that ketorolac tromethamine has no significant effect upon intraocular pressure; however, changes in intraocular pressure may occur following cataract surgery.

16 HOW SUPPLIED/STORAGE AND HANDLING

Ketorolac tromethamine ophthalmic solution 0.5% is supplied sterile, in white opaque plastic LDPE bottles with white droppers, with gray high density polyethylene (HDPE) caps as follows:

3mL in 5 mL bottle **NDC70512-790-03**

5 mL in 10 mL bottle **NDC70512-790-05**

10 mL in 10 mL bottle **NDC70512-790-10**

Storage: Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature]. Protect from light.

17 PATIENT COUNSELING INFORMATION

17.1 Slow or Delayed Healing

Patients should be informed of the possibility that slow or delayed healing may occur while using nonsteroidal anti-inflammatory drugs (NSAIDs).

17.2 Avoiding Contamination of the Product

Patients should be instructed to avoid allowing the tip of the bottle to contact the eye or surrounding structures because this could cause the tip to become contaminated by common bacteria known to cause ocular infections. Serious damage to the eye and subsequent loss of vision may result from using contaminated solutions.

Also, to avoid the potential for cross-contamination, the patient should be advised to use one bottle for each eye following bilateral ocular surgery. The use of the same bottle of topical eye drops for both eyes following bilateral ocular surgery is not recommended.

17.3 Contact Lens Wear

Patients should be advised that ketorolac tromethamine ophthalmic solution should not be administered while wearing contact lenses.

17.4 Intercurrent Ocular Conditions

Patients should be advised that if they develop an intercurrent ocular condition (e.g., trauma or infection) or have ocular surgery, they should immediately seek their physician's advice concerning the continued use of ketorolac tromethamine ophthalmic solution.

17.5 Concomitant Topical Ocular Therapy

Patients should be advised that if more than one topical ophthalmic medication is being used, the medicines should be administered at least 5 minutes apart.

Manufactured for:

SOLA Pharmaceuticals
Baton Rouge, LA 70810
Made in India

Code: TN/Drugs/TN00003457

Revised: 03/2024

22200826

Principal Display Panel Text for 3mL Container Label:

NDC 70512-790-03

**Ketrolac Tromethamine
Ophthalmic Solution**

0.5%

FOR TOPICAL OPHTHALMIC USE ONLY

Sterile

Rx only 3 mL

Each mL contains: 22200820

Active: ketorolac tromethamine 0.5%.

Preservative: benzalkonium chloride 0.01%

Usual Dosage: One drop four times a day. Refer to package insert.

Storage: Store at 20° - 25°C (68° - 77°F) [See USP Controlled Room Temperature]. **Protect from light.**

Manufactured for:

SOLA Pharmaceuticals
Baton Rouge, LA 70810
Made in India

LOT:

EXP:

Code No.:TN/Drugs/TN00003457

NDC 70512-790-03

**Ketorolac Tromethamine
Ophthalmic Solution**

0.5%

FOR TOPICAL OPHTHALMIC USE ONLY

Rx only **3 mL**

Sterile

Each mL contains: 22200820
Active: ketorolac tromethamine 0.5%.
Preservative: benzalkonium chloride 0.01%.
Usual Dosage: One drop four times a day. Refer to package insert.
Storage: Store at 20° - 25°C (68° - 77°F). [See USP Controlled Room Temperature]. **Protect from light.**
Manufactured for:
SOLA Pharmaceuticals
Baton Rouge, LA 70810
Made in India

3 70512 79003 5
Code No.:TN/Drugs/TN00003457

LOT: **17 x 7 mm**
EXP: **NUVZ**

Principal Display Panel Text for 3mL Carton Label:

NDC 70512-790-03

**Ketorolac
Tromethamine
Ophthalmic
Solution**

0.5%

**FOR TOPICAL
OPHTHALMIC USE ONLY**

Sterile

Rx Only

3mL

Each mL contains:

Active: ketorolac tromethamine 0.5%.

Preservative: benzalkonium chloride 0.01%

Inactives:edetate disodium 0.1%; octoxynol 40; water for injection; sodium chloride; and hydrochloric acid and/or sodium hydroxide to adjust pH to 7.4.

Usual Dosage: One drop four times a day. Refer to package insert.

Storage: Store at 20° - 25°C (68° - 77°F)
[See USP Controlled Room Temperature].

Protect from light.


KEEP OUT OF REACH OF CHILDREN

22200823

Manufactured for:
SOLA Pharmaceuticals
Baton Rouge, LA 70810

Made in India

Code No.:TN/Drugs/TN00003457



Principal Display Panel Text for 5mL Container Label:

NDC 70512-790-05

**Ketorolac Tromethamine
Ophthalmic Solution**

0.5%

FOR TOPICAL OPHTHALMIC USE ONLY

Sterile

Rx only 5 mL

Each mL contains: 22200831

Active: ketorolac tromethamine 0.5%.

Preservative: benzalkonium chloride 0.01%

Usual Dosage: One drop four times a day. Refer to package insert.

Storage: Store at 20° - 25°C (68° - 77°F) [See USP Controlled Room Temperature]. **Protect from light.**

Note: Bottle filled to 1/2 capacity.

Manufactured for:

SOLA Pharmaceuticals
Baton Rouge, LA 70810
Made in India

LOT:

EXP:

Code No.:TN/Drugs/TN00003457

The image shows a principal display panel for a 5mL carton label. It features a vertical blue bar on the left with the text '3 mm unavarnished area for overwrapping'. The main label area is white with orange and blue accents. At the top left, it displays 'NDC 70512-790-05'. Below this, an orange box contains 'Ketorolac Tromethamine Ophthalmic Solution' and a blue box contains '0.5%'. A white box below that says 'FOR TOPICAL OPHTHALMIC USE ONLY'. To the right of this box, the word 'Sterile' is written above an eye icon. Below the eye icon, it says 'Rx only' and '5 mL'. On the right side of the label, there is a section with 'Each mL contains: 22200831', followed by 'Active: ketorolac tromethamine 0.5%', 'Preservative: benzalkonium chloride 0.01%', 'Usual Dosage: One drop four times a day. Refer to package insert.', 'Storage: Store at 20° - 25°C (68° - 77°F). [See USP Controlled Room Temperature]. Protect from light.', 'Note: Bottle filled to 1/2 capacity.', and 'Manufactured for: SOLA Pharmaceuticals, Baton Rouge, LA 70810, Made in India'. To the right of this text is a barcode with the numbers '3 70512 79005 9' and 'Code No.:TN/Drugs/TN00003457' below it. At the bottom right, there is a light blue box containing 'LOT:' and 'EXP:' labels, and a larger light blue box containing '18 x 8.5 mm' and 'NUVZ'.

Principal Display Panel Text for 5mL Carton Label:

NDC 70512-790-05

**Ketorolac
Tromethamine
Ophthalmic
Solution**

0.5%

**FOR TOPICAL
OPHTHALMIC USE ONLY**

Sterile

Rx Only

5 mL

Each mL contains:

Active: ketorolac tromethamine 0.5%.

Preservative: benzalkonium chloride 0.01%

Inactives: edetate disodium 0.1%; octoxynol 40; water for injection; sodium chloride; and hydrochloric acid and/or sodium hydroxide to adjust pH to 7.4.

Usual Dosage: One drop four times a day. Refer to package insert.

Storage: Store at 20° - 25°C (68° - 77°F)
[See USP Controlled Room Temperature].

Protect from light.

KEEP OUT OF REACH OF CHILDREN

Note: Bottle filled to 1/2 capacity.

22200833

Manufactured for:
SOLA Pharmaceuticals
Baton Rouge, LA 70810

Made in India

Code No.: TN/Drugs/TN00003457



Principal Display Panel Text for 10mL Container Label:

NDC 70512-790-10

**Ketorolac Tromethamine
Ophthalmic Solution**

0.5%

FOR TOPICAL OPHTHALMIC USE ONLY

Sterile

Rx only 10 mL

Each mL contains: 2200832

Active: ketorolac tromethamine 0.5%.

Preservative: benzalkonium chloride 0.01%

Usual Dosage: One drop four times a day. Refer to package insert.

Storage: Store at 20° - 25°C (68° - 77°F) [See USP Controlled Room Temperature]. **Protect from light.**

Manufactured for:

SOLA Pharmaceuticals

Baton Rouge, LA 70810

Made in India

LOT:

EXP:

Code No.:TN/Drugs/TN00003457

3 mm unavarnished area for overlapping

NDC 70512-790-10

**Ketorolac Tromethamine
Ophthalmic Solution**

0.5%

FOR TOPICAL OPHTHALMIC USE ONLY

Sterile

Rx only  10 mL

Each mL contains: 22200832
Active: ketorolac tromethamine 0.5%.
Preservative: benzalkonium chloride 0.01%.
Usual Dosage: One drop four times a day. Refer to package insert.
Storage: Store at 20° - 25°C (68° - 77°F). [See USP Controlled Room Temperature]. **Protect from light.**
Manufactured for:
SOLA Pharmaceuticals
Baton Rouge, LA 70810
Made in India


3 70512 79010 3
Code No.:TN/Drugs/TN00003457

LOT: **18 x 8.5 mm
NUVZ**

EXP:

Principal Display Panel Text for 10mL Carton Label:

NDC 70512-790-10

**Ketorolac
Tromethamine
Ophthalmic
Solution**

0.5%

**FOR TOPICAL
OPHTHALMIC USE ONLY**

Sterile

Rx Only

10 mL

Each mL contains:

Active: ketorolac tromethamine 0.5%.

Preservative: benzalkonium chloride 0.01%

Inactives: edetate disodium 0.1%; octoxynol 40; water for injection; sodium chloride; and hydrochloric acid and/or sodium hydroxide to adjust pH to 7.4.

Usual Dosage: One drop four times a day. Refer to package insert.

Storage: Store at 20° - 25°C (68° - 77°F)
[See USP Controlled Room Temperature].

Protect from light.

KEEP OUT OF REACH OF CHILDREN

22200834

Manufactured for:
SOLA Pharmaceuticals
Baton Rouge, LA 70810

Made in India

Code No.:TN/Drugs/TN00003457



KETOROLAC TROMETHAMINE

ketorolac tromethamine solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70512-790
Route of Administration	OPHTHALMIC		

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
KETOROLAC TROMETHAMINE (UNII: 4EVE5946BQ) (KETOROLAC - UNII:YZI5105VOL)		KETOROLAC TROMETHAMINE	5 mg in 1 mL	
Inactive Ingredients				
Ingredient Name			Strength	
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)				
EDETATE DISODIUM (UNII: 7FLD91C86K)				
OCTOXYNOL-40 (UNII: 9T1C662FKS)				
SODIUM CHLORIDE (UNII: 451W47IQ8X)				
HYDROCHLORIC ACID (UNII: QTT17582CB)				
SODIUM HYDROXIDE (UNII: 55X04QC32I)				
WATER (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70512-790-03	1 in 1 CARTON	04/15/2024	
1		3 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		
2	NDC:70512-790-05	1 in 1 CARTON	04/15/2024	
2		5 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		
3	NDC:70512-790-10	1 in 1 CARTON	04/15/2024	
3		10 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA218204	04/15/2024		

Labeler - SOLA Pharmaceuticals, LLC (080121345)

Revised: 12/2025

SOLA Pharmaceuticals, LLC