

# **SOLIFENACIN SUCCINATE- solifenacin succinate tablet, film coated AvKARE**

---

## **HIGHLIGHTS OF PRESCRIBING INFORMATION**

### **Solifenacin Succinate Tablets**

### **Rx Only**

**These highlights do not include all the information needed to use SOLIFENACIN SUCCINATE TABLETS safely and effectively. See full prescribing information for SOLIFENACIN SUCCINATE TABLETS.**

**Initial U.S. Approval: 2004**

#### **INDICATIONS AND USAGE**

Solifenacin Succinate Tablets is a muscarinic antagonist indicated for the treatment of overactive bladder with symptoms of urge urinary incontinence, urgency, and urinary frequency (1)

#### **DOSAGE AND ADMINISTRATION**

- 5 mg tablet taken once daily, and if well tolerated may be increased to 10 mg once daily ( 2.1)
- Do not exceed 5 mg tablet once daily in patients with:
  - severe renal impairment [Creatinine Clearance] (CLcr <30 ml/min) ( 2.2)
  - moderate hepatic impairment (Child-Pugh B) ( 2.3)
  - concomitant use of potent CYP3A4 inhibitors ( 2.4)
- Use of Solifenacin Succinate Tablets is not recommended in patients with severe hepatic impairment (Child-Pugh C) ( 2.3)

#### **DOSAGE FORMS AND STRENGTHS**

Tablets: 5 mg and 10 mg (3) (3)

#### **CONTRAINDICATIONS**

- Urinary retention ( 4, 5.2)
- Gastric retention ( 4, 5.3)
- Uncontrolled narrow-angle glaucoma ( 4, 5.5)
- In patients who have demonstrated hypersensitivity to the drug ( 4, 6.2)

#### **WARNINGS AND PRECAUTIONS**

- Angioedema and anaphylactic reactions: Reports of angioedema of the face, lips and/or larynx, in some cases occurring after the first dose, have been described. Anaphylactic reactions have been reported rarely ( 5.1)
- Urinary Retention: Administer with caution to patients with clinically significant bladder outflow obstruction ( 5.2)
- Gastrointestinal Disorders: Use with caution in patients with decreased gastrointestinal motility ( 5.3)
- Central Nervous System Effects: Somnolence has been reported with Solifenacin. Advise patients not to drive or operate heavy machinery until they know how Solifenacin affects them ( 5.4)
- Controlled Narrow-Angle Glaucoma: Use with caution in patients being treated for narrow-angle glaucoma ( 5.5)
- QT Prolongation: Use with caution in patients with a known history of QT prolongation or patients who are taking medications known to prolong the QT interval ( 5.8)

#### **ADVERSE REACTIONS**

The most common adverse reactions (> 4% and > placebo) were dry mouth, and constipation at both 5 mg and 10 mg doses; and urinary tract infection, and blurred vision at the 10 mg dose ( 6.1)

To report SUSPECTED ADVERSE REACTIONS, contact AvKARE at 1-855-361-3993 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

#### **DRUG INTERACTIONS**

- Inhibitors of CYP3A4 may increase the concentration of Solifenacin ( 7.1).
- Inducers of CYP3A4 may decrease the concentration of Solifenacin ( 7.2).

#### **USE IN SPECIFIC POPULATIONS**

*Pregnancy and Nursing Mothers:* Solifenacin Succinate Tablets should be used during pregnancy only if the

potential benefit for the mother justifies the potential risk to the fetus. Solifenacin Succinate Tablets should not be administered during nursing ( 8.1, 8.3)

*Pediatric Use:* The safety and effectiveness of Solifenacin Succinate Tablets in pediatric patients have not been established ( 8.4)

**See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.**

**Revised: 9/2020**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

### **2 DOSAGE AND ADMINISTRATION**

2.1 Dosing Information

2.2 Dose Adjustment in Patients with Renal Impairment

2.3 Dose Adjustment in Patients with Hepatic Impairment

2.4 Dose Adjustment in Patients Taking CYP3A4 Inhibitors

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

5.1 Angioedema and Anaphylactic Reactions

5.2 Urinary Retention

5.3 Gastrointestinal Disorders

5.4 Central Nervous System Effects

5.5 Controlled Narrow-Angle Glaucoma

5.6 Hepatic Impairment

5.7 Renal Impairment

5.8 Patients with Congenital or Acquired QT Prolongation

### **6 ADVERSE REACTIONS**

6.1 Clinical Trials Experience

6.2 Post-Marketing Experience

### **7 DRUG INTERACTIONS**

7.1 Potent CYP3A4 Inhibitors

7.2 CYP3A4 Inducers

7.3 Drugs Metabolized by Cytochrome P450

7.4 Warfarin

7.5 Oral Contraceptives

7.6 Digoxin

### **8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Labor and Delivery

8.3 Nursing Mothers

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Renal Impairment

8.7 Hepatic Impairment

8.8 Gender

### **10 OVERDOSAGE**

### **11 DESCRIPTION**

### **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

## **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

## **14 CLINICAL STUDIES**

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Solifenacin Succinate Tablets is a muscarinic antagonist indicated for the treatment of overactive bladder with symptoms of urge urinary incontinence, urgency, and urinary frequency.

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Dosing Information**

The recommended dose of Solifenacin Succinate Tablets is 5 mg once daily. If the 5 mg dose is well tolerated, the dose may be increased to 10 mg once daily.

Solifenacin Succinate Tablets should be taken with water and swallowed whole.

Solifenacin Succinate Tablets can be administered with or without food.

#### **2.2 Dose Adjustment in Patients with Renal Impairment**

For patients with severe renal impairment ( $CL_{cr} < 30$  mL/min), a daily dose of Solifenacin Succinate Tablets greater than 5 mg is not recommended [ see *Warnings and Precautions* ( 5.7); *Use in Specific Populations* ( 8.6) ].

#### **2.3 Dose Adjustment in Patients with Hepatic Impairment**

For patients with moderate hepatic impairment (Child-Pugh B), a daily dose of Solifenacin Succinate Tablets greater than 5 mg is not recommended. Use of Solifenacin Succinate Tablets in patients with severe hepatic impairment (Child-Pugh C) is not recommended [ see *Warnings and Precautions* ( 5.6); *Use in Specific Populations* ( 8.7) ].

#### **2.4 Dose Adjustment in Patients Taking CYP3A4 Inhibitors**

When administered with potent CYP3A4 inhibitors such as ketoconazole, a daily dose of Solifenacin Succinate Tablets greater than 5 mg is not recommended [ see *Drug Interactions* ( 7.1) ].

## **3 DOSAGE FORMS AND STRENGTHS**

The 5 mg tablets are yellow, round film-coated tablets, biconvex, debossed with 'AP28' on one side, and plain on the other side

The 10 mg tablets are yellow, round film-coated tablets, biconvex, debossed with 'AP29' on one side, and plain on the other side.

## **4 CONTRAINDICATIONS**

Solifenacin is contraindicated in patients with:

- urinary retention [see *Warnings and Precautions ( 5.2) ]*,
- gastric retention [see *Warnings and Precautions ( 5.3) ]*,
- uncontrolled narrow-angle glaucoma [see *Warnings and Precautions ( 5.5) ]*, and
- in patients who have demonstrated hypersensitivity to the drug [ see *Adverse Reactions ( 6.2) ]*.

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Angioedema and Anaphylactic Reactions**

Angioedema of the face, lips, tongue, and/or larynx have been reported with solifenacin. In some cases angioedema occurred after the first dose. Cases of angioedema have been reported to occur hours after the first dose or after multiple doses. Angioedema associated with upper airway swelling may be life threatening. If involvement of the tongue, hypopharynx, or larynx occurs, solifenacin should be promptly discontinued and appropriate therapy and/or measures necessary to ensure a patent airway should be promptly provided. Anaphylactic reactions have been reported rarely in patients treated with solifenacin succinate. Solifenacin succinate should not be used in patients with a known or suspected hypersensitivity to solifenacin succinate. In patients who develop anaphylactic reactions, solifenacin succinate should be discontinued and appropriate therapy and/or measures should be taken.

### **5.2 Urinary Retention**

Solifenacin, like other anticholinergic drugs, should be administered with caution to patients with clinically significant bladder outflow obstruction because of the risk of urinary retention [ see *Contraindications ( 4) ]*.

### **5.3 Gastrointestinal Disorders**

Solifenacin, like other anticholinergics, should be used with caution in patients with decreased gastrointestinal motility [ see *Contraindications ( 4) ]*.

### **5.4 Central Nervous System Effects**

Solifenacin is associated with anticholinergic central nervous system (CNS) effects [ see *Adverse Reactions ( 6.2) ]*. A variety of CNS anticholinergic effects have been reported, including headache, confusion, hallucinations and somnolence. Patients should be monitored for signs of anticholinergic CNS effects, particularly after beginning treatment or increasing the dose. Advise patients not to drive or operate heavy machinery until they know how Solifenacin Succinate Tablets affects them. If a patient experiences

anticholinergic CNS effects, dose reduction or drug discontinuation should be considered.

## **5.5 Controlled Narrow-Angle Glaucoma**

Solifenacin should be used with caution in patients being treated for narrow-angle glaucoma [ see *Contraindications* ( 4 ) ].

## **5.6 Hepatic Impairment**

Solifenacin should be used with caution in patients with hepatic impairment. Doses of Solifenacin greater than 5 mg are not recommended in patients with moderate hepatic impairment (Child-Pugh B). Solifenacin is not recommended for patients with severe hepatic impairment (Child-Pugh C) [ see *Dosage and Administration* ( 2.3 ) and *Use in Specific Populations* ( 8.7 ) ].

## **5.7 Renal Impairment**

Solifenacin should be used with caution in patients with renal impairment. Doses of Solifenacin greater than 5 mg are not recommended in patients with severe renal impairment ( $CL_{cr} < 30 \text{ mL/min}$ ) [ see *Dosage and Administration* ( 2.2 ) and *Use in Specific Populations* ( 8.6 ) ].

## **5.8 Patients with Congenital or Acquired QT Prolongation**

In a study of the effect of solifenacin on the QT interval in 76 healthy women [ see *Clinical Pharmacology* ( 12.2 ) ] the QT prolonging effect appeared less with solifenacin 10 mg than with 30 mg (three times the maximum recommended dose), and the effect of solifenacin 30 mg did not appear as large as that of the positive control moxifloxacin at its therapeutic dose. This observation should be considered in clinical decisions to prescribe Solifenacin for patients with a known history of QT prolongation or patients who are taking medications known to prolong the QT interval.

# **6 ADVERSE REACTIONS**

## **6.1 Clinical Trials Experience**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Solifenacin has been evaluated for safety in 1811 patients in randomized, placebo-controlled trials. Expected adverse reactions of antimuscarinic agents are dry mouth, constipation, blurred vision (accommodation abnormalities), urinary retention, and dry eyes. The incidence of dry mouth and constipation in patients treated with Solifenacin was higher in the 10 mg compared to the 5 mg dose group.

In the four 12-week double-blind clinical trials, severe fecal impaction, colonic obstruction, and intestinal obstruction were reported in one patient each, all in the Solifenacin 10 mg group. Angioneurotic edema has been reported in one patient taking Solifenacin 5 mg. Compared to 12 weeks of treatment with Solifenacin, the incidence and severity of adverse reactions were similar in patients who remained on drug for up to 12

months.

The most frequent adverse reaction leading to study discontinuation was dry mouth (1.5%). Table 1 lists the rates of identified adverse reactions, derived from all reported adverse events, in randomized, placebo-controlled trials at an incidence greater than placebo and in 1% or more of patients treated with Solifenacin 5 or 10 mg once daily for up to 12 weeks.

**Table 1. Percentages of Patients with Identified Adverse Reactions, Derived from All Adverse Events Exceeding Placebo Rate and Reported by 1% or More Patients for Combined Pivotal Studies**

	Placebo (%)	Solifenacin, 5 mg (%)	Solifenacin, 10 mg (%)
Number of Patients	1216	578	1233
GASTROINTESTINAL DISORDERS			
Dry Mouth	4.2	10.9	27.6
Constipation	2.9	5.4	13.4
Nausea	2.0	1.7	3.3
Dyspepsia	1.0	1.4	3.9
Abdominal Pain Upper	1.0	1.9	1.2
Vomiting NOS	0.9	0.2	1.1
INFECTIONS AND INFESTATIONS			
Urinary Tract Infection NOS	2.8	2.8	4.8
Influenza	1.3	2.2	0.9
Pharyngitis NOS	1.0	0.3	1.1
NERVOUS SYSTEM DISORDERS			
Dizziness	1.8	1.9	1.8
EYE DISORDERS			
Vision Blurred	1.8	3.8	4.8
Dry Eyes NOS	0.6	0.3	1.6
RENAL AND URINARY DISORDERS			
Urinary Retention	0.6	0	1.4
GENERAL DISORDERS AND ADMINISTRATION SITE CONDITIONS			
Edema Lower Limb	0.7	0.3	1.1
Fatigue	1.1	1.0	2.1
PSYCHIATRIC DISORDERS			
Depression NOS	0.8	1.2	0.8
RESPIRATORY, THORACIC AND MEDIASTINAL DISORDERS			
Cough	0.2	0.2	1.1
VASCULAR DISORDERS			
Hypertension NOS	0.6	1.4	0.5

## 6.2 Post-Marketing Experience

Because these spontaneously reported events are from the worldwide postmarketing experience, the frequency of events and the role of solifenacin in their causation cannot be reliably determined.

The following events have been reported in association with solifenacin use in worldwide postmarketing experience:

General: peripheral edema, hypersensitivity reactions, including angioedema with airway obstruction, rash, pruritus, urticaria, and anaphylactic reaction;

Central Nervous: headache, confusion, hallucinations, delirium and somnolence;

Cardiovascular: QT prolongation; Torsade de Pointes, atrial fibrillation, tachycardia, palpitations;

Hepatic: liver disorders mostly characterized by abnormal liver function tests, AST (aspartate aminotransferase), ALT (alanine aminotransferase), GGT (gamma-glutamyl transferase);

Renal: renal impairment;

Metabolism and nutrition disorders: decreased appetite, hyperkalemia;

Dermatologic: exfoliative dermatitis and erythema multiforme;

Eye disorders: glaucoma;

Gastrointestinal disorders: gastroesophageal reflux disease and ileus;

Respiratory, thoracic and mediastinal disorders: dysphonia;

Musculoskeletal and connective tissue disorders: muscular weakness;

To report SUSPECTED ADVERSE REACTIONS contact AvKARE at 1-855-361-3993; email [drugsafety@avkare.com](mailto:drugsafety@avkare.com); or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

## 7 DRUG INTERACTIONS

### 7.1 Potent CYP3A4 Inhibitors

Following the administration of 10 mg of Solifenacin in presence of 400 mg of ketoconazole, a potent inhibitor of CYP3A4, the mean  $C_{max}$  and AUC of solifenacin increased by 1.5 and 2.7-fold, respectively. Therefore, it is recommended not to exceed a 5 mg daily dose of Solifenacin when administered with therapeutic doses of ketoconazole or other potent CYP3A4 inhibitors [ see *Dosage and Administration ( 2.4 )* and *Clinical Pharmacology ( 12.3 )* ]. The effects of weak or moderate CYP3A4 inhibitors were not examined.

### 7.2 CYP3A4 Inducers

There were no in vivo studies conducted to evaluate the effect of CYP3A4 inducers on Solifenacin. In vitro drug metabolism studies have shown that solifenacin is a substrate of CYP3A4. Therefore, inducers of CYP3A4 may decrease the concentration of solifenacin.

### 7.3 Drugs Metabolized by Cytochrome P450

At therapeutic concentrations, solifenacin does not inhibit CYP1A1/2, 2C9, 2C19, 2D6, or 3A4 derived from human liver microsomes.

## **7.4 Warfarin**

Solifenacin has no significant effect on the pharmacokinetics of R-warfarin or S-warfarin [ see *Clinical Pharmacology ( 12.3 )* ].

## **7.5 Oral Contraceptives**

In the presence of solifenacin there are no significant changes in the plasma concentrations of combined oral contraceptives (ethinyl estradiol/levonorgestrel) [ see *Clinical Pharmacology ( 12.3 )* ].

## **7.6 Digoxin**

Solifenacin had no significant effect on the pharmacokinetics of digoxin (0.125 mg/day) in healthy subjects [ see *Clinical Pharmacology ( 12.3 )* ].

# **8 USE IN SPECIFIC POPULATIONS**

## **8.1 Pregnancy**

### **Pregnancy Category C**

There are no adequate and well-controlled studies in pregnant women.

Reproduction studies have been performed in mice, rats and rabbits. After oral administration of <sup>14</sup>C-solifenacin succinate to pregnant mice, drug-related material was shown to cross the placental barrier. No embryotoxicity or teratogenicity was observed in mice treated with 1.2 times (30 mg/kg/day) the expected exposure at the maximum recommended human dose [MRHD] of 10 mg. Administration of solifenacin succinate to pregnant mice at 3.6 times and greater (100 mg/kg/day and greater) the exposure at the MRHD, during the major period of organ development resulted in reduced fetal body weights. Administration of 7.9 times (250 mg/kg/day) the MRHD to pregnant mice resulted in an increased incidence of cleft palate. In utero and lactational exposures to maternal doses of solifenacin succinate of 3.6 times (100 mg/kg/day) the MRHD resulted in reduced peripartum and postnatal survival, reductions in body weight gain, and delayed physical development (eye opening and vaginal patency). An increase in the percentage of male offspring was also observed in litters from offspring exposed to maternal doses of 250 mg/kg/day. No embryotoxic effects were observed in rats at up to 50 mg/kg/day (<1 times the exposure at the MRHD) or in rabbits at up to 1.8 times (50 mg/kg/day) the exposure at the MRHD. Because animal reproduction studies are not always predictive of human response, Solifenacin should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

## **8.2 Labor and Delivery**

The effect of Solifenacin on labor and delivery in humans has not been studied.

There were no effects on natural delivery in mice treated with 1.2 times (30 mg/kg/day) the expected exposure at the maximum recommended human dose [MRHD] of 10 mg. Administration of solifenacin succinate at 3.6 times (100 mg/kg/day) the exposure at the

MRHD or greater increased peripartum pup mortality.

### **8.3 Nursing Mothers**

After oral administration of  $^{14}\text{C}$ -solifenacin succinate to lactating mice, radioactivity was detected in maternal milk. There were no adverse observations in mice treated with 1.2 times (30 mg/kg/day) the expected exposure at the maximum recommended human dose [MRHD]. Pups of female mice treated with 3.6 times (100 mg/kg/day) the exposure at the MRHD or greater revealed reduced body weights, postpartum pup mortality or delays in the onset of reflex and physical development during the lactation period.

It is not known whether solifenacin is excreted in human milk. Because many drugs are excreted in human milk, Solifenacin should not be administered during nursing. A decision should be made whether to discontinue nursing or to discontinue Solifenacin in nursing mothers.

### **8.4 Pediatric Use**

The safety and effectiveness of Solifenacin in pediatric patients have not been established.

### **8.5 Geriatric Use**

In placebo-controlled clinical studies, similar safety and effectiveness were observed between older (623 patients  $\geq$  65 years and 189 patients  $\geq$  75 years) and younger patients (1188 patients  $<$  65 years) treated with Solifenacin.

Multiple dose studies of Solifenacin in elderly volunteers (65 to 80 years) showed that  $\text{C}_{\text{max}}$ ,  $\text{AUC}$  and  $\text{t}_{1/2}$  values were 20-25% higher as compared to the younger volunteers (18 to 55 years).

### **8.6 Renal Impairment**

Solifenacin should be used with caution in patients with renal impairment. There is a 2.1-fold increase in  $\text{AUC}$  and 1.6-fold increase in  $\text{t}_{1/2}$  of solifenacin in patients with severe renal impairment. Doses of Solifenacin greater than 5 mg are not recommended in patients with severe renal impairment ( $\text{CL}_{\text{cr}} < 30 \text{ mL/min}$ ) [ see *Warnings and Precautions* ( 5.7); *Dosage and Administration* ( 2.2 ) ].

### **8.7 Hepatic Impairment**

Solifenacin should be used with caution in patients with reduced hepatic function. There is a 2-fold increase in the  $\text{t}_{1/2}$  and 35% increase in  $\text{AUC}$  of solifenacin in patients with moderate hepatic impairment. Doses of Solifenacin greater than 5 mg are not recommended in patients with moderate hepatic impairment (Child-Pugh B). Solifenacin is not recommended for patients with severe hepatic impairment (Child-Pugh C) [ see *Warnings and Precautions* ( 5.6); *Dosage and Administration* ( 2.3 ) ].

### **8.8 Gender**

The pharmacokinetics of solifenacin is not significantly influenced by gender.

## **10 OVERDOSAGE**


Overdosage with Solifenacin Succinate Tablets can potentially result in severe anticholinergic effects and should be treated accordingly. The highest dose ingested in an accidental overdose of solifenacin succinate was 280 mg in a 5-hour period. This case was associated with mental status changes. Some cases reported a decrease in the level of consciousness.

Intolerable anticholinergic side effects (fixed and dilated pupils, blurred vision, failure of heel-to-toe exam, tremors and dry skin) occurred on day 3 in normal volunteers taking 50 mg daily (5 times the maximum recommended therapeutic dose) and resolved within 7 days following discontinuation of drug.

In the event of overdose with Solifenacin Succinate Tablets, treat with gastric lavage and appropriate supportive measures. ECG monitoring is also recommended.

## 11 DESCRIPTION

Solifenacin Succinate Tablets (solifenacin succinate) is a muscarinic receptor antagonist. Chemically, solifenacin succinate is butanedioic acid, compounded with (1S)-(3R)-1-azabicyclo[2.2.2]oct-3-yl 3,4-dihydro-1-phenyl-2(1H)iso-quinolinecarboxylate (1:1) having an empirical formula of  $C_{23}H_{26}N_2O_2 \cdot C_4H_6O_4$ , and a molecular weight of 480.55. The structural formula of solifenacin succinate is:



Solifenacin succinate is a white to pale-yellowish-white crystal or crystalline powder. It is freely soluble at room temperature in water, glacial acetic acid, dimethyl sulfoxide, and methanol. Each Solifenacin Succinate tablet contains 5 or 10 mg of solifenacin succinate and is formulated for oral administration. In addition to the active ingredient solifenacin succinate, each Solifenacin Succinate tablet also contains the following inert ingredients: lactose monohydrate, pregelatinized starch, hypromellose 2910, magnesium stearate, triacetin, polyethylene glycol 3350 and titanium dioxide with yellow ferric oxide and red ferric oxide.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Solifenacin is a competitive muscarinic receptor antagonist. Muscarinic receptors play an important role in several major cholinergically mediated functions, including contractions of urinary bladder smooth muscle and stimulation of salivary secretion.

### 12.2 Pharmacodynamics

## **Cardiac Electrophysiology**

The effect of 10 mg and 30 mg solifenacin succinate on the QT interval was evaluated at the time of peak plasma concentration of solifenacin in a multi-dose, randomized, double-blind, placebo and positive-controlled (moxifloxacin 400 mg) trial. Subjects were randomized to one of two treatment groups after receiving placebo and moxifloxacin sequentially. One group (n=51) went on to complete 3 additional sequential periods of dosing with solifenacin 10, 20, and 30 mg while the second group (n=25) in parallel completed a sequence of placebo and moxifloxacin. Study subjects were female volunteers aged 19 to 79 years. The 30 mg dose of solifenacin succinate (three times the highest recommended dose) was chosen for use in this study because this dose results in a solifenacin exposure that covers those observed upon co-administration of 10 mg Solifenacin Succinate Tablets with potent CYP3A4 inhibitors (e.g. ketoconazole, 400 mg). Due to the sequential dose escalating nature of the study, baseline EKG measurements were separated from the final QT assessment (of the 30 mg dose level) by 33 days.

The median difference from baseline in heart rate associated with the 10 and 30 mg doses of solifenacin succinate compared to placebo was -2 and 0 beats/minute, respectively. Because a significant period effect on QTc was observed, the QTc effects were analyzed utilizing the parallel placebo control arm rather than the pre-specified intra-patient analysis. Representative results are shown in Table 2.

**Table 2. QTc changes in msec (90%CI) from baseline at Tmax (relative to placebo)\***

<b>Drug/Dose</b>	<b>Fridericia method (using mean difference)</b>
Solifenacin 10 mg	2 (-3,6)
Solifenacin 30 mg	(4,13)

\*Results displayed are those derived from the parallel design portion of the study and represent the comparison of Group 1 to time-matched placebo effects in Group 2

Moxifloxacin was included as a positive control in this study and, given the length of the study, its effect on the QT interval was evaluated in 3 different sessions. The placebo subtracted mean changes (90% CI) in QTcF for moxifloxacin in the three sessions were 11 (7, 14), 12 (8, 17), and 16 (12, 21), respectively.

The QT interval prolonging effect appeared greater for the 30 mg compared to the 10 mg dose of solifenacin. Although the effect of the highest solifenacin dose (three times the maximum therapeutic dose) studied did not appear as large as that of the positive control moxifloxacin at its therapeutic dose, the confidence intervals overlapped. This study was not designed to draw direct statistical conclusions between the drugs or the dose levels.

## **12.3 Pharmacokinetics**

### ***Absorption***

After oral administration of Solifenacin to healthy volunteers, peak plasma levels ( $C_{max}$ ) of solifenacin are reached within 3 to 8 hours after administration, and at steady state ranged from 32.3 to 62.9 ng/mL for the 5 and 10 mg Solifenacin, respectively. The

absolute bioavailability of solifenacin is approximately 90%, and plasma concentrations of solifenacin are proportional to the dose administered.

### ***Effect of food***

Solifenacin may be administered without regard to meals. A single 10 mg dose administration of Solifenacin with food increased  $C_{max}$  and AUC by 4% and 3%, respectively.

### ***Distribution***

Solifenacin is approximately 98% (*in vivo*) bound to human plasma proteins, principally to  $\alpha$ -1-acid glycoprotein. Solifenacin is highly distributed to non-CNS tissues, having a mean steady-state volume of distribution of 600L.

### ***Metabolism***

Solifenacin is extensively metabolized in the liver. The primary pathway for elimination is by way of CYP3A4; however, alternate metabolic pathways exist. The primary metabolic routes of solifenacin are through N-oxidation of the quinuclidin ring and 4R-hydroxylation of tetrahydroisoquinoline ring. One pharmacologically active metabolite (4R-hydroxy solifenacin), occurring at low concentrations and unlikely to contribute significantly to clinical activity, and three pharmacologically inactive metabolites (N-glucuronide and the N-oxide and 4R-hydroxy-N-oxide of solifenacin) have been found in human plasma after oral dosing.

### ***Excretion***

Following the administration of 10 mg of  $^{14}C$ -solifenacin succinate to healthy volunteers, 69.2% of the radioactivity was recovered in the urine and 22.5% in the feces over 26 days. Less than 15% (as mean value) of the dose was recovered in the urine as intact solifenacin. The major metabolites identified in urine were N-oxide of solifenacin, 4R-hydroxy solifenacin and 4R-hydroxy-N-oxide of solifenacin and in feces 4R-hydroxy solifenacin. The elimination half-life of solifenacin following chronic dosing is approximately 45-68 hours.

### ***Drug Interactions***

#### Potent CYP3A4 Inhibitors

In a crossover study, following blockade of CYP3A4 by coadministration of the potent CYP3A4 inhibitor, ketoconazole 400 mg, once daily for 21 days, the mean  $C_{max}$  and AUC of solifenacin increased by 1.5 and 2.7-fold, respectively [ see *Dosage and Administration (2.4)* and *Drug Interactions (7.1)* ].

#### Warfarin

In a crossover study, subjects received a single oral dose of warfarin 25 mg on the 10<sup>th</sup> day of dosing with either solifenacin 10 mg or matching placebo once daily for 16 days. For R-warfarin when it was coadministered with solifenacin, the mean  $C_{max}$  increased by 3% and AUC decreased by 2%. For S-warfarin when it was coadministered with solifenacin, the mean  $C_{max}$  and AUC increased by 5% and 1%, respectively [ see *Drug Interactions (7.4)* ].

#### Oral Contraceptives

In a crossover study, subjects received 2 cycles of 21 days of oral contraceptives

containing 30 ug ethinyl estradiol and 150 ug levonorgestrel. During the second cycle, subjects received additional solifenacin 10 mg or matching placebo once daily for 10 days starting from 12<sup>th</sup> day of receipt of oral contraceptives. For ethinyl estradiol when it was administered with solifenacin, the mean C<sub>max</sub> and AUC increased by 2% and 3%, respectively. For levonorgestrel when it was administered with solifenacin, the mean C<sub>max</sub> and AUC decreased by 1% [ see *Drug Interactions* ( 7.5 ) ].

#### Digoxin

In a crossover study, subjects received digoxin (loading dose of 0.25 mg on day 1, followed by 0.125 mg from days 2 to 8) for 8 days. Consecutively, they received solifenacin 10 mg or matching placebo with digoxin 0.125 mg for additional 10 days. When digoxin was coadministered with solifenacin, the mean C<sub>max</sub> and AUC increased by 13% and 4%, respectively [ see *Drug Interactions* ( 7.6 ) ].

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

No increase in tumors was found following the administration of solifenacin succinate to male and female mice for 104 weeks at doses up to 200 mg/kg/day (5 and 9 times, respectively, of the exposure at the maximum recommended human dose [MRHD] of 10 mg), and male and female rats for 104 weeks at doses up to 20 and 15 mg/kg/day, respectively (<1 times the exposure at the MRHD).

Solifenacin succinate was not mutagenic in the *in vitro* *Salmonella typhimurium* or *Escherichia coli* microbial mutagenicity test or chromosomal aberration test in human peripheral blood lymphocytes with or without metabolic activation, or in the *in vivo* micronucleus test in rats.

Solifenacin succinate had no effect on reproductive function, fertility or early embryonic development of the fetus in male and female mice treated with 250 mg/kg/day (13 times the exposure at the MRHD) of solifenacin succinate, and in male rats treated with 50 mg/kg/day (<1 times the exposure at the MRHD) and female rats treated with 100 mg/kg/day (1.7 times the exposure at the MRHD) of solifenacin succinate.

## **14 CLINICAL STUDIES**

Solifenacin was evaluated in four twelve-week, double-blind, randomized, placebo-controlled, parallel group, multicenter clinical trials for the treatment of overactive bladder in patients having symptoms of urinary frequency, urgency, and/or urge or mixed incontinence (with a predominance of urge). Entry criteria required that patients have symptoms of overactive bladder for  $\geq$  3 months duration. These studies involved 3027 patients (1811 on Solifenacin and 1216 on placebo), and approximately 90% of these patients completed the 12-week studies. Two of the four studies evaluated the 5 and 10 mg Solifenacin doses and the other two evaluated only the 10 mg dose. All patients completing the 12-week studies were eligible to enter an open label, long term extension study and 81% of patients enrolling completed the additional 40-week treatment period. The majority of patients were Caucasian (93%) and female (80%) with a mean age of 58 years.

The primary endpoint in all four trials was the mean change from baseline to 12 weeks in

number of micturitions/24 hours. Secondary endpoints included mean change from baseline to 12 weeks in number of incontinence episodes/24 hours, and mean volume voided per micturition. The efficacy of Solifenacin was similar across patient age and gender. The mean reduction in the number of micturitions per 24 hours was significantly greater with Solifenacin 5 mg (2.3;  $p<0.001$ ) and Solifenacin 10 mg (2.7;  $p<0.001$ ) compared to placebo, (1.4).

The mean reduction in the number of incontinence episodes per 24 hours was significantly greater with Solifenacin 5 mg (1.5;  $p<0.001$ ) and Solifenacin 10 mg (1.8;  $p<0.001$ ) treatment groups compared to placebo (1.1). The mean increase in the volume voided per micturition was significantly greater with Solifenacin 5 mg (32.3 mL;  $p<0.001$ ) and Solifenacin 10 mg (42.5 mL;  $p<0.001$ ) compared with placebo (8.5 mL).

The results for the primary and secondary endpoints in the four individual 12-week clinical studies of Solifenacin are reported in Tables 3 through 6.

**Table 3. Mean Change from Baseline to Endpoint for Solifenacin (5 mg and 10 mg daily) and Placebo: Study 1**

Parameter	Placebo (N=253) Mean (SE)	Solifenacin 5 mg (N=266) Mean (SE)	Solifenacin 10 mg (N=264) Mean (SE)
Urinary Frequency (Number of Micturitions/24 hours)*			
Baseline Reduction	12.2 (0.26)	12.1 (0.24)	12.3 (0.24)
P value vs. placebo	1.2 (0.21)	2.2 (0.18)	2.6 (0.20)
Number of Incontinence Episodes/24 hours†			
Baseline Reduction	2.7 (0.23)	2.6 (0.22)	2.6 (0.23)
P value vs. placebo	0.8 (0.18)	1.4 (0.15)	1.5 (0.18)
Volume Voided per Micturition [mL]†			
Baseline Increase	143.8 (3.37)	149.6 (3.35)	147.2 (3.15)
P value vs. placebo	7.4 (2.28)	32.9 (2.92)	39.2 (3.11)

\*Primary endpoint

†Secondary endpoint

**Table 4. Mean Change from Baseline to Endpoint for Solifenacin (5 mg and 10 mg daily) and Placebo: Study 2**

Parameter	Placebo (N=281) Mean (SE)	Solifenacin 5 mg (N=286) Mean (SE)	Solifenacin 10 mg (N=290) Mean (SE)
Urinary Frequency (Number of Micturitions/24 hours)*			
Baseline	12.3 (0.23)	12.1 (0.23)	12.1 (0.21)

Reduction P value vs. placebo	1.7 (0.19)	2.4 (0.17) < 0.001	2.9 (0.18) < 0.001
Number of Incontinence Episodes/24 hours†			
Baseline Reduction P value vs. placebo	3.2 (0.24) 1.3 (0.19) < 0.01	2.6 (0.18) 1.6 (0.16) 1.6 (0.18)	2.8 (0.20) 0.016
Volume Voided per Micturition [mL]†			
Baseline Increase P value vs. placebo	147.2 (3.18) 11.3 (2.52) < 0.001	148.5 (3.16) 31.8 (2.94) 36.6 (3.04)	145.9 (3.42) < 0.001

\*Primary endpoint  
†Secondary endpoint

**Table 5. Mean Change from Baseline to Endpoint for Solifenacin (10 mg daily) and Placebo: Study 3**

Parameter	Placebo (N=309) Mean (SE)	Solifenacin 10 mg (N=306) Mean (SE)
Urinary Frequency (Number of Micturitions/24 hours)*		
Baseline Reduction P value vs. placebo	11.5 (0.18) 1.5 (0.15) < 0.001	11.7 (0.18) 3.0 (0.15) 3.0 (0.15)
Number of Incontinence Episodes/24 hours†		
Baseline Reduction P value vs. placebo	3.0 (0.20) 1.1 (0.16) < 0.001	3.1 (0.22) 2.0 (0.19) 2.0 (0.19)
Volume Voided per Micturition [mL]†		
Baseline Increase P value vs. placebo	190.3 (5.48) 2.7 (3.15) < 0.001	183.5 (4.97) 47.2 (3.79) 47.2 (3.79)

\*Primary endpoint  
†Secondary endpoint

**Table 6. Mean Change from Baseline to Endpoint for Solifenacin (10 mg daily) and Placebo: Study 4**

Parameter	Placebo (N=295) Mean (SE)	Solifenacin 10 mg (N=298) Mean (SE)
Urinary Frequency (Number of Micturitions/24 hours)*		
Baseline Reduction	11.8 (0.18) 1.3 (0.16)	11.5 (0.18) 2.4 (0.15)

P value vs. placebo		< 0.001
Number of Incontinence Episodes/24 hours†		
Baseline Reduction	2.9 (0.18) 1.2 (0.15)	2.9 (0.17) 2.0 (0.15)
P value vs. placebo		< 0.001
Volume Voided per Micturition [mL]†		
Baseline Increase	175.7 (4.44) 13.0 (3.45)	174.1 (4.15) 46.4 (3.73)
P value vs. placebo		< 0.001

\*Primary endpoint  
†Secondary endpoint

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Solifenacin Succinate Tablets is supplied as round film-coated tablets, biconvex, debossed on the obverse and reverse sides as follows:

	5 mg	10 mg
Obverse	AP28	AP29
Reverse*	--	--
Bottle of 30	NDC 42291-739-30	NDC 42291-740-30
Bottle of 90	NDC 42291-739-90	NDC 42291-740-90

\*Reverse sides are plain for all three strengths.

Store at 25°C (77°F) with excursions permitted from 15°C to 30°C (59°F-86°F) [see USP Controlled Room Temperature].

## 17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (Patient Information).

Patients should be informed that antimuscarinic agents such as Solifenacin Succinate Tablets have been associated with constipation and blurred vision. Patients should be advised to contact their physician if they experience severe abdominal pain or become constipated for 3 or more days. Because Solifenacin Succinate Tablets may cause blurred vision, patients should be advised to exercise caution in decisions to engage in potentially dangerous activities until the drug's effect on the patient's vision has been determined. Heat prostration (due to decreased sweating) can occur when anticholinergic drugs, such as Solifenacin Succinate Tablets, are used in a hot environment. Patients should read the patient leaflet entitled "Patient Information Solifenacin Succinate Tablets" before starting therapy with Solifenacin Succinate Tablets.

Patients should be informed that solifenacin may produce angioedema, which could result in life-threatening airway obstruction. Patients should be advised to promptly discontinue solifenacin therapy and seek immediate attention if they experience edema of the tongue or laryngopharynx, or difficulty breathing.

Manufactured for:

AvKARE

Pulaski, TN 38478

Mfg. Rev. 03/19

AV 09/20

## Patient Information

### **Solifenacin Succinate Tablets**

#### **(so-li-fen-a-cin suc-ci-nate tab-lets)**

Read the Patient Information that comes with Solifenacin Succinate Tablets before you start taking it and each time you get a refill. There may be new information. This leaflet does not take the place of talking with your doctor about your medical condition or treatment.

#### **What is Solifenacin Succinate Tablets?**

Solifenacin Succinate Tablets is a prescription medicine for **adults** used to treat the following symptoms due to a condition called **overactive bladder**:

- Urge urinary incontinence: a strong need to urinate with leaking or wetting accidents
- Urgency: a strong need to urinate right away
- Frequency: urinating often

It is not known if Solifenacin Succinate Tablets is safe and effective in children.

#### **Who should NOT take Solifenacin Succinate Tablets?**

Do not take Solifenacin Succinate Tablets if you:

- are not able to empty your bladder (urinary retention)
- have delayed or slow emptying of your stomach (gastric retention)
- have an eye problem called “uncontrolled narrow-angle glaucoma”
- are allergic to solifenacin succinate or any of the ingredients in Solifenacin Succinate Tablets. See the end of this leaflet for a complete list of ingredients.

#### **What should I tell my doctor before taking Solifenacin Succinate Tablets?**

Before you take Solifenacin Succinate Tablets, tell your doctor if you:

- have any stomach or intestinal problems or problems with constipation
- have trouble emptying your bladder or you have a weak urine stream
- have an eye problem called “narrow angle glaucoma”
- have liver problems
- have kidney problems
- have a rare heart problem called “QT prolongation”
- are pregnant or plan to become pregnant. It is not known if Solifenacin Succinate Tablets will harm your unborn baby. Talk to your doctor if you are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if Solifenacin Succinate Tablets passes into your breast milk. You and your doctor should decide if you will take Solifenacin Succinate Tablets or breastfeed. You should not do both.

**Tell your doctor about all the medicines you take**, including prescription and nonprescription medicines, vitamins, and herbal supplements. Solifenacin Succinate Tablets may affect the way other medicines work, and other medicines may affect how Solifenacin Succinate Tablets works.

### **How should I take Solifenacin Succinate Tablets?**

- Take Solifenacin Succinate Tablets exactly as your doctor tells you to take it.
- You should take 1 Solifenacin Succinate Tablets tablet 1 time a day.
- You should take Solifenacin Succinate Tablets with water and swallow the tablet whole.
- You can take Solifenacin Succinate Tablets with or without food.
- If you miss a dose of Solifenacin Succinate Tablets, begin taking Solifenacin Succinate Tablets again the next day. Do not take 2 doses of Solifenacin Succinate Tablets the same day.
- If you take too much Solifenacin Succinate Tablets, call your doctor or go to the nearest hospital emergency room right away.

### **What should I avoid while taking Solifenacin Succinate Tablets?**

Solifenacin Succinate Tablets can cause blurred vision or drowsiness. Do not drive or operate heavy machinery until you know how Solifenacin Succinate Tablets affects you.

### **What are the possible side effects of Solifenacin Succinate Tablets?**

Solifenacin Succinate Tablets may cause serious side effects including:

- Serious allergic reaction. Stop taking Solifenacin Succinate Tablets and get medical help right away if you have:

- hives, skin rash or swelling
- severe itching
- swelling of your face, mouth or tongue
- trouble breathing

The most common side effects of Solifenacin Succinate Tablets include:

- dry mouth
- constipation. Call your doctor if you get severe stomach area (abdominal) pain or become constipated for 3 or more days.
- urinary tract infection
- blurred vision
- heat exhaustion or heat-stroke. This can happen when Solifenacin Succinate Tablets is used in hot environments. Symptoms may include:

- decreased sweating
- dizziness
- tiredness
- nausea
- increase in body temperature

Tell your doctor if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of Solifenacin Succinate Tablets. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store Solifenacin Succinate Tablets?**

- Store Solifenacin Succinate Tablets at 59°F to 86°F (15°C to 30°C). Keep the bottle closed.
- Safely throw away medicine that is out of date or that you no longer need.

**Keep Solifenacin Succinate Tablets and all medicines out of the reach of children.**

**General information about Solifenacin Succinate Tablets.**

Medicines are sometimes prescribed for purposes other than those listed in Patient Information leaflets. Do not use Solifenacin Succinate Tablets for a condition for which it was not prescribed. Do not give Solifenacin Succinate Tablets to other people, even if they have the same symptoms you have. It may harm them.

This leaflet summarizes the most important information about Solifenacin Succinate Tablets. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about Solifenacin Succinate Tablets that is written for health professionals.

For more information, contact AvKARE at 1-855-361-3993.

**What are the ingredients in Solifenacin Succinate Tablets?**

**Active ingredient:** solifenacin succinate

**Inactive ingredients:** lactose monohydrate, pregelatinized starch, hypromellose 2910, magnesium stearate, triacetin, polyethylene glycol 3350 and titanium dioxide with yellow ferric oxide and red ferric oxide.

**What is overactive bladder?**

Overactive bladder occurs when you cannot control your bladder contractions. When these muscle contractions happen too often or cannot be controlled you can get symptoms of overactive bladder, which are urinary frequency, urinary urgency, and urinary incontinence (leakage).

Rx Only

Manufactured for:

AvKARE

Pulaski, TN 38478

Mfg. Rev. 03/19

AV 09/20



## Solifenacin Succinate Tablets

ONCE-DAILY

5 mg

30 Tablets

Rx Only

Each tablet contains 5 mg of solifenacin succinate.  
DOSAGE: ONCE-DAILY  
See Complete Prescribing Information.  
Store at 25°C (77°F); excursions permitted to 15°C-30°C (59°F - 86°F) (See USP Controlled Room Temperature)  
Pharmacist: Dispense in a tight container as defined in USP.

Important: Use safety closures when dispensing the product unless otherwise directed by physician or requested by purchaser.  
KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

Manufactured for:  
**AVKARE**  
Pulaski, TN 38478  
Mfg. Rev. 03/19 AV 09/20 (P)

For patient information,  
contact **AVKARE**  
at 1-855-361-3993  
Mfg. Rev. 03/19 AV 09/20 (P)

**N**  
**3 42291 73930 7**



## Solifenacin Succinate Tablets

ONCE-DAILY

5 mg

90 Tablets

Rx Only

Each tablet contains 5 mg of solifenacin succinate.  
DOSAGE: ONCE-DAILY  
See Complete Prescribing Information.  
Store at 25°C (77°F); excursions permitted to 15°C-30°C (59°F - 86°F) (See USP Controlled Room Temperature)  
Pharmacist: Dispense in a tight container as defined in USP.

Important: Use safety closures when dispensing the product unless otherwise directed by physician or requested by purchaser.  
KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

Manufactured for:  
**AVKARE**  
Pulaski, TN 38478  
Mfg. Rev. 03/19 AV 09/20 (P)

For patient information,  
contact **AVKARE**  
at 1-855-361-3993  
Mfg. Rev. 03/19 AV 09/20 (P)

**N**  
**3 42291 73990 1**



## Solifenacin Succinate Tablets

ONCE-DAILY

10 mg

30 Tablets

Rx Only

Each tablet contains 10 mg of solifenacin succinate.  
DOSAGE: ONCE-DAILY  
See Complete Prescribing Information.  
Store at 25°C (77°F); excursions permitted to 15°C-30°C (59°F - 86°F) (See USP Controlled Room Temperature)  
Pharmacist: Dispense in a tight container as defined in USP.

Important: Use safety closures when dispensing the product unless otherwise directed by physician or requested by purchaser.  
KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

Manufactured for:  
**AVKARE**  
Pulaski, TN 38478  
Mfg. Rev. 03/19 AV 09/20 (P)

For patient information,  
contact **AVKARE**  
at 1-855-361-3993  
Mfg. Rev. 03/19 AV 09/20 (P)

**N**  
**3 42291 74030 3**





# Solifenacin Succinate Tablets

ONCE-DAILY

10 mg

90 Tablets

Rx Only

Each tablet contains 10 mg of solifenacin succinate.  
DOSE: ONCE-DAILY  
See Complete Prescribing Information.  
Store at 25°C (77°F); excursions permitted to  
15°C-30°C (59°F - 86°F) (See USP Controlled  
Room Temperature)

Pharmacist: Dispense in a tight container as  
defined in USP.

Important: Use safety closures when dispensing  
the product unless otherwise directed by physician  
or requested by purchaser.  
KEEP THIS AND ALL MEDICATION OUT OF THE  
REACH OF CHILDREN.

Manufactured for:  
**AVKARE**  
Pulaski, TN 38478  
Mfg. Rev. 03/19

For patient information,  
contact AVKARE  
at 1-855-361-3993  
AV 09/20 (P)



N 3

4229174090 7

## SOLIFENACIN SUCCINATE

solifenacin succinate tablet, film coated

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:42291-739(NDC:35561-285)
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SOLIFENACIN SUCCINATE (UNII: KKA5DLD701) (SOLIFENACIN - UNII:A8910SQJ1U)	SOLIFENACIN SUCCINATE	5 mg

### Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	AP28
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42291-739-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2020	
2	NDC:42291-739-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2020	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210281	10/01/2020	

# SOLIFENACIN SUCCINATE

solifenacin succinate tablet, film coated

## Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:42291-740(NDC:35561-286)
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SOLIFENACIN SUCCINATE (UNII: KKA5DLD701) (SOLIFENACIN - UNII:A8910SQJ1U)	SOLIFENACIN SUCCINATE	10 mg

## Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	AP29
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42291-740-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2020	
2	NDC:42291-740-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2020	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210281	10/01/2020	

**Labeler** - AvKARE (796560394)