ASPIRIN- aspirin tablet Preferred Pharmaceuticals Inc.

Geri Care 44-249

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves:

- pain and fever of colds
- headache
- muscle pain
- minor pain of arthritis
- toothache
- menstrual pain

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems

take a blood thinning (anticoagulant) or steroid drug

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you have asthma
- · stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - o have bloody or black stools
 - o feel faint
 - o have stomach pain that does not get better
 - o vomit blood
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredient

corn starch

Questions or comments?

1-800-540-3765

Principal Display Panel

NDC 68788-8256

GERICARE

ASPIRIN

(NSAID) • 325 mg each

PAIN RELIEVER / FEVER REDUCER

COMPARE TO THE ACTIVE INGREDIENT OF BAYER® ASPIRIN**

1000 TABLETS

Actual Size

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

50844

REV0122D24912

**This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Aspirin

DISTRIBUTED BY:

GERI-CARE PHARMACEUTICAL CORP.

1650 63 Street, Brooklyn, NY 11204

Repackaged By: Preferred Pharmaceuticals Inc.

Aspirin Tablets 325mg

Generic for Bayer® Aspirin Each tablet contains: Aspirin 325mg (NSAID) ... pain reliever/fever reducer

Pkg Size: Exp Date:

Lot#: Batch#:


Ins:

Mfg: Geri-Care; Brooklyn, New York

Prod#:

Prod#:

Warning
Kep out of the reach of hitera. Ro's syndrome:
children and teenages who hitera. Ro's syndrome:
children and teenages who hitera for recovering from
chicken pax or flu-like symptoms should not use this
product. When using this product, it changes in
behavior with nausea and vomiting occur, consult a
behavior with nausea and vomiting occur, consult a
Reye's syndrome, a raire but serious illness. Alfery wiert
apprint may cause a severe allergie reaction. Nomich
apprint may cause a severe allergie reaction. Nomich
which may cause severe stomach bleeding. Do not use if
you are allergie to asprint or other pain relievers/fever
reducers. Store af 25 (2, 17/F); exclusions permitted
imprinted with ASPIRIN. 44.749



ASPIRIN

aspirin tablet

D		I C	
Prod	LICT	Inform	nation

Product Type HUMAN OTC DRUG **Item Code (Source)** NDC:68788-8256(NDC:57896-901)

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) **ASPIRIN** 325 mg

Inactive Ingredients

Ingredient Name Strength

STARCH, CORN (UNII: O8232NY3SJ)

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;249;ASPIRIN
Contains			

P	2		7	20	III	g
		•				м
		_		=	,===	

Ш	rackagilig				
	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
	NDC:68788- 8256-2	20 in 1 BOTTLE; Type 0: Not a Combination Product	09/16/2022		

	NDC:68788- 8256-1	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/16/2022		
Marketing Information					
	arketing .	mormation			
	Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
	Marketing	Application Number or Monograph Citation	_	_	

Labeler - Preferred Pharmaceuticals Inc. (791119022)

Registrant - Preferred Pharmaceuticals Inc. (791119022)

Establishment					
Name	Address	ID/FEI	Business Operations		
Preferred Pharmaceuticals Inc.		791119022	REPACK(68788-8256)		

Revised: 1/2025 Preferred Pharmaceuticals Inc.