

TOBRAMYCIN- tobramycin solution/ drops
NuCare Pharmaceuticals, Inc.

Rx Only

For Topical Ophthalmic Use Only


DESCRIPTION

Tobramycin ophthalmic solution is a sterile topical ophthalmic antibiotic formulation prepared specifically for topical therapy of external ophthalmic infections.

Each mL of Tobramycin ophthalmic solution USP, 0.3% contains: **Active:** tobramycin 0.3% (3 mg). **Preservative:** Benzalkonium chloride 0.01% (0.1 mg). **Inactives:** boric acid, sodium sulfate, sodium chloride, tyloxapol, sodium hydroxide and/or sulfuric acid (to adjust pH) and water for injection. Tobramycin ophthalmic solution has a pH range between 7.0 and 8.0 and an osmolality of 260-320 mOsm/kg.

Tobramycin is a water-soluble aminoglycoside antibiotic active against a wide variety of gram-negative and gram-positive ophthalmic pathogens.

The chemical structure of tobramycin is:



Molecular Weight = 467.52

Molecular Formula: C₁₈H₃₇N₅O₉

Chemical name: 0-{3-amino-3-deoxy- α -D-glucopyranosyl-(1 \rightarrow 4)}-0-{2,6-diamino-2,3,6-trideoxy- α -D-ribohexopyranosyl-(1 \rightarrow 6)}-2-deoxystreptamine.

CLINICAL PHARMACOLOGY

In Vitro Data: *In vitro* studies have demonstrated tobramycin is active against susceptible strains of the following microorganisms: *Staphylococci*, including *S. aureus* and *S. epidermidis* (coagulase-positive and coagulase-negative), including penicillin-resistant strains.

Streptococci, including some of the Group A-beta-hemolytic species, some nonhemolytic species, and some *Streptococcus pneumoniae*.

Pseudomonas aeruginosa, *Escherichia coli*, *Klebsiella pneumoniae*, *Enterobacter aerogenes*, *Proteus mirabilis*, *Morganella morganii*, most *Proteus vulgaris* strains, *Haemophilus influenzae* and *H. aegyptius*, *Moraxella lacunata*, *Acinetobacter calcoaceticus* and some *Neisseria* species. Bacterial susceptibility studies demonstrate that in some cases, microorganisms resistant to gentamicin retain susceptibility to tobramycin.

INDICATIONS AND USAGE

Tobramycin ophthalmic solution is a topical antibiotic indicated in the treatment of external infections of the eye and its adnexa caused by susceptible bacteria. Appropriate monitoring of bacterial response to topical antibiotic therapy should accompany the use of Tobramycin Ophthalmic Solution. Clinical studies have shown tobramycin to be safe and effective for use in children.

CONTRAINDICATIONS

Tobramycin ophthalmic solution is contraindicated in patients with known hyper sensitivity to any of its components.

WARNINGS

FOR TOPICAL OPHTHALMIC USE ONLY. NOT FOR INJECTION INTO THE EYE. Sensitivity to topically applied aminoglycosides may occur in some patients. If a sensitivity reaction to Tobramycin ophthalmic solution occurs, discontinue use.

PRECAUTIONS

General

As with other antibiotic preparations, prolonged use may result in overgrowth of nonsusceptible organisms, including fungi. If superinfection occurs, appropriate therapy should be initiated. Cross-sensitivity to other aminoglycoside antibiotics may occur; if hypersensitivity develops with this product, discontinue use and institute appropriate

therapy. Patients should be advised not to wear contact lenses if they have signs and symptoms of bacterial ocular infection.

Information For Patients

Do not touch dropper tip to any surface, as this may contaminate the solution.

Pregnancy Category B

Reproduction studies in three types of animals at doses up to thirty-three times the normal human systemic dose have revealed no evidence of impaired fertility or harm to the fetus due to tobramycin. There are, however, no adequate and well-controlled studies in pregnant women. Because animal studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers

Because of the potential for adverse reactions in nursing infants from Tobramycin ophthalmic solution, a decision should be made whether to discontinue nursing the infant or discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

Safety and effectiveness in pediatric patients below the age of 2 months has not been established.

Geriatric Use

No overall differences in safety or effectiveness have been observed between elderly and younger patients.

ADVERSE REACTIONS

The most frequent adverse reactions to Tobramycin ophthalmic solution are hypersensitivity and localized ocular toxicity, including lid itching and swelling, and conjunctival erythema. These reactions occur in less than three of 100 patients treated with Tobramycin ophthalmic solution. Similar reactions may occur with the topical use of other aminoglycoside antibiotics. Other adverse reactions have not been reported from Tobramycin ophthalmic solution therapy; however, if topical ocular tobramycin is administered concomitantly with systemic aminoglycoside antibiotics, care should be taken to monitor the total serum concentration.

OVERDOSAGE

Clinically apparent signs and symptoms of an overdose of Tobramycin ophthalmic solution (punctate keratitis, erythema, increased lacrimation, edema and lid itching) may be similar to adverse reaction effects seen in some patients.

DOSAGE AND ADMINISTRATION

In mild to moderate disease, instill one or two drops into the affected eye(s) every four hours. In severe infections, instill two drops into the eye(s) hourly until improvement, following which treatment should be reduced prior to discontinuation.

HOW SUPPLIED

5 mL sterile solution is packaged in a 10 mL white LDPE bottle and natural LDPE nozzle and White HDPE cap (NDC 68071-2272-5) containing Tobramycin ophthalmic solution.

TAMPER EVIDENT SEAL

Storage: Store at 2° - 25°C (36° - 77°F).

For Product Inquiry call +1-800-417-9175

Revised: October, 2019

Manufactured for:

Somerset Therapeutics, LLC

Hollywood, FL 33024

Made in India

Code No.: KR/DRUGS/KTK/28/289/97

ST-TBR11/P/03

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NuCare Pharmaceuticals, Inc.

NDC: 68071-2272-5

Tobramycin 0.3%
5mL Ophth. Soln.

See manufacturer's label for full list of ingredients.

Product #: R0343005
Rx Only

Tobramycin 0.3%
Lot: 000000 NDC: 68071-2272-05
MFR NDC: 70069-131-01 Exp.: 00-00
Serial# 0000000002

Tobramycin 0.3%
Lot: 000000 NDC: 68071-2272-05
MFR NDC: 70069-131-01 Exp.: 00-00
Serial# 0000000002

GTIN 00368071227253
Serial# 00000000002
Exp. Date 00-00
LOT#: 000000

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Manufactured for: Somerset Therapeutics, LLC
Hollywood, FL 33024
Packaged By: NuCare Pharmaceuticals, Inc.
Orange, CA 92867

Patent Instructions:
Place _____ drop(s) into the affected eye(s) every _____ hours.

Rev. 01/01/19

WARNING: KEEP OUT OF REACH OF CHILDREN

STORE AT CONTROLLED TEMPERATURE 36-77°F.

TOBRAMYCIN

tobramycin solution/ drops

Product Information

Product Type	HUMAN PRESCRIPTION	Item Code	NDC:68071-2272(NDC:70069-
--------------	--------------------	-----------	---------------------------

Product Type	DRUG	(Source)	131)	
Route of Administration	OPHTHALMIC			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	TOBRAMYCIN (UNII: VZ8RRZ51VK) (TOBRAMYCIN - UNII:VZ8RRZ51VK)	TOBRAMYCIN	3 mg in 1 mL	
Inactive Ingredients				
	Ingredient Name	Strength		
	BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)			
	BORIC ACID (UNII: R57ZHV85D4)			
	SODIUM SULFATE ANHYDROUS (UNII: 36KCS0R750)			
	TYLOXAPOL (UNII: Y27PUL9H56)			
	SODIUM CHLORIDE (UNII: 451W47IQ8X)			
	SODIUM HYDROXIDE (UNII: 55X04QC32I)			
	SULFURIC ACID (UNII: O40UQP6WCF)			
	WATER (UNII: 059QF0KO0R)			
Product Characteristics				
Color	white (Clear, colorless solution)	Score		
Shape		Size		
Flavor		Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68071-2272-5	1 in 1 CARTON	10/07/2020	
1		5 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA207444	06/28/2017		

Labeler - NuCare Pharmaceuticals, Inc. (010632300)

Establishment

Name	Address	ID/FEI	Business Operations
NuCare Pharmaceuticals, Inc.		010632300	relabel(68071-2272)

Revised: 7/2024

NuCare Pharmaceuticals, Inc.