

ACID REDUCER ORIGINAL STRENGTH- famotidine tablet, film coated
Rite Aid Corporation

Rite Aid Corporation Acid Reducer Drug Facts

Active ingredient (in each tablet)

Famotidine 10 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients

carnauba wax, hypromellose, iron oxides, magnesium stearate, microcrystalline cellulose, pregelatinized starch, polydextrose, polyethylene glycol, talc, titanium dioxide, triacetin

Questions or comments?

1-800-719-9260

Principal Display Panel

SEE NEW WARNINGS

Compare to the active ingredient of Pepcid® AC Original Strength

ACID REDUCER

Famotidine Tablets, 10 mg

acid reducer

ORIGINAL STRENGTH

Just one tablet prevents & relieves heartburn due to acid indigestion

ACTUAL SIZE

30 TABLETS

Drug Facts (continued)

Active ingredient (in each tablet) Purpose
Famotidine 10 mg.....Acid reducer

Uses
■ relieves heartburn associated with acid indigestion and sour stomach
■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use
■ if you have trouble swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
■ with other acid reducers

Ask a doctor before use if you have
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ heartburn with lightheadedness, sweating, or dizziness
■ chest pain or shoulder pain with shortness of breath, sweating, or pain radiating to arms, neck, or shoulders
■ frequent chest pain or lightheadedness

■ stomach pain ■ kidney disease
■ nausea or vomiting
■ unexplained weight loss
■ frequent wheezing, particularly with heartburn

Other information
■ read the directions and warnings before use
■ keep the carton in a cool, dry place
■ store at 20°-25°C (68°-77°F) ■ protect from moisture

Inactive ingredients camouba wax, hypromellose, croscarmellose, polyethylene glycol, pregelatinized starch, talc, titanium dioxide, triacetin

Directions
■ adults and children 12 years and over:
■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn
■ do not use more than 2 tablets in 24 hours
■ children under 12 years: ask a doctor

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

If pregnant or breast-feeding, ask a health professional before use.
■ you need to take this product for more than 14 days
■ your heartburn continues or worsens
■ stop use and ask a doctor if
■ you have heartburn more often than 14 days
■ you need to take this product for more than 14 days

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

SATISFACTION GUARANTEE

DISTRIBUTED BY: RITE AID
30 HUNTER LANE, CAMP HILL, PA 17011
www.riteaid.com

141 66 83 C3

Code Area

SEE NEW WARNINGS
NDC 11822-0141-3

Compare to the active ingredient of **Pepcid® AC Original Strength***

ACID REDUCER

Famotidine Tablets, 10 mg
acid reducer
ORIGINAL STRENGTH

Just one tablet prevents & relieves heartburn due to acid indigestion

ACTUAL SIZE



30 TABLETS

Tips for Managing Heartburn • Do not lie flat or bend over after eating • Do not wear tight-fitting clothing around the stomach • Do not eat before bedtime • Raise the head of your bed • Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables • Eat slowly and avoid big meals • If overweight, lose weight • Quit smoking

GLUTEN FREE

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Pepcid® AC Original Strength.

0 11822 45358 5



ACID REDUCER ORIGINAL STRENGTH

famotidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-0141
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	10 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

Product Characteristics

Color	PINK	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	L141
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0141-3	30 in 1 CARTON	09/10/2009	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11822-0141-4	40 in 1 CARTON	09/10/2009	10/16/2012

2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:11822-0141-1	2 in 1 CARTON	09/10/2009	
3		90 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075400	09/10/2009	

Labeler - Rite Aid Corporation (014578892)

Revised: 7/2021

Rite Aid Corporation